

Applications of Free Fields in 2D Current Algebra

$$\Sigma \quad \begin{array}{c} \diagup \quad \diagdown \\ \hline \diagdown \quad \diagup \end{array} \quad = \quad \begin{array}{c} \diagdown \quad \diagup \\ \hline \diagup \quad \diagdown \end{array} \quad \Sigma$$

Jørgen Rasmussen

Thesis submitted for the degree of Doctor of Philosophy
in the Niels Bohr Institute at the University of Copenhagen,
October 1996

Contents

1	Introduction	1
2	Conformal Field Theory	6
2.1	Some Basic Concepts	6
2.1.1	Conformal Transformations	6
2.1.2	Fields and Operator Product Expansions	8
2.1.3	Virasoro Algebra	10
2.1.4	Descendant Fields	12
2.1.5	States	13
2.1.6	Singular States and the Kac Determinant	13
2.1.7	Conformal Blocks and Greens Functions	14
2.2	Free Fields	16
2.2.1	A Digression on Lie Algebras	16
2.2.2	Bosonic Scalar Field	17
2.2.3	Ghost Fields	19
2.2.4	Fractional Calculus and Expansions	21
2.2.5	Consistency Check	24
2.3	Minimal Models	26
2.3.1	Dotsenko-Fateev Construction	27
2.3.2	A Hidden Duality	29
2.3.3	An Integral Identity	30
2.4	Affine Current Algebras	33
2.4.1	Fields and States	35
2.4.2	The Knizhnik-Zamolodchikov Equations	36
2.4.3	WZNW Models	37
3	Free Field Realizations	39
3.1	Differential Operator Realization	40
3.1.1	Differential Screening Operators	45
3.1.2	Classical Polynomial Identities	46
3.2	Wakimoto Free Field Realization	46
3.2.1	Quantum Polynomial Identities	48
3.3	Screening Currents	48
3.3.1	Screening Currents of the First Kind	49
3.3.2	Screening Currents of the Second Kind	49

3.4	Primary Fields	51
3.4.1	Case of $SL(3)$	53
3.4.2	Case of $SL(r + 1 \geq 4)$	53
3.5	A Quantum Group Structure	58
3.5.1	Hopf Algebra	58
3.5.2	Screening Current Realization	59
4	Chiral Blocks in $SL(2)$ Current Algebra	64
4.1	Dual States	66
4.1.1	Admissible Representations	69
4.2	3-point Functions	70
4.3	Fusion Rules	74
4.3.1	Decoupling of Singular States	74
4.3.2	Standard and Over-screening of 3-point Functions	76
4.3.3	Over-screening in Minimal Models	79
4.4	N -point Functions	80
4.4.1	Different Realizations	83
4.4.2	The Knizhnik-Zamolodchikov Equations	86
4.4.3	Projective Invariances	89
4.4.4	Possible Hybrid	91
4.5	Hamiltonian Reduction	92
4.5.1	At the Level of Correlators	93
4.5.2	Analysis of Explicit Correlators	97
4.5.3	Comparison with Standard Formulations of Hamiltonian Reduction . . .	100
5	4-point Greens Functions in $SL(2)$ Current Algebra	103
5.1	4-point Functions	104
5.1.1	Integration Contours	107
5.2	Andreev's Representation	114
5.2.1	Case of Fusion Rule I	115
5.2.2	Case of Fusion Rule II	116
5.3	Proof of Equivalence	118
5.4	Over-screening	128
5.5	Crossing Matrix	131
5.5.1	The Column of the Transformation Matrix	131
5.5.2	The Row of the Transformation Matrix	135
5.6	Monodromy Invariant Greens Functions	136
5.6.1	Case of Mixed Fusion Rules	139
5.7	Operator Algebra Coefficients	139
5.7.1	Case of Fusion Rule I	141
5.7.2	Case of Fusion Rule II	144
6	Conclusions and Outlook	148
	Bibliography	149

Chapter 1

Introduction

In recent years great efforts have been spent in the quest for a theory of *quantum gravity*, which probably must be a unifying theory of all known interactions. Based on the principles of quantum mechanics and local gauge invariance, the standard model is a widely recognised and experimentally tested quantum field theory of electro-weak and strong interactions. Einstein's theory of general relativity which is based on general covariance and the equivalence principle, is likewise an accepted and tested classical field theory of gravity. Though of dominant importance for large-scale phenomena due to its long-range attractive property, gravity is a weak force and negligible in particle scattering experiments done in order to test the standard model. However, in sufficiently strong gravitational fields as occurring near black hole singularities and in current descriptions of the early universe, gravity must be taken into account in a revision of the standard model. Attempts to formulate gravity as a quantum field theory have failed because of the non-renormalisability of gravity, which stems from the fact that the coupling constant (Newton's constant) has dimension as the Planck length squared in units where $\hbar = c = 1$, leading to a breakdown of standard perturbative approach.

A prime candidate for a theory of quantum gravity is (super-)string theory, or for brevity *strings*. The terminology originates from the basic description of elementary particles as excitations of one dimensional objects, strings, of the size of the Planck length. There is definitely a need for strings to be tiny since there is no experimental evidence for stringy extendedness. An intriguing observation is that the graviton, the elementary quantum of the gravitational field, is inherent in string theory as one of the excitations. Furthermore, the stringy nature introduces a natural cut-off rendering problems with short-distance divergences tractable. Among the references to strings are [116, 60, 77, 72, 1]. Very recent developments on dualities in string theory and p -branes (higher dimensional objects than strings) [114, 137, 131] tend to indicate that strings might not be favoured after all and that there exists a more fundamental paradigm yet to be revealed.

Strings are not the subject of the present thesis but serve as a motivation for studying conformal field theory (CFT). A propagating string sweeps out a two dimensional surface (the world sheet) and it is thus a two dimensional quantum field theory that describes the string coordinates as functions of the world sheet coordinates. Consistency conditions demand it to be a CFT. Studying conformally invariant *quantum* field theory within string theory may be seen as a study of *classical* solutions to string theory, with interactions according to tree diagrams. Perturbative quantum corrections are dealt with by allowing the world sheet to have

increasingly complicated topology as characterised by its genus (number of handles) which is then playing the number of loops in ordinary field theory Feynman diagrams.

The other main physical interest in CFT is due to its use in descriptions of critical phenomena in statistical mechanics. Two dimensional statistical systems enjoy scale (and indeed conformal [115, 31]) invariance at second order phase transitions due to the divergence of correlation lengths.

Besides yet other applications of CFT's in physics and in mathematics, their relevance may be argued for by somewhat more philosophical comments. A CFT may be defined without referring to any specific classical theory from the outset and is thus free of more or less well-defined quantisation procedures and subsequent renormalisation programmes. Most of the ingredients in CFT inspired by similar properties of usual quantum field theory may be seen as defining properties of CFT and are independent of regularisations and renormalisations. Such considerations naturally lead one to axiomatic approaches to CFT about which we shall have no more to say here.

Consistency in the standard bosonic string theory demands the dimension of the target space to take on the critical value 26. However, at the cost of introducing a new field the dimensionality is no longer fixed and one speaks of non-critical strings. It is the two dimensional world sheet *metric* that becomes a dynamical quantum field and in a certain gauge the surviving freedom is assigned the so-called Liouville field. Within the framework of non-critical strings, much progress has been made in describing the coupling of minimal conformal matter to $2D$ gravity. Extending the Virasoro algebra, which is inherent in every CFT, leads to more complicated and perhaps even more realistic string theories. The best known ones are superstrings based on supersymmetric extensions. However, also bosonic extensions exist and are based on W -algebras. A broad class of these may be constructed by Hamiltonian (Drinfeld-Sokolov) reduction of affine Lie algebras. The Virasoro algebra itself is obtained by Hamiltonian reduction of affine $SL(2)$ algebra and in order to produce the minimal CFT one needs to consider so-called *admissible* representations of the affine $SL(2)$ current algebra. Such representations will play a significant role in this thesis and are generalisations of the well-known integrable (unitary) representations. The former allow the spins j of the fields to be fractional, while in integrable representations $2j$ is integer.

W -strings provide us with means of defining strings consistently in various backgrounds, but they do not admit an immediately transparent way to compute correlation functions. However, H.-L. Hu and M. Yu [81] and Aharony, Ganor, Sonnenschein, Yankielowicz and Sochen [2] have outlined an alternative approach based on (topological) G/G WZNW models. In the non-critical string theory where the (extended) conformal matter is coupled to $2D$ gravity, one performs essentially two reductions in order to eliminate degrees of freedom. The matter and gravity parts are obtained using Hamiltonian reduction of WZNW models and subsequently further reduced by imposing a BRST cohomology condition to remove unwanted surviving degrees of freedom. In the G/G approach the elimination is effectuated in only one step as a BRST condition. A comparison of these two procedures was carried out in [81, 2] in the case of $G = SL(2, \mathbb{R})$. The equivalence of the $SL(2, \mathbb{R})/SL(2, \mathbb{R})$ model and the usual non-critical bosonic string, where minimal conformal matter is coupled to Liouville gravity, was established at the level of isomorphic field and state contents. Later this equivalence for $G = SL(N, \mathbb{R})$ and W_N -strings has been addressed in [3, 125]. The ultimate test in the case of $SL(2, \mathbb{R})$ is to calculate the correlation functions and compare with known results from e.g. matrix

models. For the description of coupling minimal matter to $2D$ gravity, the powerful methods of matrix models in the discrete approach are still superior to the prospects of the G/G approach. However, more general string theories do not seem tractable in terms of matrix models. For higher groups and supergroups the G/G approach might ultimately provide us with a way of formulating and understanding non-critical string theory, in a way for which computation of correlation functions is feasible.

This brings us to the subject of the present thesis, namely computation of Greens functions in CFT based on affine current algebra. We shall use free field realizations which often facilitate computations. The first obstacle is thus to work out the free field realizations of the affine current algebra, the primary fields and the screening currents. The latter constitute essential technical ingredients in the constructions. In this thesis we shall present explicit free field realizations of simple affine current algebras, thus completing the long-standing search for a generalisation of the Wakimoto construction of $SL(2)$ current algebra to any simple Lie group. In the case of the screening currents and the primary fields we still only have partial results in the most general case, but are currently witnessing further progress [113].

In the case of $SL(2)$ such realizations have been known for some time. The free field realization of the current algebra is due to Wakimoto [139], the completion of the set of screening currents is due to Bershadsky and Ooguri [24], while the construction of the primary fields in the framework of depending on an extra isotopic coordinate, is due to Furlan, Ganchev, Paunov and Petkova [64]. N -point functions of two dimensional WZNW theories based on affine $SL(2)$ current algebra have been studied much already. They are typically constructed, either by applying the free field realization of Wakimoto or by solving the Knizhnik-Zamolodchikov equations. The results of these various pieces of works are quite complete as far as unitary, integrable representations are concerned, but are incomplete in the case of admissible representations, which as already mentioned are relevant in the $SL(2)/SL(2)$ approach to non-critical bosonic strings. The origin of this incompleteness may be traced to the need in the free field approach of ghost fields raised to fractional powers. In this thesis we overcome this difficulty by showing how the techniques of fractional calculus naturally provide a way to handle such items. A major part of the thesis is then concerning applications of these techniques in the free field realization, working out general N -point functions, 4-point Greens functions and operator algebra coefficients in CFT based on affine $SL(2)$ current algebra in the case of admissible representations. In terms of free field realizations we have thereby solved completely CFT (on the sphere) based on affine $SL(2)$ current algebra for admissible representations.

This thesis is organised as follows. The chapters 2 to 5 all start with overviews. In Chapter 2 we discuss some concepts in CFT. Notably, we consider free fields and introduce the aforementioned techniques for handling rational powers of ghost fields. Since a major part of the thesis is concerned with a generalisation of the Dotsenko-Fateev construction [42] from pure CFT to CFT based on affine $SL(2)$ current algebra, we briefly review elements of that construction. In that connection we present a proof at the level of correlators of a hidden duality in those constructions. The underlying integral identity will be needed and generalised in Chapter 5.

In Chapter 3 we present explicit free field realizations of simple affine current algebras. The method employed is to first work out differential operator realizations of the simple Lie algebras using Gauss decompositions. Secondly one quantises the realization by translating

into free fields and subsequently adding appropriate anomalous terms to the lowering operators. These terms take care of multiple contractions. Furthermore, free field realizations of screening currents of both kinds and of primary fields are discussed. Many new results of general character are presented. For $SL(3)$ the results are complete. All of these free field realizations are needed for future generalisations to higher groups, of the work on $SL(2)$ presented in chapters 4 and 5 in this thesis. In connection with the construction of the screening currents we digress on a quantum group structure.

In Chapter 4 we develop integral representations for chiral N -point blocks in CFT based on affine $SL(2)$ current algebra in the case of admissible representations. We show at the level of correlators that they indeed satisfy the Knizhnik-Zamolodchikov equations and that they are projectively invariant. Considering the 3-point functions and utilising a freedom (leading to the notion of over-screening) in the choice of numbers of integrations in the realization, we re-derive the fusion rules for admissible representations within our framework. The proposal in [64] for how correlators in $SL(2)$ current algebra for admissible representations reduce to correlators in conformal minimal models is discussed and a simple proof is presented. The reduction is then verified explicitly for the blocks just found. We further discuss the relation to more standard formulations of Hamiltonian reduction.

In Chapter 5 we confine ourselves to the study of 4-point functions and perform the generalisation mentioned above of the famous work [42] by Dotsenko and Fateev on minimal CFT. A different integral realization of the 4-point functions by Andreev [8] is discussed. We present a proof at the level of highly non-trivial integral manipulations generalising the similar proof in Chapter 2 for minimal models, of the equivalence of the two representations. For both representations, the integration contours are found and shown to produce the correct singular behaviour for certain blocks. In his work Andreev does not address choosing contours. Due to simpler contours, we use Andreev's representation to work out the crossing matrix, connecting different bases of blocks, in order to determine the monodromy invariant 4-point Greens functions. We then use the monodromy coefficients to obtain the operator algebra coefficients for theories based on admissible representations. Again the notion of over-screening is utilised, and in such a way that we may normalise the operator algebra coefficients unambiguously, contrary to the results in [8].

Finally, Chapter 6 contains some concluding remarks and a brief discussion of present and future works.

Chapter 2 contains results published in [110] and [112].

Chapter 3 is based on [113]. The subject is currently under further investigation.

Chapters 4 and 5 comprise an exposition of [110, 111, 112].

Acknowledgement

The author has the greatest admiration for his supervisor Jens Lyng Petersen, both as a teacher and as a person. His tremendous efforts are sincerely appreciated.

It is with great pleasure that the author also thanks Yu Ming, without whom this work could never have been undertaken. Throughout the collaboration Yu Ming has been an indispensable source of inspiration.

The author is also indebted to Anna Tollstén for discussions at the early stages of this work. Furthermore, he wants to thank fellow students and the staff and guests at The Niels Bohr Institute for providing a pleasant and inspiring environment.

Finally, he is grateful for the very kind hospitality he met at Institute of Theoretical Physics, Academia Sinica in Beijing, where parts of the work were carried out.

Chapter 2

Conformal Field Theory

The list of references to original works and reviews on conformal field theory (CFT) seems inexhaustible. First of all there is the seminal work by Belavin, Polyakov and Zamolodchikov [21] while the following list reflects some personal preferences [108, 71, 83, 5, 109, 95, 128] and [84] is a reprint collection. These will be used without further notice and may be consulted for details.

We shall introduce techniques of fractional calculus [124, 102, 126, 96] in order to handle ghost fields raised to non-integer powers [110]. Similar techniques were recently discussed by Andreev [8]. Also fractional (indeed complex) powers of algebra generators have been successfully employed in [101, 12, 54, 10]. However, the application was to non-integral powers of algebra generators, whereas we shall consider non-integer powers of free fields where the question of Wick contractions brings in new features.

In connection with a brief review of some aspects in minimal models, we shall present at the level of integral manipulations a proof of a hidden duality in those models.

2.1 Some Basic Concepts

The content of this section serves to fix notation and present some background material.

2.1.1 Conformal Transformations

The conformal group is defined as the subgroup of the general coordinate transformations (group of diffeomorphisms) that leaves the metric invariant up to a space-time dependent scale factor, such that it acts on the metric as a Weyl transformation

$$\begin{aligned} x^\mu &\rightarrow x'^\mu(x) \\ g_{\mu\nu}(x) &\rightarrow g'_{\mu\nu}(x') = \Omega(x)g_{\mu\nu}(x) \quad , \quad \Omega(x) = e^{\omega(x)} \end{aligned} \quad (2.1)$$

It is not difficult to verify that in a flat d -dimensional space-time of signature (p, q) the generators form an algebra isomorphic to $so(p+1, q+1)$. In two dimensions the situation is radically different and in complex coordinates $z = x^1 + ix^2$ and $\bar{z} = x^1 - ix^2$ ($g_{ij} = \delta_{ij}$) the local conformal transformations can be identified with locally analytic coordinate transformations

$$z \rightarrow f(z) \quad , \quad \bar{z} \rightarrow \bar{f}(\bar{z}) \quad (2.2)$$

where $\partial = \partial_z = \frac{1}{2}(\partial_1 - i\partial_2)$ and $\bar{\partial} = \partial_{\bar{z}} = \frac{1}{2}(\partial_1 + i\partial_2)$. Being careful with the Jacobian one finds $d^2z = 2dx^1dx^2$ while the components of the metric are $g_{z\bar{z}} = g_{\bar{z}z} = \frac{1}{2}$ and $g_{zz} = g_{\bar{z}\bar{z}} = 0$. Eventually one is interested in computing correlation functions which are functions of a finite number of points. Therefore, it is indeed natural to extend the class of transformations considered to include 'truly local' or infinitesimal conformal transformations. These may be defined as analytic transformations admitting the infinitesimal coordinate change $\epsilon(z)$ to be holomorphic in the interior of some (not necessarily connected) domain D and identically zero in the exterior. In the boundary region this is not a usual conformal mapping. The emphasis here is on locality.

The infinitely many generators of the two dimensional conformal transformations are described using the basis (labelled by $n \in \mathbb{Z}$)

$$\begin{aligned} z &\rightarrow z' + \epsilon_n(z) & , & & \bar{z} &\rightarrow \bar{z}' + \bar{\epsilon}_n(\bar{z}) \\ \epsilon_n(z) &= -z^{n+1} & , & & \bar{\epsilon}_n(\bar{z}) &= -\bar{z}^{n+1} \end{aligned} \quad (2.3)$$

and are given by

$$l_n = -z^{n+1}\partial_z \quad , \quad \bar{l}_n = -\bar{z}^{n+1}\partial_{\bar{z}} \quad (2.4)$$

The local conformal algebra is achieved by working out the commutation relations

$$[l_m, l_n] = (m - n)l_{m+n} \quad , \quad [\bar{l}_m, \bar{l}_n] = (m - n)\bar{l}_{m+n} \quad , \quad [l_m, \bar{l}_n] = 0 \quad (2.5)$$

The last relation indicates the splitting of the full algebra into a direct sum of two isomorphic subalgebras (each of which is isomorphic to the Witt algebra), which justifies the common use of z and $\bar{z}$ as independent coordinates. In the bigger complexified space ($\mathbb{C} \rightarrow \mathbb{C}^2$) on which the full algebra works, one may then choose various reality conditions. Conventionally one imposes the 'physical' condition $\bar{z} = z^*$ (here $*$ means complex conjugation) and recovers the Euclidean plane. It is customary and convenient to make use of the splitting of the algebra by ignoring the anti-holomorphic part and only in the end to reconstruct the full theory simply by adding terms with bars when appropriate. This will be demonstrated in connection with Greens functions.

Only a finite subalgebra of the local algebra is well-defined globally on the Riemann sphere ($S^2 = \mathbb{C} \cup \infty$). Non-singularity of the vector field $v(z) = \sum_n a_n z^{n+1} \partial_z$ as $z \rightarrow 0$ resp. as $z \rightarrow \infty$ allows $a_n \neq 0$ only for $n \leq 1$ resp. $n \geq -1$ (seen by using the transformation $z = -1/w$). The global conformal group on S^2 is therefore generated by $\{l_{-1}, l_0, l_1\} \cup \{\bar{l}_{-1}, \bar{l}_0, \bar{l}_1\}$. The finite transformations (projective conformal transformations) form the complex Möbius group

$$z \rightarrow \frac{az + b}{cz + d} \quad a, b, c, d \in \mathbb{C} \quad , \quad ad - bc = 1 \quad (2.6)$$

and is isomorphic to $SL(2, \mathbb{C})/\mathbb{Z}_2$ where the quotient by $\mathbb{Z}_2$ is due to invariance under $(a, b, c, d) \rightarrow (-a, -b, -c, -d)$. It is well-known that these are the only analytic one-to-one maps of S^2 .

It is the existence of an infinite dimensional symmetry algebra that makes the *two* dimensional case unique by implying severe restrictions on the conformally invariant field theory. In the remaining part of the thesis considerations are confined to two dimensional CFT.

2.1.2 Fields and Operator Product Expansions

Consider the response of the correlator or expectation value

$$\langle F[\phi] \rangle = \int D\phi e^{-S[\phi]} F[\phi] \quad (2.7)$$

to a transformation

$$\phi'(x) = \phi(x) + \delta_\epsilon \phi(x) \quad (2.8)$$

ϕ is collectively denoting the fields in the system. One has

$$\begin{aligned} 0 &= \int D\phi' e^{-S[\phi']} F[\phi'] - \int D\phi e^{-S[\phi]} F[\phi] \\ &= \int \delta_\epsilon (D\phi e^{-S[\phi]}) F[\phi] + \int D\phi e^{-S[\phi]} \delta_\epsilon F[\phi] \\ &= -\frac{1}{2\pi} \int D\phi e^{-S[\phi]} \int d^2x T^{\mu\nu}(x) \partial_\mu \epsilon_\nu(x) F[\phi] + \langle \delta_\epsilon F[\phi] \rangle \end{aligned} \quad (2.9)$$

This is the definition of the energy-momentum tensor $T_{\mu\nu}$ suggested here. It is not useful to define the energy-momentum tensor in the standard way as the variation of S wrt a change in the metric $T_{\mu\nu} \sim \frac{\delta S}{\delta g^{\mu\nu}}$ since for a generally covariant theory this defines a *tensor* object. As will become clear T is often not a true tensor in CFT. Invariance under scaling transformations immediately shows that T is traceless, $\langle T^\mu{}_\mu F[\phi] \rangle = 0$, while invariance under rotations makes T symmetric, $\langle T_{\mu\nu} F[\phi] \rangle = \langle T_{\nu\mu} F[\phi] \rangle$. Invariance under translations gives a conserved (divergence free) energy-momentum tensor, $\langle \partial_\mu T^{\mu\nu} F[\phi] \rangle = 0$, since one may integrate by parts due to the vanishing outside D . In complex notation this means (introducing $T \equiv T_{zz}, \bar{T} \equiv T_{\bar{z}\bar{z}}$)

$$\langle T_{z\bar{z}} F[\phi] \rangle = \langle T_{\bar{z}z} F[\phi] \rangle = \langle \bar{\partial} T F[\phi] \rangle = \langle \partial \bar{T} F[\phi] \rangle = 0 \quad (2.10)$$

Using these properties one has

$$\begin{aligned} \langle \delta_\epsilon F[\phi] \rangle &= \frac{1}{4\pi} \int_D d^2z \langle (T^{\bar{z}\bar{z}} \bar{\partial} \epsilon_{\bar{z}} + T^{zz} \partial \epsilon_z) F[\phi] \rangle \\ &= \frac{1}{2\pi} \int_D d^2z \langle (\bar{\partial} (T_{zz} \epsilon^z) + \partial (T_{\bar{z}\bar{z}} \bar{\epsilon}^{\bar{z}})) F[\phi] \rangle \\ &= \oint_{\partial D} \left(\frac{dz}{2\pi i} \langle \epsilon(z) T(z) F[\phi] \rangle - \frac{d\bar{z}}{2\pi i} \langle \bar{\epsilon}(\bar{z}) \bar{T}(\bar{z}) F[\phi] \rangle \right) \end{aligned} \quad (2.11)$$

where the second equality is due to $(\bar{\epsilon})$ ϵ being (anti-)holomorphic except in an infinitesimal region. The last equality uses Stoke's theorem

$$\frac{1}{2\pi} \int_D d^2z \bar{\partial} f(z, \bar{z}) = \oint_{\partial D} \frac{dz}{2\pi i} f(z, \bar{z}) \quad , \quad \frac{1}{2\pi} \int_D d^2z \partial f(z, \bar{z}) = - \oint_{\partial D} \frac{d\bar{z}}{2\pi i} f(z, \bar{z}) \quad (2.12)$$

(2.11) is the conformal Ward identity and is a relation between correlation functions. It is customary to write such relations as operator equations

$$\delta_\epsilon F[\phi] = \oint_C \frac{dz}{2\pi i} \epsilon(z) T(z) F[\phi] - \oint_C \frac{d\bar{z}}{2\pi i} \bar{\epsilon}(\bar{z}) \bar{T}(\bar{z}) F[\phi] \quad (2.13)$$

and likewise $T^\mu{}_\mu = 0, \bar{\partial}T = 0, T_{z\bar{z}} = 0$ etc. remembering that they are valid within correlation functions. (2.13) is also denoted the conformal Ward identity. In the sequel the anti-holomorphic sector will often be neglected.

In quantum field theory time ordering in operator products is imposed. After the convenient conformal map

$$z = e^\zeta = e^{\tau+i\sigma} \quad (2.14)$$

from the cylinder (regularised spacetime, (τ, σ) are the Euclidean time and compactified space coordinates respectively) into the complex plane, time ordering becomes a radial ordering. In CFT operator products are always radial ordered (here only bosonic fields are encountered)

$$R(A(z)B(w)) = \begin{cases} A(z)B(w) & , \quad |w| < |z| \\ B(w)A(z) & , \quad |z| < |w| \end{cases} \quad (2.15)$$

though usually R is left understood. One speaks of a radial time $\tau \sim \ln|z|$.

The generator of infinitesimal conformal transformations (2.13) is

$$L_\epsilon = \oint_C \left(\frac{dz}{2\pi i} \epsilon(z) T(z) - \frac{d\bar{z}}{2\pi i} \bar{\epsilon}(\bar{z}) \bar{T}(\bar{z}) \right) \quad (2.16)$$

Among the fields in the theory those that transform¹ as

$$\delta_\epsilon \phi(z, \bar{z}) = \left((h\partial\epsilon(z) + \epsilon(z)\partial) + (\bar{h}\bar{\partial}\bar{\epsilon}(\bar{z}) + \bar{\epsilon}(\bar{z})\bar{\partial}) \right) \phi(z, \bar{z}) \quad (2.17)$$

showing that $\phi(z, \bar{z})(dz)^\Delta(d\bar{z})^{\bar{\Delta}}$ is conserved, are called primary or conformal tensor fields. $(\Delta, \bar{\Delta})$ are the weights of ϕ and are independent. $\Delta + \bar{\Delta}$ determines the behaviour under scalings and is called the dimension, while $\Delta - \bar{\Delta}$ determines the behaviour under rotations and is called the spin. For chiral fields $\phi(z, \bar{z}) = \phi(z)$ one has $\bar{\Delta} = 0$. The radial ordering gives the following description of the generator L_ϵ

$$[L_\epsilon, \phi(w)] = \left(\oint_{|w| < |z|} - \oint_{|z| < |w|} \right) \frac{dz}{2\pi i} \epsilon(z) T(z) \phi(w) = \oint_w \frac{dz}{2\pi i} \epsilon(z) T(z) \phi(w) \quad (2.18)$$

using a contour deformation, where the last integration is around w . Cauchy's integral formula gives an alternative way of classifying primary fields

$$T(z)\phi(w) = \frac{\Delta(\phi)}{(z-w)^2}\phi(w) + \frac{1}{z-w}\partial\phi(w) + nst \quad (2.19)$$

This is an operator product expansion (OPE) where it is assumed that the lhs is analytic in a neighbourhood around w . nst (non-singular terms) is usually not written. The algebra of the fields under operator products (the operator product algebra) is associative. The Ward identity for an n -point function of primary fields may now be written

$$\langle T(z)\phi_1(z_1)\dots\phi_n(z_n) \rangle = \sum_{i=1}^n \left(\frac{h_i}{(z-z_i)^2} + \frac{1}{z-z_i}\partial_{z_i} \right) \langle \phi_1(z_1)\dots\phi_n(z_n) \rangle \quad (2.20)$$

¹The sign in the anti-holomorphic sector is accounted for by $\oint_0 \frac{d\bar{z}}{2\pi i} \bar{z}^{-1} = -1$

Since $\delta_\epsilon \langle \prod_i \phi_i \rangle = 0$ for a global conformal transformation where ϵ is a second degree polynomial, Cauchy's integral formula on the integrated version (2.11) of (2.20) shows that

$$\langle T(z) \phi_1(z_1) \dots \phi_n(z_n) \rangle_{z \rightarrow \infty} \sim z^{-4} f(z_1, \dots, z_n) \quad (2.21)$$

for some function f .

Assume here that the operator product $A(z)B(w)$ can be expanded as

$$A(z)B(w) = \sum_{n=-n_0}^{\infty} (z-w)^n (AB)_n(w) \quad (2.22)$$

A normal ordering is a prescription for handling the singular limit $z \rightarrow w$ for $n_0 > 0$. There are many possibilities. One is to use a point-splitting regularisation [14]: Denote the normal ordered product $: A(w)B(w) :$ and define it by

$$\begin{aligned} : A(w)B(w) : &= \oint_w \frac{dz}{2\pi i} \frac{A(z)B(w)}{z-w} \\ &= (AB)_0(w) \end{aligned} \quad (2.23)$$

This is neither commutative nor associative, e.g. one finds

$$(AB)_0(z) - (BA)_0(z) = \sum_{n=1}^{n_0} \frac{(-1)^{n+1}}{n!} \partial^n (AB)_{(-n)}(z) \quad (2.24)$$

It satisfies Wick's rule

$$A(\underbrace{z})(\underbrace{B}C)(w) = \oint_w \frac{dx}{2\pi i} \frac{1}{x-w} \left(A(\underbrace{z})B(x)C(w) + B(x)A(\underbrace{z})C(w) \right) \quad (2.25)$$

where the under-brace indicates that one is only considering the contraction or singular part of that OPE.

2.1.3 Virasoro Algebra

It is a standard exercise in CFT to show that a 2-point function of quasi-primary fields is fixed up to a normalisation constant C_{12} by the global invariances

$$\langle A_1(z_1)A_2(z_2) \rangle = \frac{C_{12}}{z_{12}^{2h_1}} \delta_{h_1, h_2} \quad , \quad z_{12} = z_1 - z_2 \quad (2.26)$$

A quasi-primary field is primary under projective (global) Möbius transformations. Using (2.21) one expects

$$\langle T(z)T(w) \rangle = \frac{c/2}{(z-w)^4} \quad (2.27)$$

where the constant $\frac{1}{2}$ is chosen as to produce $c = 1$ for free bosons (see below). c is called the central charge and plays a prominent role in CFT. Thus T has $\Delta = 2$ and the Ward identity gives

$$\langle \delta_\epsilon T \rangle = \frac{1}{12} c \partial^3 \epsilon \quad (2.28)$$

For $c \neq 0$ T is not primary. The minimal deviation is in the postulate

$$\delta_\epsilon T = (2\partial\epsilon + \epsilon\partial)T + \frac{1}{12}c\partial^3\epsilon \quad (2.29)$$

or equivalently in the defining OPE

$$T(z)T(w) = \frac{c/2}{(z-w)^4} + \frac{2}{(z-w)^2}T(w) + \frac{1}{z-w}\partial T(w) \quad (2.30)$$

The finite version of (2.29) is given the ansatz under $z \rightarrow w(z)$

$$T(z) = \left(\frac{dw}{dz}\right)^2 T(w) + \frac{c}{12}\{w, z\} \quad (2.31)$$

where $\{w, z\}$ is to be determined. Two successive transformations $z \rightarrow u(z) \rightarrow w(u(z))$ result in

$$\{w, z\} = \left(\frac{du}{dz}\right)^2 \{w, u\} + \{u, z\} \quad (2.32)$$

It may be argued that the solution to this equation consistent with (2.29) is

$$\{w, z\} = \frac{\partial^3 w(z)\partial w(z) - \frac{3}{2}(\partial^2 w(z))^2}{(\partial w(z))^2} \quad (2.33)$$

which is called the Schwartzian derivative. It is easily seen to vanish for a Möbius transformation verifying that T is quasi-primary.

The commutator of two generators is found to be

$$[L_{\epsilon_1}, L_{\epsilon_2}] = L_{\partial\epsilon_1\epsilon_2 - \epsilon_1\partial\epsilon_2} + \frac{c}{12} \oint_0 \frac{dz}{2\pi i} \epsilon_2(z) \partial^3 \epsilon_1(z) \quad (2.34)$$

Choosing the basis $\epsilon_n(z) = z^{n+1}$ and performing a Laurant or mode expansion

$$T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} \quad , \quad L_n = \oint_0 \frac{dz}{2\pi i} T(z) z^{n+1} \quad (2.35)$$

the commutator becomes the conventional form of the Virasoro algebra

$$[L_n, L_m] = (n-m)L_{n+m} + \frac{c}{12}n(n^2-1)\delta_{n+m,0} \quad (2.36)$$

In terms of the modes the condition (2.17) reads

$$[L_n, \phi(z)] = z^n(\Delta(n+1) + z\partial)\phi(z) \quad (2.37)$$

Another standard exercise in CFT is to show that the Virasoro algebra (Vir) is a one-dimensional central extension of the Witt algebra. In the physics literature c is usually treated as a c-number (as in this thesis) though for Vir to be an algebra $c \in Vir$. The treatment is justified by the fact that in any representation c has constant eigenvalue. The central charge also appears in other respects. In a free bosonic scalar field theory c counts the degrees of

freedom. In a CFT defined on a curved background the tracelessness of T is broken by an anomaly proportional to c and the curvature scalar [59]. On the cylinder one has using (2.31) on (2.14)

$$T_{cyl}(z) = z^2 T(z) - \frac{c}{24} \quad (2.38)$$

so c enters in the eigenvalues of the Hamiltonian on the cylinder, $H = L_0 - \frac{c}{24} + \bar{L}_0 - \frac{\bar{c}}{24}$ (in general the central charge $\bar{c}$ of the anti-holomorphic sector is not necessarily equal to c). This is sometimes referred to as understanding the central charge as a Casimir effect, due to the finite geometry of the cylinder.

2.1.4 Descendant Fields

Descendant fields wrt ϕ are defined as finite linear combinations of fields of the form

$$[\dots[L_{\epsilon_2}, [L_{\epsilon_1}, \phi]]\dots] \quad (2.39)$$

and comprise the infinite dimensional conformal family $[\phi]$. Choosing the above conventional basis a descendant field may be written ($\vec{n} = (n_1, \dots, n_m)$)

$$\begin{aligned} \phi^{(-\vec{n})}(z) &= \hat{L}_{-n_1}(z) \dots \hat{L}_{-n_m}(z) \phi(z) \\ &= \oint_{C_1} \frac{dz_1}{2\pi i} \frac{T(z_1)}{(z_1 - z)^{n_1-1}} \dots \oint_{C_m} \frac{dz_m}{2\pi i} \frac{T(z_m)}{(z_m - z)^{n_m-1}} \phi(z) \end{aligned} \quad (2.40)$$

where C_j surrounds $z_{j+1}, \dots, z_m, z$ and $\hat{L}_n(w)$ appears in a formal expansion around w

$$T(z) = \sum_{n \in \mathbb{Z}} \frac{\hat{L}_n(w)}{(z - w)^{n+2}} \quad , \quad \hat{L}_n(w) = \oint_w \frac{dz}{2\pi i} T(z) (z - w)^{n+1} \quad (2.41)$$

Obviously one has

$$T(z) \phi(w) = \sum_{n \geq 0} (z - w)^{n-2} \phi^{(-n)}(w) \quad (2.42)$$

and by evaluating $\delta_\epsilon(T(z) \phi(w))$ one finds by straightforward calculations

$$\begin{aligned} \delta_\epsilon \phi^{(-n)}(w) &= ((\Delta + n) \partial \epsilon(w) + \epsilon(w) \partial) \phi^{(-n)}(w) \\ &+ \sum_{i=1}^n \frac{n+i}{(i+1)!} \partial^{i+1} \epsilon(w) \phi^{(i-n)}(w) + \frac{c}{12(n-2)!} \partial^{n+1} \epsilon(w) \phi(w) \end{aligned} \quad (2.43)$$

and

$$\begin{aligned} T(z) \phi^{(-n)}(w) &= \frac{c}{12} n(n^2 - 1) (z - w)^{-n-2} \phi(w) \\ &+ \sum_{i=1}^n (n+i) (z - w)^{-i-2} \phi^{(i-n)}(w) + \sum_{i \geq 0} (z - w)^{i-2} \phi^{(-i, -n)}(w) \end{aligned} \quad (2.44)$$

T itself is a descendant of the identity I , $T(z) = I^{(-2)}(z)$.

2.1.5 States

The vacuum state $|0\rangle$ may be defined by the condition that it respects the maximum number of symmetries. This means that it is annihilated by the maximum number of conserved charges (the L_n 's in pure CFT, see later for more general situations). Due to the central charge there are only two possibilities, $L_n|0\rangle = 0$ for either $n \geq -1$ or $n \leq 1$. Since the Hamiltonian is bounded (and Hermitian, allowing it to be diagonalised) from below one has

$$L_n|0\rangle = 0 \quad , \quad n \geq -1 \quad (2.45)$$

and similarly in the anti-holomorphic sector. Primary states are defined as

$$|\Delta, \bar{\Delta}\rangle = \lim_{z, \bar{z} \rightarrow 0} \phi(z, \bar{z})|0\rangle \quad (2.46)$$

and using (2.37), seen to respect

$$L_0|\Delta\rangle = \Delta|\Delta\rangle \quad , \quad L_n|\Delta\rangle = 0 \quad , \quad n \geq 1 \quad (2.47)$$

This defines a highest weight vector (state) of weight Δ . A highest weight module or Verma module (see [86, 103] for mathematical accounts) is characterised by the central charge c and the weight Δ of the highest weight vector, and consists of states created from $|\Delta\rangle$ by acting thereupon by linear combinations of lowering operators L_{-n} , $n \geq 0$. The subspace at level $N \geq 0$ spanned by vectors $L_{-n_1} \dots L_{-n_m}|\Delta\rangle$ for which $n_1 + \dots + n_m = N$, has dimension $p(N)$, where $p(N)$ is the partition of N (the number of possible ways N can be written as a sum of positive integers). $p(N)$ appears in the generating function

$$\prod_{n \geq 1} (1 - x^n)^{-1} = \sum_{n \geq 0} p(n) x^n \quad (2.48)$$

Again using (2.37) one has the correspondence $I \sim |0\rangle$. The act of taking the adjoint of a Hermitian operator in Euclidean space-time corresponds to time reversal. This translates into the CFT notion of adjoint being

$$\phi(z, \bar{z})^\dagger = \frac{1}{\bar{z}^{2\Delta}} \frac{1}{z^{2\bar{\Delta}}} \phi\left(\frac{1}{\bar{z}}, \frac{1}{z}\right) \quad (2.49)$$

and one defines $(\langle 0| = |0\rangle^\dagger)$

$$\langle \Delta, \bar{\Delta}| = \lim_{z, \bar{z} \rightarrow \infty} \langle 0| \phi(z, \bar{z}) z^{2\Delta} \bar{z}^{2\bar{\Delta}} \quad (2.50)$$

For T the notion of adjoint leads to the following hermiticity properties

$$L_n^\dagger = L_{-n} \quad (2.51)$$

2.1.6 Singular States and the Kac Determinant

A singular vector in a highest weight module is a highest weight vector itself. The existence of a singular vector at a level higher than zero renders the generically irreducible highest weight module reducible. Explicit forms for singular vectors in Verma modules for the Virasoro algebra

are considered in [16]. See [53] for a general discussion of representations of the Virasoro algebra. One may obtain knowledge of Virasoro singular vectors by considering the Kac determinants [85]. They appear as determinants of the $p(N) \times p(N)$ dimensional matrices

$$M_{\mathbf{k}\mathbf{k}'}^{(N)} = \langle \Delta | \left(\prod_{n \geq 1} (L_{-n})^{k_n} \right)^\dagger \prod_{m \geq 1} (L_{-m})^{k'_m} | \Delta \rangle \quad (2.52)$$

where $\sum n k_n = \sum m k'_m = N$, $k_n \geq 0$. The products are ordered, $\prod (L_{-n})^{k_n} = \dots (L_{-2})^{k_2} (L_{-1})^{k_1}$, such that we only consider a basis of states at level N . Such a matrix is obviously Hermitian and has zero determinant if there exists an eigenvector $\mathbf{v}^t = (v_1, \dots, v_{p(N)})$ satisfying $\mathbf{v}^t M \mathbf{v} = 0$, indicating that $\|\sum v_i |\Delta_i^{(N)}\rangle\| = 0$ where $\{|\Delta_i^{(N)}\rangle\}$ is the basis chosen. Since determinants are essentially basis independent they uniquely reveal the existence of zero norm vectors. Such vectors are called null-vectors and are easily seen to be orthogonal to all states in the Verma module. Their number at a given level N is equal to the number of zero eigenvalues of the Kac determinant K_N . Null-states appear in submodules generated by singular states different from the highest weight state $|\Delta\rangle$. Kac found the general expression for the determinant K_N [85]

$$K_N = c_N \prod_{n,m > 0, nm \leq N} (\Delta - \Delta_{n,m})^{p(N-nm)} \quad (2.53)$$

with positive constants c_N . In [51] the first proof appeared while [135, 123] contain alternative proofs. The central charge and the conformal weights (the Kac table) are given by

$$\begin{aligned} \Delta_{n,m} &= -\frac{\alpha_0^2}{2} + \frac{1}{8}(n\alpha_+ + m\alpha_-)^2, & n, m \geq 1 \\ \alpha_\pm &= \alpha_0 \pm \sqrt{\alpha_0^2 + 2} = \frac{\sqrt{1-c} \pm \sqrt{25-c}}{\sqrt{12}} \\ c = 1 - 12\alpha_0^2, & \quad 2\alpha_0 = \alpha_+ + \alpha_-, & \quad \alpha_+ \alpha_- = -2 \end{aligned} \quad (2.54)$$

For $c, \Delta > 1, 0$ the Kac determinant has no zeros. A detailed study [61] of K_N reveals that unitarity (only positive norm states) imposes severe restrictions. By looking for zeros for K_N in the plane c, Δ , separating regions of positive and negative norm states, one finds that in order to have unitarity for $c < 1$, the central charge is given by

$$c = 1 - \frac{6}{n(n+1)}, \quad n = 3, 4, 5, \dots \quad (2.55)$$

with the associated parametrisations of Δ (2.54). These are only necessary conditions for unitarity ($c < 1$) while the proof of existence is by coset constructions, see below. Minimal models, to be discussed in a subsequent section, are obtained by imposing less severe restrictions.

2.1.7 Conformal Blocks and Greens Functions

The projective invariance, generated by $\{L_{-1}, L_0, L_1\}$, puts restrictions on the N -point functions, leaving them essentially only dependent on the anharmonic ratios $z_{ijkl} = (z_i - z_j)(z_k -$

$z_l)/((z_j - z_k)(z_l - z_i))$ of which only $N - 3$ are independent. The 3-point Greens functions are thereby fixed

$$\begin{aligned} & \langle \phi_3(z_3, \bar{z}_3) \phi_2(z_2, \bar{z}_2) \phi_1(z_1, \bar{z}_1) \rangle \\ = & C_{123} (z_2 - z_1)^{-\Delta_2 - \Delta_1 + \Delta_3} (z_3 - z_2)^{-\Delta_3 - \Delta_2 + \Delta_1} (z_3 - z_1)^{-\Delta_3 - \Delta_1 + \Delta_2} \\ & \cdot (\bar{z}_2 - \bar{z}_1)^{-\bar{\Delta}_2 - \bar{\Delta}_1 + \bar{\Delta}_3} (\bar{z}_3 - \bar{z}_2)^{-\bar{\Delta}_3 - \bar{\Delta}_2 + \bar{\Delta}_1} (\bar{z}_3 - \bar{z}_1)^{-\bar{\Delta}_3 - \bar{\Delta}_1 + \bar{\Delta}_2} \end{aligned} \quad (2.56)$$

See Chapter 5 for a discussion on the structure constants C_{123} .

Due to the decomposition $Vir \oplus \bar{Vir}$, a general 4-point Greens function (in the 'standard projective limit')

$$G_{ijkl}(z, \bar{z}) = \langle \phi_l(\infty, \infty) \phi_k(1, 1) \phi_j(z, \bar{z}) \phi_i(0, 0) \rangle \quad (2.57)$$

may be expanded in conformal blocks $F_{ijkl}^{(m)}$ (or chiral blocks for bigger symmetry groups than Vir , see later)

$$G_{ijkl}(z, \bar{z}) = \sum_m C_{ij}^m C_{mkl} F_{ijkl}^{(m)}(z) \bar{F}_{ijkl}^{(m)}(\bar{z}) \quad (2.58)$$

If the fusion rule coefficients [138, 38, 105, 33, 5] are larger than one, then the sum over m must be supplemented by a degeneracy index labelling the distinct ways of coupling to the field ϕ_m . However, our prime interest lies in the study of CFT based on $SL(2)$ current algebra where such subtleties are absent, so we will ignore them here. The blocks behave like and are normalised according to

$$\lim_{z \rightarrow 0} F_{ijkl}^{(m)}(z) = z^{\Delta_m - \Delta_i - \Delta_j} (1 + \mathcal{O}(z)) \quad (2.59)$$

and are generically multi-valued functions. Yet, the physical Greens functions are supposed to be single-valued functions. The power-like singular behaviours of the blocks of a given 4-point Greens function are all distinct and determine the non-trivial monodromies for these blocks. Graphically one associates to the blocks, or to the Greens functions, Feynman skeleton diagrams for a φ^3 type quantum field theory with the fields ϕ being the external legs, while the internal lines describe the exchanged fields, see Fig. 1 in Chapter 5. The associativity of the operator product algebra leads to some duality relations (crossing symmetry or bootstrap equations) expressing the Greens function in different expansions

$$\begin{aligned} \sum_m C_{ij}^m C_{mkl} F_{ijkl}^{(m)}(z) \bar{F}_{ijkl}^{(m)}(\bar{z}) &= \sum_m C_{jk}^m C_{mil} F_{jkil}^{(m)}(1 - z) \bar{F}_{jkil}^{(m)}(1 - \bar{z}) \\ &= z^{-2\Delta_j} \bar{z}^{-2\bar{\Delta}_j} \sum_m C_{jl}^m C_{mik} F_{ikjl}^{(m)}(1/z) \bar{F}_{ikjl}^{(m)}(1/\bar{z}) \end{aligned} \quad (2.60)$$

We shall refer to these channels as s-, t- and u-channels as is standard Mandelstam terminology in relativistic kinematics describing particle scattering. Similarly one may define conformal blocks for higher N -point Greens functions. Greens functions are always supposed to be single-valued. When the summations (2.60) truncate (which is the case in rational CFT, see later), the duality means that the blocks transform into finite linear combinations of themselves under analytic continuation. Being multi-valued functions the blocks are indeed supposed to be defined first in one domain (for a given order of the arguments) and then analytically continued to other domains. Since the analytic continuation depends on the path chosen, it is the braid group and not the permutation group that acts on the blocks. In this respect a quantum group structure is revealed [105, 4, 104, 5].

The decoupling of a null field (the descendant field corresponding to the null vector in the Verma module headed by the corresponding primary field) gives rise to certain differential equations satisfied by the correlation functions and the conformal blocks. Utilising this is sometimes a convenient way of determining the blocks.

The Ward identities for insertion of symmetry generators (here only T) makes it possible to obtain correlation functions involving descendant fields from those of primary fields.

2.2 Free Fields

2.2.1 A Digression on Lie Algebras

Let $\mathfrak{g}$ be a simple Lie algebra of $\dim \mathfrak{g} = d$ and $\text{rank } \mathfrak{g} = r$. $\mathfrak{h}$ is a Cartan subalgebra of $\mathfrak{g}$. The set of (positive) roots is denoted (Δ_+) Δ and we write $\alpha > \beta$ if for two roots $\alpha - \beta \in \Delta_+$, and analogously $\alpha > 0$ means $\alpha \in \Delta_+$. The simple roots are $\{\alpha_i\}_{i=1,\dots,r}$. θ is the highest root and h^ν is the dual Coxeter number. The only non-vanishing elements of the Cartan-Killing form are

$$\kappa_{\alpha,-\alpha} = \kappa(e_\alpha, f_\alpha) = \frac{2}{\alpha^2} \quad , \quad \kappa_{ij} = \kappa(h_i, h_j) = G_{ij} \quad (2.61)$$

The Cartan matrix is $A_{ij} = \alpha_i^\nu \cdot \alpha_j = (\alpha_i^\nu, \alpha_j) = G_{ij} \alpha_j^2 / 2$, while Freudenthal-de Vries strange formula states that

$$\rho^2 = \frac{d}{24} h^\nu \theta^2 \quad (2.62)$$

Using the triangular decomposition

$$\mathfrak{g} = \mathfrak{g}_- \oplus \mathfrak{h} \oplus \mathfrak{g}_+ \quad (2.63)$$

we will denote the raising and lowering operators $e_\alpha \in \mathfrak{g}_+$ and $f_\alpha \in \mathfrak{g}_-$ respectively with $\alpha \in \Delta_+$, and Cartan operators $h_i \in \mathfrak{h}$. We use the common notation j_a for all the algebra elements. For simple roots we sometimes write $e_i = e_{\alpha_i}$, $f_i = f_{\alpha_i}$ and collectively denote the $3r$ generators e_i, h_i, f_i Chevalley generators. Their commutator relations are

$$\begin{aligned} [h_i, h_j] &= 0 & [e_i, f_j] &= \delta_{ij} h_j \\ [h_i, e_j] &= A_{ij} e_j & [h_i, f_j] &= -A_{ij} f_j \end{aligned} \quad (2.64)$$

In addition one has the Serre relations

$$\begin{aligned} (\text{ad}_{e_i})^{1-A_{ij}} e_j &= 0 & , & & i \neq j \\ (\text{ad}_{f_i})^{1-A_{ij}} f_j &= 0 & , & & i \neq j \end{aligned} \quad (2.65)$$

The Weyl vector is defined as $\rho = \frac{1}{2} \sum_{\alpha > 0} \alpha$ and satisfies $\rho \cdot \alpha_i^\nu = 1$. We use the convention $f_{-\alpha, -\beta}^{-\gamma} = -f_{\alpha\beta}^\gamma$. Furthermore

$$\begin{aligned} [e_\alpha, f_\alpha] &= h_\alpha = G^{ij}(\alpha_i^\nu, \alpha^\nu) h_j \\ h_{\alpha+\beta} &= \frac{1}{(\alpha + \beta)^2} (\alpha^2 h_\alpha + \beta^2 h_\beta) \\ [h_i, e_\alpha] &= (\alpha_i^\nu, \alpha) e_\alpha \\ [h_i, f_\alpha] &= -(\alpha_i^\nu, \alpha) f_\alpha \end{aligned} \quad (2.66)$$

2.2.2 Bosonic Scalar Field

Let us consider a set of free bosonic scalar fields $X_i(z, \bar{z})$ labelled by Cartan indices $i = 1, \dots, r$ and described by the action

$$S = -\frac{1}{4\pi} \int d^2z \partial X(z, \bar{z}) \cdot \bar{\partial} X(z, \bar{z}) \quad (2.67)$$

The metric in X -space is the Cartan part G_{ij} of the Cartan-Killing form. In this notation the fields are purely imaginary. The classical equations of motion are

$$\partial \bar{\partial} X_i(z, \bar{z}) = 0 \quad (2.68)$$

from which it follows that ∂X_i ($\bar{\partial} X_i$) is (anti-)analytic and is therefore written $\partial X_i(z)$ ($\bar{\partial} X_i(\bar{z})$). The general solution splits the scalar field into a sum of holomorphic and anti-holomorphic pieces

$$X_i(z, \bar{z}) = \varphi_i(z) + \bar{\varphi}_i(\bar{z}) \quad (2.69)$$

The property of path integrals that the integral of a total derivative vanishes, gives

$$\begin{aligned} 0 &= \int DX \frac{\delta}{\delta X^i(z, \bar{z})} (e^{-S} X_j(w, \bar{w})) \\ &= \langle G_{ij} \delta^2(z - w, \bar{z} - \bar{w}) \rangle - \frac{1}{2\pi} \partial_z \partial_{\bar{z}} \langle X_i(z, \bar{z}) X_j(w, \bar{w}) \rangle \end{aligned} \quad (2.70)$$

Using this together with

$$\partial_z \bar{\partial}_{\bar{z}} \ln |z - w|^2 = 2\pi \delta^2(z - w, \bar{z} - \bar{w}) \quad (2.71)$$

following from Stoke's theorem (2.12), we find the 2-point function

$$\langle X_i(z, \bar{z}) X_j(w, \bar{w}) \rangle = G_{ij} \ln |z - w|^2 \quad (2.72)$$

and the OPE

$$X_i(z, \bar{z}) X_j(w, \bar{w}) = G_{ij} \ln |z - w|^2 + : X_i(z, \bar{z}) X_j(w, \bar{w}) : \quad (2.73)$$

or equivalently

$$\begin{aligned} \varphi_i(z) \varphi_j(w) &= G_{ij} \ln(z - w) + : \varphi_i(z) \varphi_j(w) : \\ \varphi_i(z) \bar{\varphi}_j(\bar{w}) &= : \varphi_i(z) \bar{\varphi}_j(\bar{w}) : \\ \bar{\varphi}_i(\bar{z}) \bar{\varphi}_j(\bar{w}) &= G_{ij} \ln(\bar{z} - \bar{w}) + : \bar{\varphi}_i(\bar{z}) \bar{\varphi}_j(\bar{w}) : \end{aligned} \quad (2.74)$$

whereby we have introduced the normal ordering of free scalar fields by subtracting the singular parts. The 2-point function demonstrates that the free fields do not have well-defined scaling dimensions, hence they do not rigorously exist as quantum fields [35]. Nevertheless, we shall only need to consider constructions such as ∂X_i of dimension 1 and vertex operators to be defined in (2.82). According to (2.71) the normal ordered product satisfies the classical equations of motion and may be Taylor expanded. The first term in the expansion precisely corresponds to the definition of composite operators in (2.23). We have found that inside the

normal ordering the scalar fields act as classical fields. The corresponding energy-momentum tensor is

$$T = \frac{1}{2} : \partial X \cdot \partial X : = \frac{1}{2} : \partial \varphi \cdot \partial \varphi : \quad (2.75)$$

and has central charge $c = r$. This result is the origin of the conventional factor of $1/2$ in front of c in (2.30). The notion of normal ordering is easily generalised along the lines of

$$X_i(z, \bar{z}) : \prod_{j=1}^n X_{i_j}(z_j, \bar{z}_j) := \sum_{j=1}^n G_{ii_j} \ln |z - z_j|^2 : \prod_{l \neq j} X_{i_l}(z_l, \bar{z}_l) : + : X_i(z, \bar{z}) \prod_{j=1}^n X_{i_j}(z_j, \bar{z}_j) : \quad (2.76)$$

and one has that the OPE of two normal ordered operators can be written as the formal expression

$$: G[X] : : F[X] := \exp \left(\int d^2 z d^2 w G_{ij} \ln |z - w|^2 \frac{\delta G}{\delta X_i(z, \bar{z})} \frac{\delta F}{\delta X_j(w, \bar{w})} \right) : G[X] F[X] : \quad (2.77)$$

The outset for a much richer structure is the introduction of a background charge $2a_0$ for the fields X_i [51, 42, 60]. It amounts to adding to the action a term

$$S_{2a_0} = \frac{-a_0}{8\pi} \int d^2 z \sqrt{g} R \rho \cdot X \quad (2.78)$$

coupling the fields X_i to the curvature of the underlying surface. In what follows we will mostly be interested in the Riemann sphere S^2 in which case it is possible to localise the curvature at the point of infinity (the north pole). Hence, everywhere except in the vicinity of the north pole we may use a flat metric and still consider the fields to be free. The equations of motion become

$$\partial \bar{\partial} X_i = \frac{1}{4} a_0 (\alpha_i^\nu, \rho) \sqrt{g} R \quad (2.79)$$

The corresponding energy-momentum tensor is (found by using the conformal gauge or isothermal coordinates $g_{\mu\nu}(z) = e^{\sigma(z)} \delta_{\mu\nu}$)

$$T_\varphi = \frac{1}{2} : \partial \varphi \cdot \partial \varphi : - a_0 \rho \cdot \partial^2 \varphi \quad (2.80)$$

which has central charge

$$c_\varphi = r - 12a_0^2 \rho^2 = r - \frac{\theta^2}{2} a_0^2 h^\nu d \quad (2.81)$$

where we have used (2.62). In the sequel we will concentrate on the holomorphic part φ and put $a_0 = 1/\sqrt{t}$, see (4.2), (4.3).

The vertex operators later to be used in building primary fields are

$$V_\lambda(z) =: e^{\frac{1}{\sqrt{t}} \lambda \cdot \varphi(z)} : \quad (2.82)$$

and are seen to have conformal weights

$$\Delta(V_\lambda) = \frac{1}{2t} (\lambda, \lambda + 2\rho) \quad (2.83)$$

Furthermore

$$V_\lambda(z) V_{\lambda'}(w) = (z - w)^{\frac{1}{t} \lambda \cdot \lambda'} : e^{\frac{1}{\sqrt{t}} (\lambda \cdot \varphi(z) + \lambda' \cdot \varphi(w))} : \quad (2.84)$$

2.2.3 Ghost Fields

Ghost fields appear in gauge theory in the Faddeev-Popov procedure of gauge fixing and in string theory. Usually one denotes a fermionic ghost system b, c arising when gauge fixing a bosonic symmetry, and a bosonic ghost system β, γ arising when gauge fixing a fermionic symmetry. They are described by first order actions (characterised by having first order kinetic terms as for the free Majorana fermion) and are conveniently treated simultaneously by introducing the parameter η , $\eta = -1$ for bosonic statistics and $\eta = +1$ for fermionic statistics. Hence, we will now let b, c denote any ghost pair disregarding their statistics and write

$$S_\eta = \frac{\eta}{2\pi} \int d^2z \left(b\bar{\partial}c + \bar{b}\partial\bar{c} \right) \quad (2.85)$$

The rationale for discussing ghost fields in this thesis is not a wish to treat gauge fixing procedures, rather they appear as building blocks in Wakimoto free field realizations, see Chapter 3. The equations of motion ensure that the fields are (anti-)holomorphic. This action is conformally invariant if b and c have dimensions $(\lambda, 0)$ and $(1 - \lambda, 0)$ respectively, in which case the energy-momentum tensor becomes (focusing on the holomorphic part)

$$T = \eta((1 - \lambda) : \partial b c : - \lambda : b \partial c :) \quad (2.86)$$

As for the free scalar field the 2-point function is easy to determine, this time using $0 = \int \frac{\delta}{\delta b} (e^{-S} b)$, and one arrives at the OPE²

$$\begin{aligned} c(z)b(w) &= \frac{\eta}{z-w} + :c(z)b(w): \\ b(z)c(w) &= \frac{1}{z-w} + :b(z)c(w): \end{aligned} \quad (2.87)$$

The conformal dimensions are readily verified and the central charge is easily found to be

$$c = -2\eta(6\lambda^2 - 6\lambda + 1) = \eta(1 - 3Q^2) \quad (2.88)$$

where $Q = \eta(2\lambda - 1)$.

The standard mode expansions read

$$b(z) = \sum_n b_n z^{-n-\lambda}, \quad c(z) = \sum_n c_n z^{-n-(1-\lambda)} \quad (2.89)$$

and give rise to the (anti-)commutation relation

$$[b_m, c_n]_\eta = \delta_{m+n,0} \quad (2.90)$$

where $[\cdot, \cdot]_\eta$ is a (anti-)commutator for $\eta = -1$ ($\eta = +1$). The ranges for parameters n depend on the choice of sector (Neveu-Schwarz versus Ramond). A zero-mode is one for which $\bar{\partial}$ vanishes throughout the Riemann surface, here the sphere S^2 . Using that for the primary field b , $b(z)(dz)^\lambda$ is conserved under the conformal map $z \rightarrow 1/z$, one finds the zero-modes of b as the ones multiplying functions of z well-defined both in the neighbourhood of 0 and in the

²The somewhat unconventional factor η in the action was chosen to produce $\beta(z)\gamma(w) = \frac{+1}{z-w}$.

neighbourhood of ∞ . This analysis gives an index theorem (Riemann-Roch theorem) for ghost fields on the sphere. In general [60]

$$\sharp(\text{zero-modes of } c) - \sharp(\text{zero-modes of } b) = \frac{1}{2}\eta Q\chi \quad (2.91)$$

where $\chi = 2(1 - g)$ is the Euler characteristic of the Riemann surface and g is the genus.

The ghost number operator is

$$N_{gh} = \oint_0 \frac{dz}{2\pi i} j(z) \quad , \quad j(z) = \eta : b(z)c(z) : \quad (2.92)$$

and satisfies $[N_{gh}, b(w)] = b(w)$ and $[N_{gh}, c(w)] = -c(w)$. The ghost number current is non-primary (or anomalous) for $Q \neq 0$

$$T(z)j(w) = \frac{Q}{(z-w)^3} + \frac{j(w)}{(z-w)^2} + \frac{\partial j(w)}{z-w} \quad (2.93)$$

The hermiticity properties are

$$\begin{aligned} b_n^\dagger &= \eta b_{-n} \\ c_n^\dagger &= c_{-n} \\ j_n^\dagger &= -j_{-n} - Q\delta_{n,0} \end{aligned} \quad (2.94)$$

In Chapter 4 we elaborate on the case of bosonic ghost fields β, γ of dimension $(1,0)$.

In the applications considered in this thesis we shall only need the bosonic ghost fields. It is clear that the following rules apply in computations of OPE's when either β or γ fields appear with integral powers

$$\begin{aligned} \beta(z)^n F(\gamma(w)) &= : (\beta(z) + \frac{1}{z-w} \partial_{\gamma(w)})^n F(\gamma(w)) : \\ \gamma(z)^n F(\beta(w)) &= : (\gamma(z) - \frac{1}{z-w} \partial_{\beta(w)})^n F(\beta(w)) : \end{aligned} \quad (2.95)$$

It turns out (see Chapter 4) that we need to be able to treat ghost fields raised to fractional powers. The proposal in [110] to deal with those cases consists in generalising (2.95) as

$$G(\beta(z))F(\gamma(w)) =: G(\beta(z) + \frac{1}{z-w} \partial_{\gamma(w)})F(\gamma(w)) : \quad (2.96)$$

where the asymptotic expansions for $G(\beta(z) + \frac{1}{z-w} \partial_{\gamma(w)})$ and $F(\gamma(w))$ would depend on their monodromy conditions in the z and w variables respectively. An example relevant in later chapters is

$$\begin{aligned} \beta(z)^{-t} F(\gamma(w)) &= : (\beta(z) + \frac{1}{z-w} \partial_{\gamma(w)})^{-t} F(\gamma(w)) : \\ &= \sum_{n=0}^{\infty} \binom{-t}{n} : \beta^n(z) (z-w)^{t+n} \partial_{\gamma(w)}^{-t-n} F(\gamma(w)) : \end{aligned} \quad (2.97)$$

where t is fractional for admissible representations (to be introduced in Chapter 4) and we then see the need for fractional calculus [124, 102, 126, 96]. As an example of how the technique works we provide in a subsequent section an explicit non-trivial proof (2.121) that

$$(\beta^a(z)\gamma^a(w))(\beta^b(z)\gamma^b(w)) = \beta^{a+b}(z)\gamma^{a+b}(w) \quad (2.98)$$

Additionally, the explicit verification that the N -point functions in $SL(2)$ current algebra based on this proposal satisfy the Knizhnik-Zamolodchikov equations (see Chapter 4), may be viewed as a check that fractional calculus does indeed provide us with the requisite properties for Wick contractions as defined in (2.97).

An alternative to the proposal (2.96) is to write (2.95) as in (2.77)

$$G(\beta(z))F(\gamma(w)) = \exp\left(\frac{1}{z-w}\partial_{\beta(z)}\partial_{\gamma(w)}\right) : G(\beta(z))F(\gamma(w)) : \quad (2.99)$$

and then generalise it by allowing asymptotic expansions (see below) of the exponential as in

$$e^z = \partial^a e^z = \sum_{n \in \mathbb{Z}} \frac{1}{\Gamma(n-a+1)} z^{n-a} \quad , \quad a \in \mathbb{C} \quad (2.100)$$

To reproduce (2.97) we put $a = t$. The gamma-function $\Gamma(a+1)$ is the standard analytic continuation of 'factorial': $\Gamma(n+1) = n!$, n non-negative integer.

For every positive root $\alpha > 0$ we now introduce a pair of free bosonic ghost fields $(\beta_\alpha, \gamma^\alpha)$ of conformal weights (1,0) satisfying the OPE

$$\beta_\alpha(z)\gamma^\beta(w) = \frac{\delta_\alpha^\beta}{z-w} \quad (2.101)$$

The corresponding energy-momentum tensor is

$$T_{\beta\gamma} =: \beta_\alpha \partial \gamma^\alpha : \quad (2.102)$$

with central charge

$$c_{\beta\gamma} = d - r \quad (2.103)$$

We will understand 'properly' repeated root indices as in (2.102) to be summed over the positive roots.

2.2.4 Fractional Calculus and Expansions

Here we briefly introduce fractional calculus [124, 102, 126, 96]. For an analytic function $f(z)$ the fractional derivative (or integration) $\partial^a f(z)$ is defined for complex number a . It satisfies the following axioms:

1. If $f(z)$ is an analytic function, the fractional derivative $\partial^a f(z)$ is an analytic function of z and a .
2. For a integer, the result must agree with ordinary differentiation (a positive) or integration (a negative). By default the integration constants are put to zero, so that the function together with a maximum number of derivatives vanish at some point, like $z = 0$.

3. $a = 0$ is the identity.
4. Fractional differentiation is linear.
5. For fractional integration, $\Re a > 0, \Re b > 0$

$$\partial^{-a}\partial^{-b}f(z) = \partial^{-(a+b)}f(z)$$

For $\Re a > 0$, fractional integration ∂^{-a} may be represented by the Riemann-Liouville operator

$$\partial^{-a}f(z) = \frac{1}{\Gamma(a)} \int_0^z (z-t)^{a-1} f(t) dt \quad (2.104)$$

By induction in the integer n it is shown that for $a = n$ this coincides with ordinary n -fold integration given by

$$\partial^{-n}f(z) = \int_0^z dz_1 \int_0^{z_1} dz_2 \dots \int_0^{dz_{n-1}} dz_n f(z_n) \quad (2.105)$$

The third axiom follows from a Taylor expansion of $f(t)$. The semi-group property of axiom 5 follows from

$$\begin{aligned} \partial^{-a}\partial^{-b}f(z) &= \frac{1}{\Gamma(a)\Gamma(b)} \int_0^z dz_1 \int_{z_1}^z dz_2 (z-z_2)^{a-1} (z_2-z_1)^{b-1} f(z_1) \\ &= \partial^{-(a+b)}f(z) \end{aligned} \quad (2.106)$$

where we have used the substitution $u = \frac{z_2-z_1}{x-z_1}$ and the Euler integral

$$\int_0^1 u^{b-1} (1-u)^{a-1} du = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)} \quad (2.107)$$

Having defined fractional differentiation for arbitrary negative powers, we may obtain fractional derivatives of arbitrary positive powers from these by acting with an integer number of ordinary differentiations. Thus, for $a > 0$ we write $a = n - a'$ where n is a positive integer and $0 \leq a' < 1$, and define

$$\partial^a f(z) = \partial^n \left(\frac{1}{\Gamma(a')} \int_0^z (z-t)^{a'-1} f(t) dt \right) \quad (2.108)$$

For every pair of complex numbers a, b we have $\partial^a \partial^b = \partial^{a+b}$. We now consider the particular case $f(z) = z^b$ and find

$$\partial^a z^b = \frac{\Gamma(b+1)}{\Gamma(b-a+1)} z^{b-a} \quad (2.109)$$

This holds for both fractional integration and fractional differentiation. Notice that $\partial^a 1 \neq 0$ for a not a positive integer.

Let us conclude this section by commenting on various expansions as the one employed in (2.100). Since we shall need contractions with the operator (see chapters 4 and 5)

$$(1 + \gamma(z)x)^{2j} \quad (2.110)$$

we shall find it convenient to represent the expansion of the associated analytic function

$$f(z) = (1+z)^{2j} \quad (2.111)$$

in several different ways. Indeed, from

$$(1+z)^{2j} = z^{-a}[(1+z)^{2j}z^a] \quad (2.112)$$

we may imagine that the last bracket is expanded in integer powers of z in a way convergent on the unit circle. Although $[(1+z)^{2j}z^a]$ has a discontinuity on the circle, it vanishes when we take it to be for negative z , corresponding to choosing the branch cuts along the negative real axis. This renders many equivalent representations for the function

$$(1+z)^{2j} = (1+z)^{2j}_{(-a)} = \sum_{n \in \mathbb{Z}} \binom{2j}{n-a} z^{n-a} \quad (2.113)$$

which are all equivalent in the sense of analytic function theory but which correspond to expansions with different monodromies for the individual terms.

When deciding on what expansion to adopt for the operator

$$(1+\gamma(z)x)^{2j}_{(a)} = \sum_{n \in \mathbb{Z}} \binom{2j}{n+a} (\gamma(z)x)^{n+a} \quad (2.114)$$

we use the criterion, that *after* all Wick contractions have been performed, powers of β and γ inside normal ordering signs are non-negative *integers*. Only then, these terms have an obvious interpretation when sandwiched between states. In other terms, the existence of external states and other primary fields in the correlator decides what monodromies to choose for individual terms in expansions. All of that will be illustrated further in chapters 4 and 5.

Next, consider the fractional derivative of the exponential function

$$D^a \exp(z) = \sum_{n \in \mathbb{Z}} \frac{1}{\Gamma(n-a+1)} z^{n-a}, \quad a \in \mathbb{C} \quad (2.115)$$

The right hand side is covariant under further differentiation corresponding to the fact that it represents the original exponential function. The representation is a peculiar realization of the exponential function, which converges asymptotically for $|z| \rightarrow \infty$. Again the representation may be better understood by writing

$$\exp(z) = z^{-a}[\exp(z)z^a] \quad (2.116)$$

and then introducing for the last bracket a Fourier expansion with integer powers of z on a circle in the complex z plane. On the circle

$$[\exp(z)z^a] \quad (2.117)$$

has a discontinuity which we may take to be for negative z . The Fourier coefficients depend on the radius of the circle. However, for large $|z|$ where the discontinuity becomes vanishingly small, they converge to the expansion coefficients in (2.115). Thus we take

$$D^a \exp(z) = \exp(z) \quad (2.118)$$

for any a . However, we shall find it convenient to use the fractional derivative to represent a generating functional for the integrals

$$\lim_{R \rightarrow \infty} \oint_{RS^1} \frac{du}{2\pi i} \frac{e^u}{u^{a+n+1}} = \frac{1}{\Gamma(a+n+1)} \quad (2.119)$$

Different a 's give rise to representations or expansions of the exponential function in which individual terms have different non-trivial monodromies. We shall use $D^a \exp(z)$ simply as a reminder of the appropriate expansion needed. Let us notice the following identity for fractional derivatives of exponentials

$$D_z^a \exp(z) D_w^b \exp(w) = D^{a+b} \exp(z+w) \quad (2.120)$$

2.2.5 Consistency Check

In this section we illustrate the non-trivial nature of the workings of the Wick contractions proposed in (2.96), by considering the evaluation [110] of

$$I_{a,b}(z, w) = (\beta^a(z) \gamma^a(w)) (\beta^b(z) \gamma^b(w)) \quad (2.121)$$

First we notice the OPE

$$\beta^a(z) \gamma^a(w) = \frac{\Gamma(a+1)}{(z-w)^a} \sum_n \binom{a}{n} \frac{1}{n!} (z-w)^n : (\beta(z) \gamma(w))^n : \quad (2.122)$$

following from (2.96) and (2.109) (or alternatively from (2.99-2.100) and (2.109)). Now, (2.121) may be evaluated in two ways: either (i) by first using (2.122) for both parentheses to reduce both of them to normal ordered products of integer powers of β and γ , and then subsequently carrying out all remaining contractions, or else (ii) by simply using (2.122) with $a+b$ replacing a . Obviously these two ways should lead to the same result for consistency. This requirement is part of the associativity properties for operators, and we consider them to be justified by the fact that the N -point functions of Chapter 4 satisfy the Knizhnik-Zamolodchikov equations. Here we demonstrate that the above condition gives rise to non-trivial identities for which we indicate an independent elementary proof.

The general contractions between integer powers of β and γ are carried out using (2.99)

$$\begin{aligned} \beta^n(z) \gamma^m(w) &= \exp \left(\frac{1}{z-w} \partial_{\beta(z)} \partial_{\gamma(w)} \right) : \beta^n(z) \gamma^m(w) : \\ &= \sum_{\ell} \frac{1}{(z-w)^{\ell} \ell!} \frac{\Gamma(n+1)}{\Gamma(n-\ell+1)} \frac{\Gamma(m+1)}{\Gamma(m-\ell+1)} : \beta^{n-\ell}(z) \gamma^{m-\ell}(w) : \end{aligned} \quad (2.123)$$

Then one obtains after a few steps

$$\begin{aligned} I_{a,b}(z, w) &= \frac{\Gamma(a+1) \Gamma(b+1)}{(z-w)^{a+b}} \sum_{m,n,k,\ell} \binom{a}{m} \binom{b}{n} \binom{m}{k} \binom{n}{\ell} \\ &\cdot \frac{(z-w)^{m+n-k-\ell}}{(m-\ell)! (n-k)!} : (\beta(z) \gamma(w))^{m+n-k-\ell} : \end{aligned} \quad (2.124)$$

On the other hand the second way of evaluation simply gives the result

$$I_{a,b}(z, w) = \frac{\Gamma(a+b+1)}{(z-w)^{a+b}} \sum_N \binom{a+b}{N} (z-w)^N \frac{(\beta(z)\gamma(w))^N}{N!} : \quad (2.125)$$

Defining the generating functions

$$\begin{aligned} F_a(x) &= \sum_N \binom{a}{N} \frac{x^N}{N!} \\ G_{ab}(x) &= \sum_{m,n,k,\ell} \binom{a}{m} \binom{b}{n} \binom{m}{k} \binom{n}{\ell} \frac{x^{m+n-k-\ell}}{(m-\ell)!(n-k)!} \end{aligned} \quad (2.126)$$

we see that the consistency condition may be expressed as

$$G_{ab}(x) = \binom{a+b}{b} F_{a+b}(x) \quad (2.127)$$

We now briefly indicate how this identity may be proven. First we notice (see below) that we may write

$$\binom{a+b}{a} = \binom{a+b}{b} = \oint \frac{dt}{2\pi i} \frac{(1+t)^{a+b}}{t^{a+1}} \quad (2.128)$$

where the contour may be taken as the unit circle, passing through the branch point at $t = -1$ of the integrand (for suitable values of the exponents). Then we write

$$\begin{aligned} G_{ab}(x) &= \sum_{p,q,k,\ell} \binom{a}{q+\ell} \binom{b}{p+k} \binom{q+\ell}{k} \binom{p+k}{\ell} \frac{x^{p+q}}{p!q!} \\ &= \sum_{p,q,k,\ell} \prod_{i=1}^4 \oint_{C_0} \frac{dt_i}{2\pi i} \frac{1}{t_i} \frac{(1+t_1)^a}{t_1^{q+\ell}} \frac{(1+t_2)^b}{t_2^{p+k}} \frac{(1+t_3)^{q+\ell}}{t_3^k} \frac{(1+t_4)^{p+k}}{t_4^\ell} \frac{x^{p+q}}{p!q!} \end{aligned} \quad (2.129)$$

The identity is now obtained by successively doing 1) the sum over ℓ , 2) the integral over t_4 , 3) the sum over k , 4) the integral over t_3 , 5) the integral over t_1 and t_2 in any order, and 6) the sum over q with $p+q = N$. In each case one picks up residues after suitable deformations of the contours.

Let us illustrate techniques of contour deformations by proving (2.128). We consider

$$\oint_{S^1} \frac{dt}{2\pi i} (1+t)^a t^{b-1} = - \oint_{S^1} \frac{dt}{2\pi i} (1-t)^a (-t)^{b-1} \quad (2.130)$$

On the rhs we take the branch cuts to be along the positive real axis. We may then deform the contour to be one starting in 1 (where the integrand vanishes) and surrounding 0 in a way such that it 'runs' first just above and then just below the interval from 0 to 1. Compared to the negative real axis on which the integrand is positive, we pick up phases when being just above/below the positive real axis. Since the contour is always to the left of 1 we only need to consider the power of $-t$. We find

$$\begin{aligned} \oint_{S^1} \frac{dt}{2\pi i} (1+t)^a t^{b-1} &= -\frac{1}{2\pi i} \left(\int_1^0 (1-t)^a t^{b-1} e^{-i\pi(b-1)} dt + \int_0^1 (1-t)^a t^{b-1} e^{i\pi(b-1)} dt \right) \\ &= \binom{a}{-b} \end{aligned} \quad (2.131)$$

where we have used the Euler integral (2.107) and the gamma-function identity

$$\Gamma(b)\Gamma(1-b) = \frac{\pi}{s(b)} \quad (2.132)$$

with

$$s(b) = \sin(\pi b) \quad (2.133)$$

Similarly, one may show that

$$\oint_{S^1} \frac{dt}{2\pi i} (1-t)^a t^{b-1} = e^{i\pi b} \begin{pmatrix} a \\ -b \end{pmatrix} \quad (2.134)$$

Now, to make the lhs well-defined (S^1 a closed contour) we must choose the cuts as before. Letting the function on the upper sheet be real near the positive real axis between 0 and 1, we find

$$\begin{aligned} \oint_{S^1} \frac{dt}{2\pi i} (1-t)^a t^{b-1} &= \frac{1 - e^{2\pi i(b-1)}}{2\pi i} \int_1^0 (1-t)^a t^{b-1} dt \\ &= e^{i\pi b} \begin{pmatrix} a \\ -b \end{pmatrix} \end{aligned} \quad (2.135)$$

These formulas and the techniques of contour manipulations will be used repeatedly in the following chapters.

2.3 Minimal Models

Referring to the notation in (2.54) it can be shown [21] that for α_-/α_+ being rational

$$\begin{aligned} \alpha_-/\alpha_+ &= -p/q \quad , \quad c = 1 - \frac{6(p-q)^2}{pq} \\ \Delta_{n,m} &= \frac{(nq - mp)^2 - (p-q)^2}{4pq} \end{aligned} \quad (2.136)$$

where p and q are co-prime, the operator algebra becomes finite dimensional. In general, a CFT with a finite number of primary fields (wrt the chiral algebra, see below) is called a rational CFT (RCFT). The nomenclature is due to the fact that the weights and the central charge can then be shown to be rational [138, 136, 7]. The CFT's based on (2.136) are called minimal models and have multiplicities one in their spectra of primary fields. The reflection symmetry $\Delta_{n,m} = \Delta_{p-n, q-m}$ leads to the fundamental region

$$0 < n < p \quad , \quad 0 < m < q \quad , \quad p < q \quad (2.137)$$

For $q = p + 1$ one recovers the unitary subset (2.55).

In Chapter 4 we discuss Hamiltonian reduction of $SL(2)$ current algebra based on admissible representation to minimal models. There we put $p/q = t$ whereby $\alpha_+ = \sqrt{2/t}$ and $\alpha_- = -\sqrt{2t}$, while the background charge becomes $-2\alpha_0 = -\sqrt{2/t} + \sqrt{2t}$.

2.3.1 Dotsenko-Fateev Construction

Here we discuss a realization of the minimal models due to Dotsenko and Fateev [42] following ideas of Feigin and Fuks [52]. It is based on a single bosonic scalar field φ coupled to a background charge $-2\alpha_0$, as described in a preceding section

$$\begin{aligned}\varphi(z)\varphi(w) &= \ln(z-w) \\ T(z) &= \frac{1}{2} : \partial\varphi(z)\partial\varphi(z) : + \alpha_0 \partial^2\varphi(z) \\ c &= 1 - 12\alpha_0^2 \\ V_p(z) &= : e^{p\varphi(z)} : \\ \Delta_p &= \frac{1}{2}p(p - 2\alpha_0)\end{aligned}\tag{2.138}$$

V_p is a free field realization of a primary field and $\Delta(V_p) = \Delta_p = \Delta_{2\alpha_0 - p}$. We shall be interested in conformal blocks

$$\left\langle \prod_{i=1}^{\mathcal{N}} V_{p_i}(z_i) \right\rangle \tag{2.139}$$

subject to the charge conservation $\sum_{i=1}^{\mathcal{N}} p_i = 2\alpha_0$. Charges are defined wrt the (anomalous) current $j = \partial\varphi$

$$[Q, V_p(w)] = pV_p(w) \quad , \quad Q = \oint \frac{dz}{2\pi i} j(z) \tag{2.140}$$

Let us define screening currents as primary fields (vertex operators) of dimension 1. We easily find the two solutions

$$S_{\pm}(z) =: V_{\alpha_{\pm}}(z) : \quad , \quad \alpha_{\pm} = \alpha_0 \pm \sqrt{\alpha_0^2 + 2} \tag{2.141}$$

These currents are useful since their conformal properties ensure that the corresponding screening charges (integrated currents) can be inserted into correlators without violating the conformal structure. They only affect the boundary configuration or total charge, and may be inserted into (2.139) in order that the charge conservation be met.

The mode expansions read

$$\begin{aligned}\varphi(z) &= q_{\varphi} + a_0 \ln z + \sum_{n \in \mathbb{Z}} \frac{a_n}{-n} z^{-n} \\ L_n &= \frac{1}{2} \sum_{m \in \mathbb{Z}} : a_{n-m} a_m : - \alpha_0(n+1)a_n \\ [a_n, a_m] &= n\delta_{n+m,0} \quad , \quad [a_0, q_{\varphi}] = 1 \\ : e^{-p\varphi(z)} : &= \exp\left(-p \sum_{n>0} \frac{a_{-n}}{n} z^n\right) e^{-pq_{\varphi}} z^{-pa_0} \exp\left(-p \sum_{n>0} \frac{a_n}{-n} z^{-n}\right)\end{aligned}\tag{2.142}$$

and the projective ($SL(2)$) invariant ket-vacuum $|0\rangle = |sl_2\rangle$ satisfies

$$\begin{aligned}L_n|0\rangle &= 0 \quad , \quad n \geq -1 \\ a_n|0\rangle &= 0 \quad , \quad n \geq 0\end{aligned}\tag{2.143}$$

The $SL(2)$ invariant bra-vacuum $\langle sl_2|$ satisfies

$$\begin{aligned} \langle sl_2|L_n &= 0 & , & & n \leq 1 \\ \langle sl_2|a_n &= 0 & , & & n \leq -1 \end{aligned} \quad (2.144)$$

From considering $\langle sl_2|L_0$ one finds the hermiticity property

$$a_n^\dagger = a_{-n} - 2\alpha_0\delta_{n,0} \quad (2.145)$$

Let us define an $SL(2)$ non-invariant dual bra-vacuum

$$\langle 0| = \lim_{z \rightarrow \infty} \langle sl_2| : e^{2\alpha_0\varphi(z)} := \langle sl_2| e^{2\alpha_0q\varphi} \quad (2.146)$$

which has the advantage that it satisfies

$$\langle 0|a_0 = 0 \quad , \quad \langle 0|0\rangle = 1 \quad (2.147)$$

It corresponds to inserting at infinity the vertex operator $: e^{2\alpha_0\varphi(z)} :$ of conformal dimension zero. The ket-states are defined by

$$|p\rangle = \lim_{z \rightarrow 0} : e^{p\varphi(z)} : |0\rangle \quad (2.148)$$

while dual (bra-)states are defined by

$$\langle p| = \lim_{z \rightarrow \infty} \langle sl_2| : e^{(2\alpha_0-p)\varphi(z)} : z^{2\Delta_p} = \langle 0| e^{-pq\varphi} \quad (2.149)$$

Introduce intertwining fields [56, 22, 26]

$$\begin{aligned} [\phi_{p_2}(z, x)]_{p_1}^{p_3} &= \int \prod_{k=1}^n dv_k \prod_{i=1}^m dw_i \phi_{p_2}(z, x) P(v_1, \dots, v_n; w_1, \dots, w_m) \\ P(v_1, \dots, v_n; w_1, \dots, w_m) &= \prod_{k=1}^n S_+(v_k) \prod_{i=1}^m S_-(w_i) = \prod_{k=1}^n : e^{\alpha_+\varphi(v_k)} : \prod_{i=1}^m : e^{\alpha_-\varphi(w_i)} : \end{aligned} \quad (2.150)$$

with charge conservation $p_1 + p_2 - p_3 + n\alpha_+ + m\alpha_- = 0$. Now, in the $\mathcal{N}$ -point conformal block

$$W_{\mathcal{N}} = \langle p_{\mathcal{N}} | [\phi_{p_{\mathcal{N}-1}}(z_{\mathcal{N}-1})]_{\kappa_{\mathcal{N}-2}}^{p_{\mathcal{N}}} \dots [\phi_{p_j}(z_j)]_{\kappa_{j-1}}^{\kappa_j} \dots [\phi_{p_2}(z_2)]_{p_1}^{\kappa_2} | p_1 \rangle \quad (2.151)$$

we have the overall charge conservation

$$p_1 + \dots + p_{\mathcal{N}-1} - p_{\mathcal{N}} + N\alpha_+ + M\alpha_- = 0 \quad (2.152)$$

where N, M are the total numbers of screening charges inserted. The notation follows from the pictorial version

$$\begin{array}{ccccccc} & & p_{\mathcal{N}-1} & & p_j & & p_3 & p_2 \\ & & | & & | & & | & | \\ p_{\mathcal{N}} & \text{---} & \text{---} & \dots & \text{---} & \text{---} & \text{---} & \text{---} \\ & & \kappa_{\mathcal{N}-2} & & \kappa_j & & \kappa_3 & \kappa_2 & p_1 \end{array} \quad (2.153)$$

where

$$\begin{aligned}
p_1 + p_2 - \kappa_2 + n_2\alpha_+ + m_2\alpha_- &= 0 \\
\kappa_2 + p_3 - \kappa_3 + n_3\alpha_+ + m_3\alpha_- &= 0 \\
&\vdots \\
\kappa_{j-1} + p_j - \kappa_j + n_j\alpha_+ + m_j\alpha_- &= 0 \\
&\vdots \\
\kappa_{N-2} + p_{N-1} - p_N + n_{N-1}\alpha_+ + m_{N-1}\alpha_- &= 0 \\
n_2 + \dots + n_{N-1} &= N \\
m_2 + \dots + m_{N-1} &= M
\end{aligned} \tag{2.154}$$

with n_j, m_j non-negative integers. The charge conservation ensures that formal powers of ∞ arising from the contractions involving $: e^{(2\alpha_0 - p_N)\varphi(\infty)} :$, is cancelled by $\infty^{2\Delta_{p_N}}$ from (2.149). Thus we may concentrate on the remaining contractions and we find

$$\begin{aligned}
W_N &= \prod_{1 \leq j < j' < N} (z_j - z_{j'})^{p_j p_{j'}} \int \prod_{k=1}^N dv_k \prod_{i=1}^M dw_i \prod_{j,k} (z_j - v_k)^{p_j \alpha_+} \prod_{j,i} (z_j - w_i)^{p_j \alpha_-} \\
&\cdot \prod_{k < k'} (v_k - v_{k'})^{\alpha_+^2} \prod_{i < i'} (w_i - w_{i'})^{\alpha_-^2} \prod_{k,i} (v_k - w_i)^{-2}
\end{aligned} \tag{2.155}$$

For a 4-point function with all $p_i = p$, the charge conservation reads $p = -(N\alpha_+ + M\alpha_-)/2$ and leads to a discretisation or quantisation of the common conformal weights

$$\Delta_p = \frac{1}{8} ((N+1)\alpha_+ + (M+1)\alpha_-)^2 - \frac{1}{2}\alpha_0^2 \tag{2.156}$$

which re-produces the Kac table (2.54) since here $N, M \geq 0$. An alternative derivation of this quantisation also based on free fields is discussed in [93, 48]. In Chapter 4 we will use the notation

$$\alpha_{n,m} = p_{n,m} = \frac{1-n}{2}\alpha_+ + \frac{1-m}{2}\alpha_- \tag{2.157}$$

Here we will not go into a discussion of choosing contours, finding the fusion rules, the monodromy invariant Greens functions or the operator product algebra coefficients, but refer to the original works by Dotsenko and Fateev [42, 43] and by Felder [56]. However, in chapters 4 and 5 we will consider such issues in the case of a CFT based on $SL(2)$ current algebra. In that connection we shall comment on the fusion rules for minimal models.

2.3.2 A Hidden Duality

For 4-point functions in the limit $(z_1, z_2, z_3, z_4) \rightarrow (0, z, 1, \infty)$, (2.155) reduces to

$$\begin{aligned}
W_4 &= z^{p_1 p_2} (1-z)^{p_1 p_3} \int \prod_{k=1}^N dv_k \prod_{i=1}^M dw_i \prod_{k=1}^N v_k^{p_1 \alpha_+} (z - v_k)^{p_2 \alpha_+} (1 - v_k)^{p_3 \alpha_+} \prod_{k < k'} (v_k - v_{k'})^{\alpha_+^2} \\
&\cdot \prod_{i=1}^M w_i^{p_1 \alpha_-} (z - w_i)^{p_2 \alpha_-} (1 - w_i)^{p_3 \alpha_-} \prod_{i < i'} (w_i - w_{i'})^{\alpha_-^2} \prod_{k,i} (v_k - w_i)^{-2}
\end{aligned} \tag{2.158}$$

For minimal models where the momenta or charges are quantised

$$p_i = \frac{1 - n_i}{2}\alpha_+ + \frac{1 - m_i}{2}\alpha_- \quad (2.159)$$

we write the corresponding primary fields as ϕ_{n_i, m_i} . This notation allows us to state a hidden symmetry or duality in the minimal models (in fact, in any DF construction of degenerate models) as

$$\langle \prod_{i=1}^4 \phi_{n_i, m_i} \rangle \propto \langle \prod_{i=1}^4 \phi_{\hat{n}_i, \hat{m}_i} \rangle \quad (2.160)$$

where the fields are subject to the charge conservations

$$p_1 + p_2 + p_3 - p_4 + N\alpha_+ + M\alpha_- = \hat{p}_1 + \hat{p}_2 + \hat{p}_3 - \hat{p}_4 + N\alpha_+ + M\alpha_- = 0 \quad (2.161)$$

and where

$$\begin{aligned} \hat{n}_1 &= \frac{n_1 + n_2 - n_3 - n_4}{2} & , & & \hat{m}_1 &= \frac{m_1 + m_2 - m_3 - m_4}{2} \\ \hat{n}_2 &= \frac{n_1 + n_2 + n_3 + n_4}{2} & , & & \hat{m}_2 &= \frac{m_1 + m_2 + m_3 + m_4}{2} \\ \hat{n}_3 &= \frac{-n_1 + n_2 + n_3 - n_4}{2} & , & & \hat{m}_3 &= \frac{-m_1 + m_2 + m_3 - m_4}{2} \\ \hat{n}_4 &= \frac{-n_1 + n_2 - n_3 + n_4}{2} & , & & \hat{m}_4 &= \frac{-m_1 + m_2 - m_3 + m_4}{2} \end{aligned} \quad (2.162)$$

This relation is denoted 'mysterious' by Andreev [8]. Nevertheless, in the next section we will present a proof of it [112].

2.3.3 An Integral Identity

Here we want to establish the integral identity [112] underlying (2.160) in the DF constructions.

Proposition

$$\begin{aligned} & \int_0^1 \prod_{k=1}^N dv_k v_k^{a'} (1 - v_k)^{b'} (1 - zv_k)^{c'} \prod_{k < k'} (v_k - v_{k'})^{2\rho'} \\ & \cdot \prod_{i=1}^M dw_i w_i^a (1 - w_i)^b (1 - zw_i)^c \prod_{i < i'} (w_i - w_{i'})^{2\rho} \prod_{k,i}^{N,M} (v_k - w_i)^{-2} \\ & = K_{NM} \int_0^1 \prod_{k=1}^N dv_k v_k^{a' - \delta'} (1 - v_k)^{b' + \delta'} (1 - zv_k)^{c' - \delta'} \prod_{k < k'} (v_k - v_{k'})^{2\rho'} \\ & \cdot \prod_{i=1}^M dw_i w_i^{a - \delta} (1 - w_i)^{b + \delta} (1 - zw_i)^{c - \delta} \prod_{i < i'} (w_i - w_{i'})^{2\rho} \prod_{k,i}^{N,M} (v_k - w_i)^{-2} \end{aligned} \quad (2.163)$$

where

$$\begin{aligned} a' &= -\rho' a & b' &= -\rho' b & c' &= -\rho' c & \delta' &= -\rho' \delta & \rho' &= 1/\rho \\ \delta &= a + c + 1 - N + (M - 1)\rho \\ \delta' &= a' + c' + 1 - M + (N - 1)\rho' \end{aligned} \quad (2.164)$$

and

$$K_{NM} = \prod_{i=0}^{N-1} \frac{\Gamma(a' + 1 + i\rho')\Gamma(b' + 1 + i\rho')}{\Gamma(-c' + M + (-N + 1 + i)\rho')\Gamma(a' + b' + c' + 2 - M + (N - 1 + i)\rho')} \cdot \prod_{i=0}^{M-1} \frac{\Gamma(a + 1 - N + i\rho)\Gamma(b + 1 - N + i\rho)}{\Gamma(-c + (-M + 1 + i)\rho)\Gamma(a + b + c + 2 - 2N + (M - 1 + i)\rho)} \quad (2.165)$$

The left hand side of (2.163) has the structure of the standard integral realization for minimal models [42], leaving out some irrelevant pre-factors (2.158). There are N, M screening charges of the two kinds and they are at positions v_k and w_i , except they have been scaled by z , as to scale their integration regions $\int_0^z \rightarrow \int_0^1$. The fact that all integrations are like that corresponds to considering only one kind of conformal block, where all the integrations are between 0 and z [42], see also Chapter 5. The integrations are taken to be time ordered in the sense

$$T \prod_{i=1}^n \int_0^1 dt_i \sim \int_0^1 dt_1 \int_{t_1}^1 dt_2 \dots \int_{t_{n-1}}^1 dt_n \sim \frac{1}{n!} \prod_{i=1}^n \int_0^1 dt_i \quad (2.166)$$

Proof

The proof [112] we present of this identity is by brute force and takes several lengthy calculations. For the understanding of (2.160) it is interesting in it's own right. Though, we shall mostly be interested in a similar but much more complicated integral identity (5.64) between $SL(2)$ blocks. However, the techniques involved are to a large extend the same in the two cases, so let us illustrate them by going over some of the details of the present proof.

The idea is simply to consider both sides of (2.163) as functions of z , and notice that the only singularities occur when $z \rightarrow 0, 1, \infty$.

The limit $z \rightarrow 0$ is simple. In that limit both sides of (2.163) are holomorphic in z ($1 - zw \neq 0$) and we may simply put $z = 0$. Then both sides may be computed in terms of the Dotsenko-Fateev integral (4.58). This gives immediately the normalisation K_{NM} .

The limits $z \rightarrow 1, \infty$ are much more complicated. Here there will be several different power singularities of the form $(1 - z)^A$ and z^B . We must isolate those and compute their strengths and demonstrate that we get the same results for both sides of (2.163).

$z \rightarrow 1$

Now $(1 - zw)$ is close to 0 for w close to 1, so we split the integration region from 0 to 1 using a small positive ϵ as follows (the integrations are time-ordered throughout)

$$\int_0^1 = \int_0^{1-\epsilon} + \int_{1-\epsilon}^1$$

$$\int_0^1 \prod_{k=1}^N dv_k \prod_{i=1}^M dw_i = \sum_{n,m} \int_0^{1-\epsilon} \prod_{k=N-n+1}^N dv_k \prod_{i=M-m+1}^M dw_i \int_{1-\epsilon}^1 \prod_{l=1}^{N-n} dv_l \prod_{j=1}^{M-m} dw_j \quad (2.167)$$

It is not difficult to check that a particular n, m term will give rise to a particular power of $(1 - z)$. One performs the following scalings of the $\int_{1-\epsilon}^1$ integration variables, $w \sim v_l, w_j$,

$$w \rightarrow 1 - (1 - z) \frac{1 - w}{w}$$

$$\begin{aligned}
dw &\rightarrow \frac{1-z}{w^2} dw \\
1-w &\rightarrow (1-z) \frac{1-w}{w} \\
1-zw &\rightarrow (1-z) \frac{(1-z)w+z}{w} \sim \frac{1-z}{w} \\
\int_{1-\epsilon}^1 &\rightarrow \int_{\frac{1-z}{\epsilon+1-z}}^1 \sim \int_0^1
\end{aligned} \tag{2.168}$$

One rather easily finds that the power of $(1-z)$ occurring on both sides of (2.163) is

$$(1-z)^{(N-n)(b'+c'+1)+(N-n)(N-n-1)\rho'+(M-m)(b+c+1)+(M-m)(M-m-1)\rho-2(N-n)(M-m)} \tag{2.169}$$

The coefficient of this singularity may also be evaluated on both sides in terms of DF integrals. It is not, however, immediately obvious that these coefficients are equal. Both sides involves many products of terms involving ratios of gamma-functions. One employs over and over again the simple identity

$$\frac{\Gamma(X)}{\Gamma(X-L)} = \prod_{j=0}^{L-1} (X-1-j) \tag{2.170}$$

Thus, it turns out for example that there are factors on both sides involving gamma-functions with argument involving a' only (no b, c'). On the left hand side we have

$$\prod_{i=0}^{n-1} \Gamma(a' + 1 + i\rho') \tag{2.171}$$

whereas on the right hand side there are similar factors of the form

$$\frac{\prod_{i=0}^{N-1} \Gamma(a' + 1 + i\rho')}{\prod_{i=0}^{N-n-1} \Gamma(a' + 1 - M + (n+i)\rho')} \tag{2.172}$$

Using the identity (2.170) we find that the ratio between (2.171) and (2.172) is

$$\prod_{i=0}^{N-n-1} \prod_{j=0}^{M-1} \frac{1}{(a' - j + (n+i)\rho')} \tag{2.173}$$

Similarly one finds that the ratio of terms only involving a is

$$\prod_{i=0}^{N-n-1} \prod_{j=0}^{M-1} (a - n - i + j\rho) \tag{2.174}$$

Up to powers of ρ these cancel according to (2.164). By going over the several other different factors on both sides and working out the ratios, one finally shows that the product of all ratios equals 1.

This completes the proof that the singularities are identical in the limit $z \rightarrow 1$.

$z \rightarrow \infty$

The strategy is entirely analogous. Now $(1 - zw)$ is close to 0 (or indefinite) for w close to 0, so we make the split (time-ordered integrations)

$$\int_0^1 = \int_0^\epsilon + \int_\epsilon^1$$

$$\int_0^1 \prod_{k=1}^N dv_k \prod_{i=1}^M dw_i = \sum_{n,m} \int_0^\epsilon \prod_{k=N-n+1}^N dv_k \prod_{i=M-m+1}^M dw_i \int_\epsilon^1 \prod_{l=1}^{N-n} dv_l \prod_{j=1}^{M-m} dw_j \quad (2.175)$$

The $\int_0^\epsilon$ integration variables are scaled according to

$$w \rightarrow \frac{1-w}{-zw}$$

$$\int_0^\epsilon dw \rightarrow \int_1^{\frac{1}{1-z\epsilon}} \frac{dw}{zw^2} \sim (-z)^{-1} \int_0^1 \frac{dw}{w^2} \quad (2.176)$$

This time there is a subtlety in the identification of the two sides since it turns out that we must identify the left hand side with n, m with the right hand side with $N - n, M - m$. It is then simple to verify that the power of z on both sides are

$$(-z)^{-na' + (N-n)c' - n - n(n-1)\rho' - ma + (M-m)c - m - m(m-1)\rho + 2nm} \quad (2.177)$$

To check that the coefficients also agree, as before one carries out explicitly the integrations in terms of DF integrals resulting in many products of ratios of gamma-functions. Finally, one carries out the cumbersome computation showing that the ratios indeed multiply up to 1.

□

Let us remark that the trivial identity

$$F(\alpha, \beta; \gamma; z) = F(\beta, \alpha; \gamma; z) \quad (2.178)$$

for hyper-geometric functions

$$F(\alpha, \beta; \gamma; z) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\beta)} \sum_{n \geq 0} \frac{\Gamma(\alpha + n)\Gamma(\beta + n)}{\Gamma(\gamma + n)n!} z^n \quad (2.179)$$

is less transparent at the level of integral representations

$$F(\alpha, \beta; \gamma; z) = \frac{\Gamma(\gamma)}{\Gamma(\beta)\Gamma(\gamma - \beta)} \int_0^1 t^{\beta-1} (1-t)^{\gamma-\beta-1} (1-zt)^{-\alpha} dt \quad (2.180)$$

However, it is the simplest example of the integral identity (2.163) with $(N, M) = (1, 0)$ or $(N, M) = (0, 1)$. Hence, (2.163) may be viewed as a highly non-trivial generalisation of (2.178).

2.4 Affine Current Algebras

For every Cartan-Weyl generator j_a of a simple Lie algebra $\mathfrak{g}$ let there be associated a spin 1 field $J_a(z)$ (of conformal dimension $(1, 0)$). Dimensional analysis leads us to define the operator product expansion of these $(2D)$ current algebra generators to be

$$J_a(z)J_b(w) = \frac{\kappa_{ab}k}{(z-w)^2} + \frac{f_{ab}^c J_c(w)}{z-w} \quad (2.181)$$

where regular terms have been omitted. k is the central extension (see below), commuting with all currents and takes on a constant value in any representation, so we will treat it as a constant. This is analogue to the treatment of the algebra element c (the central charge) in the Virasoro algebra as a constant. $k^\nu = \frac{2k}{\theta^2}$ is the level. The Chevalley currents (generators) have OPE's

$$\begin{aligned}
H_i(z)H_j(w) &= \frac{kG_{ij}}{(z-w)^2} \\
H_i(z)E_j(w) &= \frac{A_{ij}}{z-w}E_j(w) \\
H_i(z)F_j(w) &= \frac{-A_{ij}}{z-w}F_j(w) \\
E_i(z)F_j(w) &= \frac{\frac{2k}{\alpha_i^2}\delta_{ij}}{(z-w)^2} + \frac{\delta_{ij}}{z-w}H_j(w)
\end{aligned} \tag{2.182}$$

It is wrt the Sugawara construction (originally considered in [133, 132])

$$\begin{aligned}
T(z) &= \frac{1}{\theta^2(k^\nu + h^\nu)}\kappa^{ab} : J_a J_b : (z) \\
&= \frac{1}{t} : \sum_{\alpha>0} \frac{1}{\alpha^2} (E_\alpha F_\alpha + F_\alpha E_\alpha) + \frac{1}{2} (H, H) :
\end{aligned} \tag{2.183}$$

that the currents have dimension (1,0). Here we have introduced the parameter

$$t = \frac{\theta^2}{2} (k^\nu + h^\nu) \tag{2.184}$$

where h^ν is the dual Coxeter number. The central charge is easily found to be

$$c = \frac{k^\nu d}{k^\nu + h^\nu} \tag{2.185}$$

The mode expansion reads

$$J_a(z) = \sum_{n \in \mathbb{Z}} J_a^n z^{-n-1} \tag{2.186}$$

In terms of the modes the algebra is expressed as

$$[J_a^n, J_b^m] = f_{ab}^c J_c^{n+m} + \kappa_{ab} k n \delta^{n+m,0} \tag{2.187}$$

For $n = m = 0$ one obtains a subalgebra isomorphic to $\mathfrak{g}$. It is often denoted the horizontal algebra. One obtains an affine Lie algebra (special case of a Kac-Moody algebra, see [85, 73, 18, 63, 103]) by including a derivation element D satisfying $[D, J_a^n] = n J_a^n$ and $[D, k] = 0$. In view of the fact that the currents have spin one

$$[L_n, J_a^m] = -m J_a^{n+m} \tag{2.188}$$

we may take $D = -L_0$ whenever we consider affine Lie algebras in connection with the Virasoro algebra. D plays an important role in the study of the root system of the affine Lie algebra.

Precisely as the Virasoro algebra is a central extension of the Witt algebra, (2.187) is a central extension [119, 88] of the loop algebra $\mathbf{g}_t = \mathbf{g} \otimes \mathbb{C}[t, t^{-1}]$, whose basis elements are written $j_a^n = j_a \otimes t^n = j_a(n)$ and satisfies

$$[j_a(n), j_b(m)] = [j_a, j_b](n+m) \quad (2.189)$$

The restriction from $t \in \mathbb{C} \setminus \{0\}$ to $t \in S^1$ reveals the terminology of loop algebra since then $\mathbf{g}_t$ may be viewed as the space of smooth mappings, loops, from S^1 to the Lie algebra $\mathbf{g}$.

Generalisations of the Sugawara construction is in the basis of recent developments, see [78] and references therein. It has given insight in irrational CFT.

2.4.1 Fields and States

Having introduced the affine Lie (current) algebra we are in a situation with an enlarged chiral algebra, being the semi-direct sum of the Virasoro algebra (based on the Sugawara construction) and the affine Lie algebra (with corresponding simple Lie algebra $\mathbf{g}$). By the (anti-)chiral algebra we mean the (anti-)holomorphic part of the symmetry algebra of the theory. It is natural to try to classify the field content not only wrt the Virasoro algebra but wrt the whole chiral algebra. This is part of the foundation of the study of W algebras which generally are (non-linear) extensions of the Virasoro algebra. Here we confine ourselves to the case of extending by an affine current algebra and refer to the pioneering work on W_3 algebra [142] by Zamolodchikov, some work on Hamiltonian reduction [44, 15, 24, 49], the review paper [27], the reprint collection [28] and [122]. The latter includes several details and unpublished material on Hamiltonian reduction of affine $SL(3)$ current algebra and on free field realization of W_3 algebra. In [48, 47] W_N minimal models are considered. The references mentioned are mainly concerning bosonic extensions. However, the literature includes a huge amount of material on fermionic extensions such as the $N = 1$ [121, 106, 46, 23, 62] and $N = 2$ (see e.g. [58] and references therein) superconformal algebras.

Now, let us define a primary field (multiplet) ϕ wrt the current algebra by

$$J_a(z)\phi(w) = \frac{t_a}{z-w}\phi(w) \quad (2.190)$$

Here $t_a\phi$ is a shorthand notation for the action of the matrix t_a on the field multiplet, $t_a\phi^i = (t_a)^i_j\phi^j$. Here we do not specify any further the representation that ϕ forms of the Lie algebra $\mathbf{g}$ generated by the horizontal subalgebra. Similarly, the highest weight state

$$|\lambda\rangle = \lim_{z \rightarrow 0} \phi(z)|0\rangle \quad (2.191)$$

(see (2.202) for choice of notation) provides a representation of $\{J_a^0\}$ satisfying

$$\begin{aligned} J_a^0|\lambda\rangle &= t_a|\lambda\rangle \\ J_a^n|\lambda\rangle &= 0 \quad , \quad n > 0 \end{aligned} \quad (2.192)$$

where, in addition to (2.45), the vacuum satisfies

$$J_a^n|0\rangle = 0 \quad , \quad n \geq 0 \quad (2.193)$$

Note that just like T is not a conformal primary field, J_a is not an affine primary field. The (affine) descendants are obtained by acting on the highest weight $|\lambda\rangle$ with the raising operators J_a^{-n} , $n > 0$. One may specify the highest weight state further using the triangular decomposition of $\mathfrak{g}$

$$E_\alpha^0|\lambda\rangle = 0 = J_a^n|\lambda\rangle \text{ for } n > 0, \quad H_i^0|\lambda\rangle = (\alpha_i^\nu, \lambda)|\lambda\rangle \quad (2.194)$$

Integrable (unitary) highest weight representations satisfy

$$0 \leq \lambda \cdot \theta \leq k, \quad k^\nu \in \mathbb{Z} \quad (2.195)$$

Let us end this section by briefly considering the GKO or coset construction [74]. We have a situation with two Lie algebras $\mathfrak{g}, \mathfrak{h}$ where $\mathfrak{h} \subset \mathfrak{g}$. In the corresponding affine Lie algebras we choose the generators of $\mathfrak{h}$ to be the first $\dim \mathfrak{h}$ generators among the generators J_a , $a = 1, \dots, \dim \mathfrak{g}$, of $\mathfrak{g}$. Wrt the difference $T_{g/h} = T_g - T_h$ we then have

$$T_{g/h}(z)J_a(w) = 0, \quad a = 1, \dots, \dim \mathfrak{h} \quad (2.196)$$

and

$$c_{g/h} = c_g - c_h \quad (2.197)$$

The level k_h^ν of $\mathfrak{h}$ depends on the level k_g^ν of $\mathfrak{g}$ through the Dynkin index $I_{h \subset g}$ of the embedding $\mathfrak{h} \subset \mathfrak{g}$, $k_h^\nu/k_g^\nu = I_{h \subset g}$ [129, 13, 63]. Of particular interest are the cases

$$\mathfrak{g} = (A_1)_l \oplus (A_1)_1, \quad \mathfrak{h} = (A_1)_{l+1} \quad (2.198)$$

(the out-most subscripts are the integer levels) which produces the discrete series of central charge values (2.55) in the unitary minimal models

$$c_{g/h} = \frac{3l}{l+2} + \frac{3}{1+2} - \frac{3(l+1)}{l+1+2} = 1 - \frac{6}{(l+2)(l+3)} \quad (2.199)$$

It is the GKO construction that proves the existence of unitary minimal models ($p = l + 2$). Similar constructions of the $N = 1$ minimal models are also considered in [74] while the $N = 2$ minimal models are considered in [40, 144, 25, 45].

2.4.2 The Knizhnik-Zamolodchikov Equations

The currents J_a generate the affine transformations as T generates the conformal transformations ($\epsilon(z) = \epsilon^a(z)t_a$)

$$\delta_\epsilon F(\phi(w)) = \oint \frac{dz}{2\pi i} \epsilon^a(z) J_a(z) F(\phi(w)) \quad (2.200)$$

This is sometimes referred to as the affine Ward identity. Notably, one has

$$\langle J_a(z) \phi_1(w_1) \dots \phi_N(w_N) \rangle = \sum_{j=1}^N \frac{t_a}{z - w_j} \langle \phi_1(w_1) \dots \phi_j(w_j) \dots \phi_N(w_N) \rangle \quad (2.201)$$

The index j on ϕ in ϕ_j is just a labelling. Let us now see what we learn from performing the OPE $T\phi$ using the Sugawara construction and then compare the result with the definition (2.19) of ϕ being a conformal primary field. From the double pole we find

$$\Delta(\phi) = \frac{\kappa^{ab} t_a t_b}{\theta^2(k^\nu + h^\nu)} = \frac{(\lambda, \lambda + 2\rho)}{2t} \quad (2.202)$$

where λ is the highest weight of the representation t_a , thus ϕ is characterised and sometimes labelled by this weight ($\phi = \phi_\lambda$). From the single pole we find

$$\begin{aligned}\partial\phi(w) &= \frac{\kappa^{ab}t_a}{t} : J_b(w)\phi(w) : \\ &= \frac{\kappa^{ab}t_a}{t} \oint_w \frac{dz}{2\pi i} \frac{J_b(z)\phi(w)}{z-w}\end{aligned}\tag{2.203}$$

Inserting this in an N -point function, deforming the contour into a sum of contours encircling the other fields in the correlator and using (2.201), we obtain

$$\left(t\partial_{w_j} + \sum_{i \neq j} \frac{\kappa^{ab}t_a(\lambda_j)t_b(\lambda_i)}{w_i - w_j}\right) \langle \phi_1(w_1) \dots \phi_N(w_N) \rangle = 0\tag{2.204}$$

for all w_j . Here we have specified the representations by $t_a(\lambda_i)$. This set of equations are called the Knizhnik-Zamolodchikov equations due to their first appearance in the seminal work [97]. In terms of modes (2.204) expresses the decoupling of the combined (Virasoro/affine Lie algebra) singular state

$$\left(L_{-1} - \frac{1}{t}\kappa^{ab}t_a J_b^{-1}\right) |\phi\rangle\tag{2.205}$$

Decoupling of pure affine null vectors are studied in [69]. In a subsequent section we will follow [12] and use affine singular vectors to determine the fusion rules for $SL(2)$ current algebra.

The above considerations pertain to $\mathfrak{g}$ being simple. For a semi-simple Lie algebra one simply has to add appropriate terms in various places, in particular both the Casimir in (2.202) and the second term in the bracket in (2.204) become sums of similar terms.

2.4.3 WZNW Models

A Lagrangian realization of a CFT with an affine symmetry is the Wess-Zumino-Novikov-Witten (WZNW) theory [140, 107, 141]. It is a non-linear σ -model with a compact semi-simple Lie group manifold G as target space and is based on the action

$$\begin{aligned}S = k^\nu S^\pm &= \frac{k^\nu}{4\pi} \int_\Sigma d^2z \text{Tr}(g^{-1}\partial g g^{-1}\bar{\partial} g) \\ &\pm \frac{k^\nu}{12\pi} \int_B d^3y \epsilon^{\alpha\beta\gamma} \text{Tr}(g^{-1}\partial_\alpha g g^{-1}\partial_\beta g g^{-1}\partial_\gamma g)\end{aligned}\tag{2.206}$$

where k^ν is the level of the corresponding affine Lie algebra, while $\partial B = \Sigma$. In the first term g is a map from the two dimensional manifold Σ to G , $g : \Sigma \rightarrow G$, while in the second term we use the same symbol to represent the extension to the three manifold B , $g : B \rightarrow G$. The action is conformally invariant for the ratio given of the normalisations of the two terms. Different extensions of g from Σ to B may give different values of the second term in the action, but following Witten [141] these amount to additional constants $2\pi k^\nu n$, n integer. Thus, for the path integral weight e^{iS} to be unique, k^ν is necessarily an integer [141, 30]. This is related to the second and third integer homology groups of G being $H_2(G; \mathbb{Z}) \simeq 1$ and $H_3(G; \mathbb{Z}) \simeq \mathbb{Z}$ in the cases of interest [63]. The action respects the Polyakov-Wiegman identity [118, 39]

$$S^+(gh) = S^+(g) + S^+(h) + \frac{1}{2\pi} \int_\Sigma d^2z \text{Tr}(g^{-1}\bar{\partial} g \partial h h^{-1})\tag{2.207}$$

and similarly for S^- , found by $S^-(g) = S^+(g^{-1})$. Exploring the identity, the action is seen to be invariant under

$$g(z, \bar{z}) \rightarrow \Omega(z)g(z, \bar{z})\bar{\Omega}(\bar{z}) \quad (2.208)$$

giving rise to the conserved currents

$$J = k^\nu g^{-1} \partial g \quad , \quad \bar{J} = k^\nu \bar{\partial} g g^{-1} \quad , \quad \bar{\partial} J = \partial \bar{J} = 0 \quad (2.209)$$

satisfying classical versions of (2.181). The models possess unitary highest weight representations since the level is integer. The Hilbert space of a WZNW model decomposes into a direct sum of highest weight representations [69], and due to $0 \leq \lambda \cdot \theta^\nu \leq k^\nu$, the summation over λ truncates and the WZNW model based on a compact Lie group G is a RCFT.

For non-compact Lie groups such as $SL(N, \mathbb{R})$, one is more 'sloppy' and does not restrict k^ν to be integer, since Witten's argument is no longer applicable, and the theory may be unbounded from below. However, the final quantum theory of interest still makes sense when gauging and constraining the WZNW theory [90, 91, 15] restore the boundedness of the spectrum [67, 36]. Gauged WZNW theories appear e.g. in path integral realizations of the GKO construction [68, 90] and in the G/G approach to non-critical string theory.

Chapter 3

Free Field Realizations

Since the work by Wakimoto [139] on free field realizations of affine $SL(2)$ current algebra much effort has been made in obtaining similar constructions in more general cases [50, 99, 26, 70, 82]. Many partial results are found in the literature but a general construction is still lacking. However, here we present an explicit (generalised) Wakimoto free field realization of affine Lie algebras based on simple Lie algebras [113]. The method employed is a generalisation of and a much more detailed version than the one by Awata *et al* [11]. First one finds differential operator realizations of the simple Lie algebras using Gauss decompositions, which we work out in details. This was carried out to second order in x in [11] and by similar methods to all orders in [134], though some essential parts are only presented there as recursion relations. The framework of isotopic x variables in these works is a generalisation of the situation in $SL(2)$ [143, 64, 110, 8] to any simple group. Secondly one quantises the realization by translating into free fields and then adding appropriate anomalous terms to the lowering operators [50, 99, 11]. Such terms were previously only known explicitly in the case of simple roots [82] and for $SL(n)$ [26, 50], but here explicit expressions for all roots are presented. The anomalous terms take care of multiple contractions, or in other words of the normal ordering.

As shown by Dotsenko and Fateev [42] in the case of minimal models and briefly reviewed in Chapter 2, the use of screening currents in constructing conformal blocks and Greens functions is essential. In chapters 4 and 5 we shall generalise their work to admissible representations [89] (to be introduced in Chapter 4) of affine $SL(2)$ current algebras. Thereby we find use of the screening current of the second kind originally introduced by Bershadsky and Ooguri [24]. Here we recapitulate the known results [26, 11] on screening currents of the first kind in the general case. To our knowledge, a completely satisfactory proof of their existence is lacking. We shall present some sufficient conditions which are then checked in some details. To be able to treat admissible representations of affine $SL(2)$ current algebra we shall find it necessary in chapters 4 and 5 to include also the screening currents of the second kind. Hence, we shall undertake a discussion of those in the general case. In the case of $SL(n)$ we show that they indeed exist if screening currents of the first kind exist. We also comment on a quantum group structure in the braiding-commutation algebra of the screening currents following ideas of Gomez and Sierra [75, 120, 76].

Finally, one needs explicit free field realizations of primary fields. In the framework of the x variables, such realizations are known for $SL(2)$ [64, 110] but for more general groups little is known. Here we initiate the discussion and present some partial results on $SL(n)$ and the

complete solution for $SL(3)$ and review the solution for $SL(2)$.

Throughout we choose to avoid normalising the root system of the Lie algebra by keeping the length of the highest root a free parameter.

3.1 Differential Operator Realization

Following the idea of [11] we here discuss [113] a differential operator realization of a simple Lie algebra $\mathfrak{g}$ on the polynomial ring $\mathbb{C}[x^\alpha]$, given by the following right action

$$\begin{aligned} E_\alpha(x, \partial) \langle \lambda | \mathcal{Z} &= \langle \lambda | \mathcal{Z} e_\alpha \\ H_i(x, \partial, \lambda) \langle \lambda | \mathcal{Z} &= \langle \lambda | \mathcal{Z} h_i \\ F_\alpha(x, \partial, \lambda) \langle \lambda | \mathcal{Z} &= \langle \lambda | \mathcal{Z} f_\alpha \end{aligned} \quad (3.1)$$

where $\mathcal{Z} = e^{x^\alpha e_\alpha}$ and where $\langle \lambda |$ is a lowest weight vector

$$\langle \lambda | f_\alpha = 0 \quad , \quad \langle \lambda | h_i = \langle \lambda | \lambda_i \quad , \quad \lambda_i = \langle \lambda, h_i \rangle = (\lambda, \alpha_i^\vee) \quad (3.2)$$

There is an isotopic coordinate x^α for every positive root $\alpha > 0$ and for brevity we sometimes write $x^i = x^{\alpha_i}$. Before working out the explicit form of the λ dependent first order differential operators E_α, H_i, F_α , let us briefly argue that they indeed represent a realization of $\mathfrak{g}$. The verification is based on the associativity

$$\langle \lambda | \left(\mathcal{Z} e^{s j_a} \right) e^{t j_b} = \langle \lambda | \mathcal{Z} \left(e^{s j_a} e^{t j_b} \right) \quad (3.3)$$

where a comparison of terms linear in st gives the desired commutator of the differential operators $[J_a, J_b] = f_{ab}^c J_c$. As for the affine currents we use the common notation J_a to denote the differential operators. We hope it does not lead to any confusion.

The Gauss decomposition of $\langle \lambda | \mathcal{Z} e^{t j_a}$ is (see also (3.23))

$$\begin{aligned} \langle \lambda | \mathcal{Z} \exp(t e_\alpha) &= \langle \lambda | \exp \left(x^\gamma e_\gamma + t V_\alpha^\beta(x) e_\beta + \mathcal{O}(t^2) \right) \\ &= \langle \lambda | \exp \left(t V_\alpha^\beta(x) \partial_\beta + \mathcal{O}(t^2) \right) \mathcal{Z} \\ \langle \lambda | \mathcal{Z} \exp(t h_i) &= \langle \lambda | \exp(t h_i) \exp \left(x^\gamma e_\gamma + t V_i^\beta(x) e_\beta + \mathcal{O}(t^2) \right) \\ &= \langle \lambda | \exp \left(t \left(V_i^\beta(x) \partial_\beta + \lambda_i \right) + \mathcal{O}(t^2) \right) \mathcal{Z} \\ \langle \lambda | \mathcal{Z} \exp(t f_\alpha) &= \langle \lambda | \exp \left(t Q_{-\alpha}^{-\beta}(x) f_\beta + \mathcal{O}(t^2) \right) \exp \left(t P_{-\alpha}^j(x) h_j + \mathcal{O}(t^2) \right) \\ &\quad \cdot \exp \left(x^\gamma e_\gamma + t V_{-\alpha}^\beta(x) e_\beta + \mathcal{O}(t^2) \right) \\ &= \langle \lambda | \exp \left(t \left(P_{-\alpha}^j(x) \lambda_j + V_{-\alpha}^\beta(x) \partial_\beta \right) + \mathcal{O}(t^2) \right) \mathcal{Z} \end{aligned} \quad (3.4)$$

hence

$$\begin{aligned} E_\alpha(x, \partial) &= V_\alpha^\beta(x) \partial_\beta \\ H_i(x, \partial, \lambda) &= V_i^\beta(x) \partial_\beta + \lambda_i \\ F_\alpha(x, \partial, \lambda) &= V_{-\alpha}^\beta(x) \partial_\beta + P_{-\alpha}^j(x) \lambda_j \end{aligned} \quad (3.5)$$

Here the notation $\partial_\beta = \partial_{x^\beta}$ has been introduced. Since $E_\alpha(x, \partial)$ is independent of λ it may be defined through a Gauss decomposition alone. The remaining part of this section is devoted to determine the polynomials V , P and Q .

The following versions [79, 134] of the Campbell-Baker-Hausdorff (CBH) formula will be used repeatedly

Lemma

$$\begin{aligned}
e^A e^{tB} &= \exp \left\{ A + t \sum_{p \geq 0} M_p (\text{ad}_A)^p B + \mathcal{O}(t^2) \right\} \\
e^{tB} e^A &= \exp \left\{ A + t \sum_{p \geq 0} M_p (-\text{ad}_A)^p B + \mathcal{O}(t^2) \right\} \\
e^{A+tB+tC} &= e^{tB} \exp \left\{ A + tC - t \sum_{p \geq 1} M_p (-\text{ad}_A)^p B + \mathcal{O}(t^2) \right\} \\
e^A e^{tB} &= e^{tB} \exp \left\{ A + t[A, B] + \mathcal{O}(t^2) \right\} \\
M_p &= (-1)^p \frac{B_p}{p!}
\end{aligned} \tag{3.6}$$

where the coefficients B_p are the Bernoulli numbers

$$\begin{aligned}
B(u) &= \frac{u}{e^u - 1} = \sum_{n \geq 0} \frac{B_n}{n!} u^n \\
B_{2m+1} &= 0 \quad \text{for } m \geq 1 \\
B_0 = 1, \quad B_1 &= -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_4 = -\frac{1}{30}, \dots
\end{aligned} \tag{3.7}$$

Proof

In order to solve for C the equation of the second version

$$e^{tB} e^A = e^{A+tC+\mathcal{O}(t^2)} \tag{3.8}$$

one may introduce the two-parameter group element

$$g(t, s) = e^{s(A+tC)} \tag{3.9}$$

and consider

$$\partial_s \partial_t g(t, s)|_{t=0} = Cg(0, s) + A \partial_t g(t, s)|_{t=0} \tag{3.10}$$

Furthermore, using

$$e^{-s \text{ad}_A} B = e^{-sA} B e^{sA} \tag{3.11}$$

following from

$$(-s \text{ad}_A)^N B = \sum_n \binom{N}{n} s^N (-1)^n A^n B A^{N-n} \tag{3.12}$$

which is easily shown by induction, one has

$$\partial_s \left(g(0, s)^{-1} \partial_t g(t, s)|_{t=0} \right) = e^{-s \text{ad}_A} C \tag{3.13}$$

Now consider the integral

$$e^{-\text{ad}_A} B = \int_0^1 \partial_s \left(g(0, s)^{-1} \partial_t g(t, s)|_{t=0} \right) ds = \frac{1 - e^{-\text{ad}_A}}{\text{ad}_A} C \quad (3.14)$$

From this it immediately follows that

$$C = \sum_{n \geq 0} M_n (-\text{ad}_A)^n B \quad (3.15)$$

The remaining versions of the lemma are simple rewritings of the second version.

□

In [134] the V 's are determined by an approach very similar to the one employed in the following. However, here and in [113] the Gauss decomposition (3.4) is carried out explicitly whereby one is able to determine also the P 's and Q 's. In [134] functions similar to the P 's are given by recursion relations while functions similar to the Q 's are not discussed. In [11] the polynomials V_α^β are worked out only to second order in x while only the Chevalley generators amongst the lowering operators are considered and similarly worked out to second order. As indicated in (3.4) one does not need the entire decomposition at this point but later on we will make use of it.

Proposition

$$\begin{aligned} V_\alpha^\beta(x) &= \delta_\alpha^\beta + \sum_{n \geq 1} M_n C_{\alpha; \beta_1, \dots, \beta_n}^\beta x^{\beta_1} \dots x^{\beta_n} \\ V_i^\beta(x) &= -(\alpha_i^\nu, \beta) x^\beta \\ V_{-\alpha}^\beta(x) &= \sum_{n \geq 1} \sum_{l=0}^{n-m(-\alpha, \beta_1, \dots, \beta_n)} \frac{B_l}{l!(n-l)!} C_{-\alpha; \beta_1, \dots, \beta_n}^\beta x^{\beta_1} \dots x^{\beta_n} \\ P_{-\alpha}^j(x) &= \sum_{n \geq 1} \frac{1}{n!} C_{-\alpha; \beta_1, \dots, \beta_n}^j x^{\beta_1} \dots x^{\beta_n} \\ Q_{-\alpha}^{-\beta}(x) &= \sum_{n \geq 0} \frac{1}{n!} C_{-\alpha; \beta_1, \dots, \beta_n}^{-\beta} x^{\beta_1} \dots x^{\beta_n} \end{aligned} \quad (3.16)$$

where the contracted structure constants are

$$\begin{aligned} C_{a; \beta_1, \dots, \beta_n}^b &= f_{\beta_1 a}^{a_1} f_{\beta_2 a_1}^{a_2} \dots f_{\beta_n a_{n-1}}^{a_n} \\ C_{a; \beta_1, \dots, \beta_n}^b &= \delta_a^b \quad \text{for} \quad n = 0 \end{aligned} \quad (3.17)$$

and where $m = m(-\alpha, \beta_1, \dots, \beta_n)$ is defined for a given sequence of roots $(-\alpha, \beta_1, \dots, \beta_n)$ as the minimum integer for which $-\alpha + \beta_1 + \dots + \beta_m > 0$.

Lemma

In any (formal) expansion of the form

$$\sum_{s \geq 0} \left(- \sum_{n \geq 1} b_n x^n \right)^s = \sum_{n \geq 0} a_n x^n \quad (3.18)$$

the following recursion relation is valid [113]

$$\begin{aligned} a_0 &= 1 \\ a_n &= - \sum_{l=0}^{n-1} b_{n-l} a_l \quad , \quad \text{for } n > 0 \end{aligned} \quad (3.19)$$

Proof of lemma

We have

$$\begin{aligned} \sum_{n \geq 0} a_n x^n &= \sum_{s \geq 0} \left(- \sum_{n \geq 1} b_n x^n \right)^s \\ &= 1 + \sum_{1 \leq s} (-1)^s \sum_{1 \leq i_1 < i_2 < \dots < i_s} b_{i_1} b_{i_2 - i_1} \dots b_{i_s - i_{s-1}} x^{i_s} \\ &= 1 + \sum_{0 \leq s} (-1)^{s+1} \sum_{0 = i_0 < i_1 < \dots < i_s < l} b_{i_1} b_{i_2 - i_1} \dots b_{i_s - i_{s-1}} b_{l - i_s} x^l \\ &= 1 - \sum_{1 \leq n} b_n x^n - \sum_{n \geq 2} \sum_{0 \leq s} (-1)^{s+1} \sum_{0 = i_0 < i_1 < \dots < i_s < l < n} b_{i_1} b_{i_2 - i_1} \dots b_{l - i_s} b_{n-l} x^n \\ &= 1 - b_1 x - \sum_{n \geq 2} \left(b_n + \sum_{l=1}^{n-1} b_{n-l} a_l \right) x^n \end{aligned} \quad (3.20)$$

and the lemma is easily read off.

□

Proof of proposition

Let us notice the following almost trivial relations

$$\begin{aligned} (\text{ad}_{x^\beta e_\beta})^n j_a &= C_{a; \beta_1, \dots, \beta_n}^b x^{\beta_1} \dots x^{\beta_n} j_b \\ C_{a; \beta_1, \dots, \beta_n}^b &= C_{a; \beta_1, \dots, \beta_m}^c C_{c; \beta_{m+1}, \dots, \beta_n}^b \end{aligned} \quad (3.21)$$

where $j_a \in \{e_\alpha, h_i, f_\alpha\}$. Of course one could be more general and let $j_a \in \text{span}\{e_\alpha, h_i, f_\alpha\}$ but we may concentrate on basis elements. It follows that

$$\mathcal{Z} e^{t j_a} = \exp \left\{ x^\beta e_\beta + t \sum_{n \geq 0} M_n C_{a; \beta_1, \dots, \beta_n}^b x^{\beta_1} \dots x^{\beta_n} j_b + \mathcal{O}(t^2) \right\} \quad (3.22)$$

and one may read off V_α^β in the case of $a = \alpha$. For $a = i$ one may simply use (3.6d) and thereby determine V_i^β . The last and by far the most complicated case is $a = -\alpha$ corresponding to a lowering operator $j_a = f_\alpha$. Using the CBH formula one obtains

$$\begin{aligned} \mathcal{Z} e^{t f_\alpha} &= \exp \left\{ t \sum_{l \geq 1, n_l > \dots > n_1 \geq 0} M_{n_1} M_{n_2 - n_1} \dots M_{n_l - n_{l-1}} (-1)^{n_l - n_1 + l - 1} \right. \\ &\quad \cdot C_{-\alpha; \beta_1, \dots, \beta_{n_1}}^{-\gamma_1} C_{-\gamma_1; \beta_{n_1+1}, \dots, \beta_{n_2}}^{-\gamma_2} \dots C_{-\gamma_{l-1}; \beta_{n_{l-1}+1}, \dots, \beta_{n_l}}^{-\beta} x^{\beta_1} \dots x^{\beta_{n_l}} f_\beta + \mathcal{O}(t^2) \Big\} \\ &\quad \cdot \exp \left\{ t \sum_{l \geq 1, n_l > \dots > n_1 \geq 0} M_{n_1} M_{n_2 - n_1} \dots M_{n_l - n_{l-1}} (-1)^{n_l - n_1 + l - 1} \right. \end{aligned}$$

$$\begin{aligned}
& \cdot C_{-\alpha;\beta_1,\dots,\beta_{n_1}}^{-\gamma_1} C_{-\gamma_1;\beta_{n_1+1},\dots,\beta_{n_2}}^{-\gamma_2} \dots C_{-\gamma_{l-1};\beta_{n_{l-1}+1},\dots,\beta_{n_l}}^i x^{\beta_1} \dots x^{\beta_{n_l}} h_i + \mathcal{O}(t^2) \Big\} \\
& \cdot \exp \left\{ x^\beta e_\beta + t \sum_{l \geq 1, n_l > \dots > n_1 \geq 0} M_{n_1} M_{n_2-n_1} \dots M_{n_l-n_{l-1}} (-1)^{n_l-n_1+l-1} \right. \\
& \cdot C_{-\alpha;\beta_1,\dots,\beta_{n_1}}^{-\gamma_1} C_{-\gamma_1;\beta_{n_1+1},\dots,\beta_{n_2}}^{-\gamma_2} \dots C_{-\gamma_{l-2};\beta_{n_{l-2}+1},\dots,\beta_{n_{l-1}}}^{a_-} C_{a_-;\beta_{n_{l-1}+1},\dots,\beta_{n_l}}^\beta \\
& \cdot x^{\beta_1} \dots x^{\beta_{n_l}} e_\beta + \mathcal{O}(t^2) \Big\} \\
& = \exp \left\{ t \sum_{n \geq 0} K_{n,n} C_{-\alpha;\beta_1,\dots,\beta_n}^{-\beta} x^{\beta_1} \dots x^{\beta_n} f_\beta + \mathcal{O}(t^2) \right\} \\
& \cdot \exp \left\{ t \sum_{n \geq 1} K_{n,n} C_{-\alpha;\beta_1,\dots,\beta_n}^i x^{\beta_1} \dots x^{\beta_n} h_i + \mathcal{O}(t^2) \right\} \\
& \cdot \exp \left\{ x^\beta e_\beta + t \sum_{n \geq 1} \sum_{\beta_1, \dots, \beta_n > 0} K_{n,m} C_{-\alpha;\beta_1,\dots,\beta_n}^\beta x^{\beta_1} \dots x^{\beta_n} e_\beta + \mathcal{O}(t^2) \right\} \quad (3.23)
\end{aligned}$$

where $a_- \in \{-\alpha, i\}$ and where we have introduced the coefficients

$$\begin{aligned}
K_{n,m} &= M_n + \sum_{1 \leq l, 0 \leq n_1 < \dots < n_l < m} (-1)^{n-n_1+l} M_{n_1} M_{n_2-n_1} \dots M_{n_l-n_{l-1}} M_{n-n_l} \\
&= \begin{cases} \delta_{n,1} & , \quad m = 1 \\ (-1)^n M_{n-1} + \sum_{l \geq 1, 1 < n_1 < \dots < n_l < m} (-1)^{n+l} M_{n_1-1} M_{n_2-n_1} \dots M_{n-n_l} & , \quad m > 1 \end{cases} \quad (3.24)
\end{aligned}$$

where $m = m(-\alpha, \beta_1, \dots, \beta_n)$. In the contraction of the contracted structure constants $C_{a;\beta_1,\dots,\beta_n}^b$ it has been used that the summation over c in (3.17) reduces to a summation over $-\gamma$ when $a = -\alpha$ and $b = a_-$. Furthermore, we have introduced the parameter $m = m(-\alpha, \beta_1, \dots, \beta_n)$ in order to perform the contraction over a_- in the last exponential in the Gauss decomposition. The rewriting in (3.24) is due to $(1 + (-1)^{n+1}) M_n = \delta_{n,1}$. For $m > 1$ one now has the generating function

$$\sum_{n \geq m \geq 2} K_{n,m} y^{n-m} x^n = \sum_{l > k \geq 0} (-1)^{l+1} M_l y^k x^{l+1} \sum_{s \geq 0} (1 - B(x))^s \quad (3.25)$$

In order to simplify the expression for K_{nm} we use the lemma on

$$\sum_{s \geq 0} (1 - B(x))^s = \sum_{s \geq 0} \left(- \sum_{n \geq 1} \frac{B_n}{n!} x^n \right)^s = \sum_{n \geq 0} A_n x^n \quad (3.26)$$

and find the recursion relation

$$A_n = - \sum_{l=0}^{n-1} \frac{B_{n-l}}{(n-l)!} A_l \quad , \quad A_0 = 1 \quad (3.27)$$

The solution is easily proven by induction to be

$$A_n = \frac{1}{(n+1)!} \quad (3.28)$$

The lemma also proves itself useful in considering

$$B(x) = \frac{x}{e^x - 1} = \sum_{s \geq 0} \left(- \sum_{n \geq 1} \frac{x^n}{(n+1)!} \right)^s = \sum_{n \geq 0} a_n x^n \quad (3.29)$$

which then yields

$$\sum_{l=0}^{n-1} B_l \binom{n}{l} = \begin{cases} 0 & , \quad n > 1 \\ 1 & , \quad n = 1 \end{cases} \quad (3.30)$$

Finally, from the generating function we find using (3.30) that for $m > 1$

$$K_{n,m} = \sum_{l=0}^{n-m} \frac{B_l}{l!(n-l)!} \quad (3.31)$$

while it follows immediately from (3.30) that this relation is also valid for $m = 1$.

□

In the case of a simple root $\alpha = \alpha_j$, $V_{-\alpha}^\beta(x)$ reduces to

$$V_{-\alpha_j}^\beta(x) = f_{\gamma, -\alpha_j}^\beta x^\gamma - \sum_{n \geq 1} (-1)^n M_n C_{j; \beta_1, \dots, \beta_n}^\beta x^{\alpha_j} x^{\beta_1} \dots x^{\beta_n} \quad (3.32)$$

3.1.1 Differential Screening Operators

By a left action we may define the differential operator S_α

$$\begin{aligned} \exp\{-te_\alpha\} \mathcal{Z} &= \exp\{tS_\alpha(x, \partial) + \mathcal{O}(t^2)\} \mathcal{Z} \\ S_\alpha(x, \partial) &= S_\alpha^\beta(x) \partial_\beta \end{aligned} \quad (3.33)$$

It is easily seen that

$$S_\alpha(x, \partial) = E_\alpha(-x, -\partial) \quad (3.34)$$

This indicates that

$$S_\alpha^\beta(x) = -V_\alpha^\beta(x) + f_{\gamma\alpha}^\beta x^\gamma \quad (3.35)$$

The term differential screening operator is justified by the use of the polynomials S_α^β in constructing screening currents later in this chapter. From the associativity property

$$\langle \lambda | \left(e^{-se_\alpha} \mathcal{Z} \right) e^{tj_a} = \langle \lambda | e^{-se_\alpha} \left(\mathcal{Z} e^{tj_a} \right) \quad (3.36)$$

and the Gauss decomposition (3.4), (3.23) one deduces

$$\begin{aligned} [E_\alpha(x, \partial), S_\beta(x, \partial)] &= 0 \\ [H_i(x, \partial, \lambda), S_\beta(x, \partial)] &= (\alpha_i^\nu, \beta) S_\beta(x, \partial) \\ [F_\alpha(x, \partial, \lambda), S_\beta(x, \partial)] &= P_{-\alpha}^j(x) (\alpha_j^\nu, \beta) S_\beta(x, \partial) + Q_{-\alpha}^{-\gamma}(x) (\delta_{\beta\gamma}(\beta^\nu, \lambda) - f_{\beta, -\gamma}^\sigma S_\sigma(x, \partial)) \\ [S_\alpha(x, \partial), S_\beta(x, \partial)] &= f_{\alpha\beta}^\gamma S_\gamma(x, \partial) \end{aligned} \quad (3.37)$$

The last commutator follows from the associativity of $e^{-se_\alpha} e^{-te_\beta} \mathcal{Z}$. For $\alpha = \alpha_i$ and $\beta = \alpha_j$ the commutator $[F_\alpha(x, \partial, \lambda), S_\beta(x, \partial)]$ becomes

$$[F_{\alpha_i}(x, \partial, \lambda), S_{\alpha_j}(x, \partial)] = A_{ij} x^{\alpha_i} S_{\alpha_j}(x, \partial) + \lambda_i \delta_{ij} \quad (3.38)$$

3.1.2 Classical Polynomial Identities

From the fact that E_α, H_i and F_α constitute a differential operator realization of $\mathfrak{g}$ one may easily deduce several interesting polynomial identities amongst the V 's and P 's

$$\begin{aligned}
(\alpha_i^\nu, \beta - \alpha)V_\alpha^\beta(x) &= (\alpha_i^\nu, \gamma)x^\gamma \partial_\gamma V_\alpha^\beta(x) \\
(\alpha_i^\nu, \gamma + \alpha)V_{-\alpha}^\gamma(x) &= (\alpha_i^\nu, \beta)x^\beta \partial_\beta V_{-\alpha}^\gamma(x) \\
V_\alpha^\gamma(x) \partial_\gamma V_\beta^\sigma(x) - V_\beta^\gamma(x) \partial_\gamma V_\alpha^\sigma(x) &= f_{\alpha\beta}^\gamma V_\gamma^\sigma(x) \\
V_\alpha^\gamma(x) \partial_\gamma V_{-\beta}^\sigma(x) - V_{-\beta}^\gamma(x) \partial_\gamma V_\alpha^\sigma(x) &= f_{\alpha, -\beta}^\gamma V_\gamma^\sigma(x) + f_{\alpha, -\beta}^{-\gamma} V_{-\gamma}^\sigma(x) - \delta_{\alpha\beta}(\alpha^\nu, \sigma)x^\sigma \\
V_{-\alpha}^\gamma(x) \partial_\gamma V_{-\beta}^\sigma(x) - V_{-\beta}^\gamma(x) \partial_\gamma V_{-\alpha}^\sigma(x) &= -f_{\alpha\beta}^\gamma V_{-\gamma}^\sigma(x) \\
V_\alpha^\beta(x) \partial_\beta P_{-\alpha}^j(x) &= G^{ij}(\alpha_i^\nu, \alpha^\nu) \\
V_\alpha^\gamma(x) \partial_\gamma P_{-\beta}^j(x) &= f_{\alpha, -\beta}^{-\gamma} P_{-\gamma}^j(x) \\
(\alpha_i^\nu, \beta)x^\beta \partial_\beta P_{-\alpha}^j(x) &= (\alpha_i^\nu, \alpha)P_{-\alpha}^j(x) \\
V_{-\alpha}^\gamma(x) \partial_\gamma P_{-\beta}^j(x) - V_{-\beta}^\gamma(x) \partial_\gamma P_{-\alpha}^j(x) &= -f_{\alpha\beta}^\gamma P_{-\gamma}^j(x)
\end{aligned} \tag{3.39}$$

These we will denote classical identities since they are based on realizations of Lie algebras. Additional relations determined using the Wakimoto free field realizations (of affine current algebras) will be denoted quantum identities. This terminology is justified by the point of view that the Wakimoto construction corresponds to a quantisation of the differential operator realisation. Similarly, (3.37) gives the classical identities

$$\begin{aligned}
V_\alpha^\gamma(x) \partial_\gamma S_\beta^\sigma(x) - S_\beta^\gamma(x) \partial_\gamma V_\alpha^\sigma(x) &= 0 \\
(\alpha_i^\nu, \beta - \alpha)S_\alpha^\beta(x) &= (\alpha_i^\nu, \gamma)x^\gamma \partial_\gamma S_\alpha^\beta(x) \\
V_{-\alpha}^\gamma(x) \partial_\gamma S_\beta^\sigma(x) - S_\beta^\gamma(x) \partial_\gamma V_{-\alpha}^\sigma(x) &= P_{-\alpha}^j(x)(\alpha_j^\nu, \beta)S_\beta^\sigma(x) - Q_{-\alpha}^{-\gamma}(x)f_{\beta, -\gamma}^\mu S_\mu^\sigma(x) \\
S_\alpha^\gamma(x) \partial_\gamma S_\beta^\sigma(x) - S_\beta^\gamma(x) \partial_\gamma S_\alpha^\sigma(x) &= f_{\alpha\beta}^\gamma S_\gamma^\sigma(x) \\
S_\beta^\gamma(x) \partial_\gamma P_{-\alpha}^j(x) &= -Q_{-\alpha}^{-\beta}(x)(\alpha_i^\nu, \beta^\nu)G^{ij}
\end{aligned} \tag{3.40}$$

3.2 Wakimoto Free Field Realization

The free field realization [50, 99, 11] is obtained from the differential operator realization by the substitution

$$\partial_\alpha \rightarrow \beta_\alpha(z) \quad , \quad x^\alpha \rightarrow \gamma^\alpha(z) \quad , \quad \lambda_i \rightarrow \sqrt{t} \partial \varphi_i(z) \tag{3.41}$$

and a subsequent renormalisation by adding an anomalous term, $F_\alpha^a(\gamma(z), \partial \gamma(z))$, to the lowering part. This is summarized in

Proposition

$$\begin{aligned}
E_\alpha(z) &= : V_\alpha^\beta(\gamma(z)) \beta_\beta(z) : \\
H_i(z) &= : V_i^\beta(\gamma(z)) \beta_\beta(z) : + \sqrt{t} \partial \varphi_i(z) \\
F_\alpha(z) &= : V_{-\alpha}^\beta(\gamma(z)) \beta_\beta(z) : + \sqrt{t} \partial \varphi_j(z) P_{-\alpha}^j(\gamma(z)) + F_\alpha^a(\gamma(z), \partial \gamma(z)) \\
\Delta(J_a) &= 1
\end{aligned} \tag{3.42}$$

where the anomalous part for a simple root is (this result was originally found in [82])

$$\begin{aligned} F_{\alpha_i}^a(\gamma(z), \partial\gamma(z)) &= \left(\frac{2k}{\alpha_i^2} + \frac{h^\nu \theta^2 / 2 - \alpha_i^2}{\alpha_i^2} \right) \partial\gamma^{\alpha_i}(z) \\ &= \left(\frac{k+t}{\alpha_i^2} - 1 \right) \partial\gamma^{\alpha_i}(z) \end{aligned} \quad (3.43)$$

while for a non-simple root, $\alpha = \alpha_{i_1} + \dots + \alpha_{i_n}$, $n > 1$, the anomalous part [113] is

$$\begin{aligned} & C_{\alpha_{i_1}; \alpha_{i_2}, \dots, \alpha_{i_n}}^{\alpha_{i_1} + \dots + \alpha_{i_n}} F_{\alpha_{i_1} + \dots + \alpha_{i_n}}^a(\gamma(z), \partial\gamma(z)) \\ &= (-1)^{n-1} D_n D_{n-1} \dots D_3 \left(\left(\frac{k+t}{\alpha_{i_1}^2} - 1 \right) \partial\gamma^\sigma(z) \partial_\sigma V_{-\alpha_{i_2}}^{\alpha_{i_1}}(\gamma(z)) \right) \\ &+ \sum_{m=1}^{n-1} (-1)^{n-m-1} C_{\alpha_{i_1}; \alpha_{i_2}, \dots, \alpha_{i_m}}^{\alpha_{i_1} + \dots + \alpha_{i_m}} D_n D_{n-1} \dots D_{m+2} \\ &\cdot \left(\partial\gamma^\sigma(z) \partial_\sigma \partial_\gamma V_{-\alpha_{i_{m+1}}}^\mu(\gamma(z)) \partial_\mu V_{-(\alpha_{i_1} + \dots + \alpha_{i_m})}^\gamma(\gamma(z)) \right. \\ &- \left. t G_{i_{m+1}l} \partial\gamma^{\alpha_{i_{m+1}}}(z) P_{-(\alpha_{i_1} + \dots + \alpha_{i_m})}^l(\gamma(z)) \right) \end{aligned} \quad (3.44)$$

Here we have introduced the differential operators

$$D_m = V_{-\alpha_{i_m}}^\gamma(\gamma(z)) \partial_\gamma + \partial\gamma^\sigma(z) \partial_\sigma V_{-\alpha_{i_m}}(\gamma(z)) \partial_{\partial\gamma} \quad (3.45)$$

and defined the product $D_n D_{n-1} \dots D_{n+1} = 1$, in the sense that $D_n D_{n-1} \dots D_3 = 1$ if $n = 2$, and $D_n D_{n-1} \dots D_{m+2} = 1$ for $m = n - 1$. We recall (3.17) that for $m = 1$ we have $C_{\alpha_{i_1}; \alpha_{i_2}, \dots, \alpha_{i_m}}^{\alpha_{i_1} + \dots + \alpha_{i_m}} = \delta_{\alpha_{i_1}}^{\alpha_{i_1}} = 1$. The anomalous part $F_\alpha^a(\gamma(z), \partial\gamma(z))$ is seen to be linear in $\partial\gamma$. Indeed, this is necessary for F_α to be of weight one.

Lemma

$$\begin{aligned} & : V_a^\mu(\gamma(z)) \beta_\mu(z) :: V_b^\nu(\gamma(w)) \beta_\nu(w) : \\ &= \frac{1}{z-w} : (V_a^\nu \partial_\nu V_b^\mu - V_b^\nu \partial_\nu V_a^\mu)(\gamma(w)) \beta_\mu(w) : \\ &+ \frac{-1}{(z-w)^2} \{ \partial_\mu V_a^\nu \partial_\nu V_b^\mu(\gamma(w)) + (z-w) \partial\gamma^\rho(w) \partial_\rho \partial_\mu V_a^\nu \partial_\nu V_b^\mu(\gamma(w)) \} \end{aligned} \quad (3.46)$$

The proof of this is obvious.

Proof of proposition

The strategy is straightforward, namely one performs the OPE's $J_a(z) J_b(w)$ and $T(z) J_a(w)$ using the lemma after which a comparison with (2.181), using the classical polynomial identities (3.39), reduces the problem to the set of quantum polynomial identities of the next section. One of those is a recursion relation determining the anomalous part (3.44) from $F_{\alpha_i}^a$ while the remaining ones may be seen as consistency conditions on the solution (3.44). The verification of the expression in the case of a simple root, $F_{\alpha_i}^a$, is a fairly simple matter. Since it is known [50, 99, 11] that the Wakimoto realization (3.42) exists and the anomalous part is of the form found, $F_\alpha^a(\gamma, \partial\gamma) = G_{\alpha\beta}^a(\gamma) \partial\gamma^\beta$, we may view the consistency conditions merely as interesting quantum polynomial identities. Referring to the next section for details on the solution of the

recursion relation, this completes the proof.

□

Let us emphasise that the new result in [113] over [50, 26, 70, 99, 11] is the general and explicit expression (3.44) for the anomalous part of the lowering operator in the Wakimoto realization (3.42), in addition to the explicit expressions for the polynomials V and P (and Q). As earlier stated, these were already worked out to some extent in [11, 134].

3.2.1 Quantum Polynomial Identities

The only non-trivial quantum polynomial identities, besides the already found classical identities, arise when performing OPE's involving a lowering operator F_β . We find

$$\begin{aligned}
\frac{2k}{\alpha^2} \delta_{\alpha,\beta} &= -\partial_\sigma V_\alpha^\gamma \partial_\gamma V_{-\beta}^\sigma + V_\alpha^\gamma \partial_{\partial\gamma\gamma} F_\beta^a \\
f_{\alpha,-\beta}^{-\gamma} F_\gamma^a &= -\partial\gamma^\sigma \partial_\sigma \partial_\mu V_\alpha^\gamma \partial_\gamma V_{-\beta}^\mu + V_\alpha^\gamma \partial_\gamma F_\beta^a + \partial\gamma^\sigma \partial_\sigma V_\alpha^\gamma \partial_{\partial\gamma\gamma} F_\beta^a \\
0 &= (\alpha_i^\nu, \sigma) \partial_\sigma V_{-\beta}^\sigma - (\alpha_i^\nu, \alpha) \gamma^\alpha \partial_{\partial\gamma\alpha} F_\beta^a + t G_{ij} P_{-\beta}^j \\
(\alpha_i^\nu, \beta) F_\beta^a &= (\alpha_i^\nu, \alpha) \gamma^\alpha \partial_\alpha F_\beta^a + (\alpha_i^\nu, \alpha) \partial\gamma^\alpha \partial_{\partial\gamma\alpha} F_\beta^a \\
0 &= 2(\rho, \alpha_j^\nu) P_{-\alpha}^j + \partial_\gamma V_{-\alpha}^\gamma \\
F_\alpha^a &= \partial\gamma^\gamma \partial_{\partial\gamma\gamma} F_\alpha^a \\
\partial_\gamma V_{-\alpha}^\sigma \partial_\sigma V_{-\beta}^\gamma &= t G_{ij} P_{-\alpha}^i P_{-\beta}^j + V_{-\alpha}^\gamma \partial_{\partial\gamma\gamma} F_\beta^a + V_{-\beta}^\gamma \partial_{\partial\gamma\gamma} F_\alpha^a \\
f_{\alpha\beta}^\gamma F_\gamma^a &= \partial\gamma^\sigma \partial_\sigma \partial_\gamma V_{-\alpha}^\mu \partial_\mu V_{-\beta}^\gamma - V_{-\alpha}^\gamma \partial_\gamma F_\beta^a + V_{-\beta}^\gamma \partial_\gamma F_\alpha^a \\
&\quad - t G_{ij} \partial\gamma^\sigma \partial_\sigma P_{-\alpha}^i P_{-\beta}^j - \partial\gamma^\sigma \partial_\sigma V_{-\alpha}^\gamma \partial_{\partial\gamma\gamma} F_\beta^a - V_{-\beta}^\gamma \partial\gamma^\sigma \partial_\sigma \partial_{\partial\gamma\gamma} F_\alpha^a \quad (3.47)
\end{aligned}$$

from the OPE's EF, HF, TF, FF , two identities from each. Not all the identities are independent, e.g. the second to last one follows from the last. The last one is the aforementioned recursion relation determining the anomalous part for a non-simple root. The proof of (3.44) goes as follows. In the recursion relation we expand $\gamma = \alpha_{i_1} + \dots + \alpha_{i_n}$ in simple roots and let $\alpha = \alpha_{i_n}$, whereby the two terms including F_α^a vanish. We then find

$$\begin{aligned}
f_{\alpha_{i_n}, \alpha_{i_1} + \dots + \alpha_{i_{n-1}}}^{\alpha_{i_1} + \dots + \alpha_{i_n}} F_{\alpha_{i_1} + \dots + \alpha_{i_n}}^a &= \partial\gamma^\sigma \partial_\sigma \partial_\gamma V_{-\alpha_{i_n}}^\mu \partial_\mu V_{-(\alpha_{i_1} + \dots + \alpha_{i_{n-1}})}^\gamma \\
&\quad - t G_{ijn} \partial\gamma^{\alpha_{i_n}} P_{-(\alpha_{i_1} + \dots + \alpha_{i_{n-1}})}^j - D_n F_{\alpha_{i_1} + \dots + \alpha_{i_{n-1}}}^a \quad (3.48)
\end{aligned}$$

Now the procedure is to replace the anomalous part of RHS using the recursion relation as above letting α be the 'last' simple root. In the final step where $\gamma = \alpha_{i_1} + \alpha_{i_2}$ we encounter

$$D_2 F_{\alpha_{i_1}}^a = \left(\frac{k+t}{\alpha_{i_1}^2} - 1 \right) \partial\gamma^\sigma(z) \partial_\sigma V_{-\alpha_{i_2}}^{\alpha_{i_1}}(\gamma(z)) \quad (3.49)$$

explaining the origin of the first term in (3.44).

3.3 Screening Currents

3.3.1 Screening Currents of the First Kind

The screening currents of the first kind are [26, 11]

$$s_j(z) =: S_{\alpha_j}^\alpha(\gamma(z))\beta_\alpha(z)e^{-\frac{1}{\sqrt{t}}\alpha_j\cdot\varphi(z)} : \quad (3.50)$$

and they satisfy the OPE's

Proposition

$$\begin{aligned} E_\alpha(z)s_j(w) &= 0 \\ H_i(z)s_j(w) &= 0 \\ F_{\alpha_i}(z)s_j(w) &= -\frac{2t}{\alpha_j^2}\delta_{ij}\frac{\partial}{\partial w}\left(\frac{1}{z-w} : e^{-\frac{1}{\sqrt{t}}\alpha_j\cdot\varphi(w)} : \right) \\ T(z)s_j(w) &= \frac{\partial}{\partial w}\left(\frac{1}{z-w}s_j(w)\right) \end{aligned} \quad (3.51)$$

The last OPE expresses that the screening currents have conformal weights $\Delta(s_j) = 1$. In the formalism presented here the proof of the proposition is a matter of direct verification using the classical polynomial identities (3.40). In the case of F_α for α a non-simple root it has not yet been proven that the singular part of $F_\alpha(z)s_j(w)$ is a total derivative. However, if we assume that there exists a polynomial $A(\gamma(w))$ such that

$$F_\alpha(z)s_j(w) = \frac{\partial}{\partial w}\left(\frac{1}{z-w}A(\gamma(w)) : e^{-\frac{1}{\sqrt{t}}\alpha_j\cdot\varphi(w)} : \right) \quad (3.52)$$

we find that a necessary condition is

$$F_\alpha(z)s_j(w) = -\frac{2t}{\alpha_j^2}\frac{\partial}{\partial w}\left(\frac{1}{z-w}Q_{-\alpha}^{-\alpha_j}(\gamma(w)) : e^{-\frac{1}{\sqrt{t}}\alpha_j\cdot\varphi(w)} : \right) \quad (3.53)$$

In the process we derive as sufficient conditions the following two relations

$$\begin{aligned} \partial_\gamma V_{-\alpha}^\beta \partial_\beta S_{\alpha_j}^\gamma &= 2\left(1 - \frac{t}{\alpha_j^2}\right) S_{\alpha_j}^\beta \partial_\beta P_{-\alpha}^j + S_{\alpha_j}^\gamma \partial_{\partial\gamma\gamma} F_\alpha^a \\ S_{\alpha_j}^\beta \partial_\beta F_\alpha^a &= \partial\gamma^\sigma \partial_\gamma V_{-\alpha}^\beta \partial_\beta \partial_\sigma S_{\alpha_j}^\gamma - A_{ij} \partial\gamma^\sigma \partial_\sigma S_{\alpha_j}^\beta \partial_\beta P_{-\alpha}^i - \partial\gamma^\sigma \partial_\sigma S_{\alpha_j}^\gamma \partial_{\partial\gamma\gamma} F_\alpha^a \end{aligned} \quad (3.54)$$

They have been checked explicitly for $\alpha = \alpha_{i_1} + \alpha_{i_2}$ [113]. Furthermore [113], it has been checked that for $\alpha = \beta_1 + \beta_2$ (β_i is not necessarily a simple root) the first identity follows from the recursion relation (3.47) and the assumption that (3.54) is satisfied for β_1 and β_2 . However, the very final step in the proof is still missing. The minimal conclusion is that the first identity is satisfied for α being a sum of 3 simple roots while the second identity is satisfied for α being a sum of 2 simple roots. In the case of $SL(3)$ the highest root is $\theta = \alpha_1 + \alpha_2$ so in that case (3.50) and (3.53) are seen to be true.

3.3.2 Screening Currents of the Second Kind

The only well-known screening current of the second kind is the one by Bershadsky and Ooguri for $SL(2)$ [24, 110]. In [113] a generalisation to $SL(n)$ and possibly to all simple groups (see also

[82]) is discussed. The construction of screening currents of the second kind $\tilde{s}_j$ is summarised in

Proposition

$$\tilde{s}_j(w) =: \left(S_{\alpha_j}^\beta(\gamma(w)) \beta_\beta(w) e^{-\frac{1}{\sqrt{t}} \alpha_j \cdot \varphi(w)} \right)^{-\frac{2t}{\alpha_j^2}} : \quad (3.55)$$

$$\begin{aligned} E_\alpha(z) \tilde{s}_j(w) &= 0 \\ H_i(z) \tilde{s}_j(w) &= 0 \\ F_{\alpha_i}(z) \tilde{s}_j(w) &= -\frac{2t}{\alpha_j^2} \delta_{ij} \frac{\partial}{\partial w} \left(\frac{1}{z-w} : \left(S_{\alpha_j}^\beta(\gamma(w)) \beta_\beta(w) \right)^{-\frac{2t}{\alpha_j^2}-1} e^{\sqrt{t} \alpha_j^\nu \cdot \varphi(w)} : \right) \\ T(z) \tilde{s}_j(w) &= \frac{\partial}{\partial w} \left(\frac{1}{z-w} \tilde{s}_j(w) \right) \end{aligned} \quad (3.56)$$

Furthermore

$$F_\alpha(z) \tilde{s}_j(w) = -\frac{2t}{\alpha_j^2} \frac{\partial}{\partial w} \left(\frac{1}{z-w} : Q_{-\alpha}^{-\alpha_j}(\gamma(w)) \left(S_{\alpha_j}^\beta(\gamma(w)) \beta_\beta(w) \right)^{-\frac{2t}{\alpha_j^2}-1} e^{\sqrt{t} \alpha_j^\nu \cdot \varphi(w)} : \right) \quad (3.57)$$

Proof

It is straightforward to check that $H_i(z) \tilde{s}_j(w) = 0$ and $\Delta(\tilde{s}_j) = 1$. Let us introduce the shorthand notation

$$S_j^u(z) =: \left(S_{\alpha_j}^\beta(\gamma(z)) \beta_\beta(z) \right)^u : \quad (3.58)$$

Using the explicit expressions for V_α^β and $S_{\alpha_j}^\gamma$, we find

$$\begin{aligned} E_\alpha(z) S_j^u(w) &= \sum_{l \geq 1} \frac{1}{(z-w)^l} (-1)^l \binom{u}{l} \left(: \partial_{\gamma_1} \dots \partial_{\gamma_l} V_\alpha^\beta \beta_\beta(z) S_{\alpha_j}^{\gamma_1} \dots S_{\alpha_j}^{\gamma_l} S_j^{u-l} : \right. \\ &\quad \left. - l : \partial_{\gamma_1} \dots \partial_{\gamma_{l-1}} V_\alpha^\beta(z) S_{\alpha_j}^{\gamma_1} \dots S_{\alpha_j}^{\gamma_{l-1}} \partial_\beta S_{\alpha_j}^{\gamma_l} \beta_{\gamma_l} S_j^{u-l} : \right) \end{aligned} \quad (3.59)$$

and in the case of $SL(n)$ where a simple root (here α_j) appears at most once in the decomposition of a root, all terms for $l > 1$ vanish. Hence, in that case we find

$$\begin{aligned} E_\alpha(z) S_j^u(w) &= \frac{u}{z-w} : \left(V_\alpha^\gamma \partial_\gamma S_{\alpha_j}^\beta - S_{\alpha_j}^\gamma \partial_\gamma V_\alpha^\beta \right) \beta_\beta S_j^{u-1} : \\ &= 0 \end{aligned} \quad (3.60)$$

Similarly for the lowering operator we find in the case of $SL(n)$

$$\begin{aligned} : V_{-\alpha_i}^\beta(\gamma(z)) \beta_\beta(z) : S_j^u(w) &= \frac{u A_{ij}}{z-w} : \gamma^{\alpha_i} S_j^u(w) : - \frac{u \delta_{ij}}{(z-w)^2} \frac{h^\nu \theta^2 / 2 - \alpha_i^2}{\alpha_i^2} S_j^{u-1}(w) \\ &+ \frac{u(u-1)}{(z-w)^2} \left(\frac{1}{2} : \partial_{\gamma_1} \partial_{\gamma_2} V_{-\alpha_i}^\beta(z) \beta_\beta(z) S_{\alpha_j}^{\gamma_1}(w) S_j^{\gamma_2}(w) S_j^{u-2}(w) : \right. \\ &\quad \left. - \delta_{ij} : \partial_{\gamma_1} V_{-\alpha_i}^\beta(z) S_{\alpha_j}^{\gamma_1}(w) \partial_\beta S_{\alpha_j}^{\gamma_2}(w) \beta_{\gamma_2}(w) S_j^{u-2}(w) : \right) \end{aligned} \quad (3.61)$$

Including the more simple contributions from the $P_{-\alpha_i}^l$ and $F_{\alpha_i}^a$ parts we find

$$\begin{aligned}
F_{\alpha_i}(z)\tilde{s}_j(w) &= \frac{u}{(z-w)^2} : \left((2-u)\delta_{ij}S_j^{u-1} + \frac{u-1}{2}S_{\alpha_j}^{\gamma_1}S_{\alpha_j}^{\gamma_2}\partial_{\gamma_1}\partial_{\gamma_2}V_{-\alpha_i}^\beta\beta_\beta S_j^{u-2} \right. \\
&\quad \left. - (u-1)\delta_{ij}S_{\alpha_j}^{\gamma_1}\partial_{\gamma_1}V_{-\alpha_i}^\beta\partial_\beta S_{\alpha_j}^{\gamma_2}\beta_{\gamma_2}S_j^{u-2} \right) e^{\frac{1}{\sqrt{t}}\lambda\cdot\varphi(w)} : \\
&\quad + \frac{u}{z-w} : \left(\sqrt{t}\delta_{ij}\partial\varphi_i S_j^{u-1} + \frac{u-1}{2}\partial\gamma^\alpha\partial_\alpha\partial_{\gamma_1}\partial_{\gamma_2}V_{-\alpha_j}^\beta\beta_\beta S_{\alpha_j}^{\gamma_1}S_{\alpha_j}^{\gamma_2}S_j^{u-2} \right. \\
&\quad \left. + \frac{u-1}{2}S_{\alpha_j}^{\gamma_1}S_{\alpha_j}^{\gamma_2}\partial_{\gamma_1}\partial_{\gamma_2}V_{-\alpha_i}^\beta\partial_\beta S_j^{u-2} \right. \\
&\quad \left. - (u-1)\delta_{ij}\partial\gamma^\alpha\partial_\alpha\partial_{\gamma_1}V_{-\alpha_i}^\beta S_{\alpha_j}^{\gamma_1}\partial_\beta S_{\alpha_j}^{\gamma_2}\beta_{\gamma_2}S_j^{u-2} \right) e^{\frac{1}{\sqrt{t}}\lambda\cdot\varphi(w)} : \tag{3.62}
\end{aligned}$$

where $u = -2t/\alpha_j^2$ and $\lambda = t\alpha_j^\nu$. A comparison with (3.56) yields the consistency conditions

$$\begin{aligned}
\delta_{ij}S_{\alpha_j}^\beta &= \frac{1}{2}S_{\alpha_j}^{\gamma_1}S_{\alpha_j}^{\gamma_2}\partial_{\gamma_1}\partial_{\gamma_2}V_{-\alpha_i}^\beta - \delta_{ij}S_{\alpha_j}^{\gamma_1}\partial_{\gamma_1}V_{-\alpha_i}^{\gamma_2}\partial_{\gamma_2}S_{\alpha_j}^\beta \\
\delta_{ij}\partial_\alpha S_{\alpha_j}^\beta &= \frac{1}{2}\partial_\alpha\partial_{\gamma_1}\partial_{\gamma_2}V_{-\alpha_i}^\beta S_{\alpha_j}^{\gamma_1}S_{\alpha_j}^{\gamma_2} - \delta_{ij}\partial_\alpha\partial_{\gamma_1}V_{-\alpha_i}^{\gamma_2}S_{\alpha_j}^{\gamma_1}\partial_{\gamma_2}S_{\alpha_j}^\beta \\
\delta_{ij}S_{\alpha_j}^\beta &= \frac{1}{2}S_{\alpha_j}^{\gamma_1}S_{\alpha_j}^{\gamma_2}\partial_{\gamma_1}\partial_{\gamma_2}V_{-\alpha_i}^\beta \tag{3.63}
\end{aligned}$$

which are checked using the classical polynomial identities. We may conclude that (3.55) and (3.56) work for $SL(n)$. For the more general statement in (3.57) to be satisfied in the case of $SL(n)$, the following relations are sufficient conditions

$$\begin{aligned}
Q_{-\alpha}^{-\alpha_j}S_{\alpha_j}^\beta &= \frac{1}{2}S_{\alpha_j}^{\gamma_1}S_{\alpha_j}^{\gamma_2}\partial_{\gamma_1}\partial_{\gamma_2}V_{-\alpha}^\beta - S_{\alpha_j}^\sigma\partial_\sigma V_{-\alpha}^\gamma\partial_\gamma S_{\alpha_j}^\beta \\
S_{\alpha_j}^\gamma\partial_\gamma F_\alpha^a S_{\alpha_j}^\beta &= (u-1)\partial\gamma^\sigma \left(\frac{1}{2}S_{\alpha_j}^{\gamma_1}S_{\alpha_j}^{\gamma_2}\partial_\sigma\partial_{\gamma_1}\partial_{\gamma_2}V_{-\alpha}^\beta - S_{\alpha_j}^{\gamma_1}\partial_\sigma\partial_{\gamma_1}V_{-\alpha}^{\gamma_2}\partial_{\gamma_2}S_{\alpha_j}^\beta - Q_{-\alpha}^{-\alpha_j}\partial_\sigma S_{\alpha_j}^\beta \right) \\
&\quad + \partial\gamma^\sigma \left(S_{\alpha_j}^\gamma\partial_\sigma\partial_{\gamma\gamma}F_\alpha^a - \partial_\sigma Q_{-\alpha}^{-\alpha_j} - \partial_\sigma\partial_\gamma V_{-\alpha}^\mu\partial_\mu S_{\alpha_j}^\gamma - S_{\alpha_j}^\gamma\partial_\gamma\partial_\sigma P_{-\alpha}^i\lambda_i \right) S_{\alpha_j}^\beta \\
Q_{-\alpha}^{-\alpha_j}S_{\alpha_j}^\beta &= \frac{1}{2}S_{\alpha_j}^{\gamma_1}S_{\alpha_j}^{\gamma_2}\partial_{\gamma_1}\partial_{\gamma_2}V_{-\alpha}^\beta \tag{3.64}
\end{aligned}$$

One can verify these conditions using the classical polynomial identities together with the consistency conditions (3.54). Hence, we conclude that in the case of $SL(n)$ the screening currents of the second kind (3.56) exist and are well-defined if the screening currents of the first kind (3.50) exist and are well-defined. Referring to the discussion of $\alpha = \alpha_{i_1} + \alpha_{i_2}$ in (3.54) we find that both types of screening currents exist in the case of $SL(3)$.

□

3.4 Primary Fields

The primary field $\phi(w, x)$ is defined by

$$\begin{aligned}
J_a(z)\phi(w, x) &= \frac{1}{z-w}\tilde{J}_a(x)\phi(w, x) \\
T(z)\phi(w, x) &= \frac{\Delta(\phi)}{(z-w)^2}\phi(w, x) + \frac{1}{z-w}\partial\phi(w, x) \tag{3.65}
\end{aligned}$$

where

$$\tilde{J}_a = \sigma(J_a) \quad (3.66)$$

and σ is the involution

$$\begin{aligned} \sigma(E_\alpha) &= F_\alpha \\ \sigma(H_i) &= H_i \\ \sigma(F_\alpha) &= E_\alpha \end{aligned} \quad (3.67)$$

$\Delta(\phi)$ is the conformal dimension of ϕ .

In the simple case of $SL(2)$ a primary field of spin j , $\lambda = 2j$, may be written [64, 110]

$$\phi(z, x) = (1 + x\gamma(z))^\lambda V_\lambda(z) \quad (3.68)$$

and has conformal dimension $\Delta(\phi) = \frac{1}{2t}(\lambda, \lambda + 2\rho)$. This is easily verified using

$$\begin{aligned} \tilde{E}(x) &= -xx\partial + \lambda x \\ \tilde{H}(x) &= -2x\partial + \lambda \\ \tilde{F}(x) &= \partial \\ E(z) &= \beta(z) \\ H(z) &= -2 : \gamma\beta(z) : + \sqrt{t}\partial\varphi(z) \\ F(z) &= - : \gamma\gamma\beta(z) : + \sqrt{t}\partial\varphi(z)\gamma(z) + k^\nu\partial\gamma(z) \end{aligned} \quad (3.69)$$

In general the primary field is labelled by and depends on a highest weight λ (generalising the notion of spin) in such a way that for $x = 0$ it becomes a highest weight field of weight λ , annihilated by the raising operators E_α . Highest weight fields are exactly the vertex operators V_λ and since the conformal dimension of the primary field should remain unchanged when $x \neq 0$ one should expect

$$\phi(w, x) = \phi'(\gamma(w), x) V_\lambda(w) \quad (3.70)$$

Indeed such a field is conformally primary and has conformal dimension $\Delta(\phi) = \frac{1}{2t}(\lambda, \lambda + 2\rho)$. In order to comply with (3.65) for $J_a = H_i$, we find that ϕ' is symmetric in $\gamma(w)$ and x . Due to the fact that effectively one can forget the anomalous part when contracting with ϕ' , $F_\alpha^a \phi' \sim 0$, it is then sufficient to consider the cases $J_a = E_\alpha$, and we obtain the following sufficient conditions, one for each $\alpha > 0$

$$V_\alpha^\beta(\gamma)\partial_{\gamma\beta}\phi' = V_{-\alpha}^\beta(x)\partial_{x\beta}\phi' + P_{-\alpha}^j(x)\lambda_j\phi' \quad (3.71)$$

One can say even more. By induction in addition of positive roots one can use the classical recursion relations (3.39) in V_α^β , $V_{-\alpha}^\beta$ and $P_{-\alpha}^j$ to demonstrate that if ϕ is a primary field wrt E_α and E_β , then it is a primary field wrt $f_{\alpha\beta}^\gamma E_\gamma$. This means that there are only r sufficient conditions a primary field (3.70) must satisfy:

$$V_{\alpha_i}^\beta(\gamma)\partial_{\gamma\beta}\phi' = V_{-\alpha_i}^\beta(x)\partial_{x\beta}\phi' + x^{\alpha_i}\lambda_i\phi' \quad (3.72)$$

Only in a few cases, to be discussed in the following, solutions have been found [113], though we believe that solutions exist in the general case.

3.4.1 Case of $SL(3)$

In the case of $SL(3)$ the primary field ϕ is given by

Proposition

$$\begin{aligned}
\phi(z, x) &= \prod_{j=1}^2 \left(1 + \sum_{\alpha > 0} P_{-\alpha}^j(x) P_{-\alpha}^j(\gamma(z)) \right)^{\lambda_j} V_{\lambda}(z) \\
&= \left(1 + x^1 \gamma^1(z) + \left(\frac{1}{2} x^1 x^2 + x^{12} \right) \left(\frac{1}{2} \gamma^1(z) \gamma^2(z) + \gamma^{12}(z) \right) \right)^{\lambda_1} \\
&\quad \cdot \left(1 + x^2 \gamma^2(z) + \left(\frac{1}{2} x^1 x^2 - x^{12} \right) \left(\frac{1}{2} \gamma^1(z) \gamma^2(z) - \gamma^{12}(z) \right) \right)^{\lambda_2} V_{\lambda}(z) \quad (3.73)
\end{aligned}$$

and has conformal dimension $\Delta(\phi) = \frac{1}{2t}(\lambda, \lambda + 2\rho)$. This can be checked using the relevant polynomials or the proposition (3.116) of the next section. Below we have listed all the polynomials in the case of $SL(3)$ found by reducing the general expressions of the preceding sections:

$$\begin{aligned}
V_{\alpha_1}^{\beta}(x) &= \delta_{\alpha_1}^{\beta} - \frac{1}{2} x^2 \delta_{\theta}^{\beta} \\
V_{\alpha_2}^{\beta}(x) &= \delta_{\alpha_2}^{\beta} + \frac{1}{2} x^1 \delta_{\theta}^{\beta} \\
V_1^{\beta}(x) &= -2x^1 \delta_{\alpha_1}^{\beta} + x^2 \delta_{\alpha_2}^{\beta} - x^{12} \delta_{\theta}^{\beta} \\
V_2^{\beta}(x) &= x^1 \delta_{\alpha_1}^{\beta} - 2x^2 \delta_{\alpha_2}^{\beta} - x^{12} \delta_{\theta}^{\beta} \\
V_{-\alpha_1}^{\beta}(x) &= -x^1 x^1 \delta_{\alpha_1}^{\beta} + \left(\frac{1}{2} x^1 x^2 - x^{12} \right) \delta_{\alpha_2}^{\beta} - \frac{1}{2} x^1 \left(\frac{1}{2} x^1 x^2 + x^{12} \right) \delta_{\theta}^{\beta} \\
V_{-\alpha_2}^{\beta}(x) &= \left(\frac{1}{2} x^1 x^2 + x^{12} \right) \delta_{\alpha_1}^{\beta} - x^2 x^2 \delta_{\alpha_2}^{\beta} + \frac{1}{2} x^2 \left(\frac{1}{2} x^1 x^2 - x^{12} \right) \delta_{\theta}^{\beta} \\
V_{-\theta}^{\beta}(x) &= -x^1 \left(\frac{1}{2} x^1 x^2 + x^{12} \right) \delta_{\alpha_1}^{\beta} + x^2 \left(\frac{1}{2} x^1 x^2 - x^{12} \right) \delta_{\alpha_2}^{\beta} - \left(\frac{1}{4} x^1 x^1 x^2 x^2 + x^{12} x^{12} \right) \delta_{\theta}^{\beta} \\
P_{-\theta}^j(x) &= \left(\frac{1}{2} x^1 x^2 + x^{12} \right) \delta_1^j + \left(-\frac{1}{2} x^1 x^2 + x^{12} \right) \delta_2^j \\
Q_{-\theta}^{-\beta}(x) &= x^2 \delta_{\alpha_1}^{\beta} - x^1 \delta_{\alpha_2}^{\beta} + \delta_{\theta}^{\beta} \\
F_{\theta}^a(\gamma, \partial\gamma) &= \frac{1}{2} (k^{\nu} + 1) (\partial\gamma^1 \gamma^2 - \gamma^1 \partial\gamma^2) + k^{\nu} \partial\gamma^{12} \\
S_{\alpha_1}^{\beta}(x) &= -\delta_{\alpha_1}^{\beta} - \frac{1}{2} x^2 \delta_{\theta}^{\beta} \\
S_{\alpha_2}^{\beta}(x) &= -\delta_{\alpha_2}^{\beta} + \frac{1}{2} x^1 \delta_{\theta}^{\beta} \quad (3.74)
\end{aligned}$$

in addition to the trivial ones $V_{\theta}^{\beta}(x) = \delta_{\theta}^{\beta}$, $S_{\theta}^{\beta} = -\delta_{\theta}^{\beta}$, $P_{-\alpha_i}^j(x) = x^j \delta_i^j$, $Q_{-\alpha}^{-\alpha}(x) = 1$ and $F_{\alpha_i}^a(\gamma, \partial\gamma) = (k^{\nu} + 1/2) \partial\gamma^i$.

3.4.2 Case of $SL(r+1 \geq 4)$

Let us summarise some facts on the simply-laced Lie algebras $sl(r+1) = A_r$. The positive roots are written

$$\alpha_{ij} = \alpha_i + \dots + \alpha_j \quad , \quad 1 \leq i \leq j \leq r \quad (3.75)$$

where r is the rank. The dual Coxeter number is given by $h^\nu = r + 1$ while the Weyl vector can be written

$$\rho = \frac{1}{2} \sum_{i=1}^r i(r+1-i)\alpha_i \quad (3.76)$$

The structure coefficients are found using

$$[e_{\alpha_{ij}}, e_{\alpha_{kl}}] = \delta_{k-j,1} e_{\alpha_{il}} - \delta_{i-l,1} e_{\alpha_{kj}} \quad (3.77)$$

and the standard symmetries such as $f_{\alpha,\beta}^{\alpha+\beta} = f_{\beta,-\alpha-\beta}^{-\alpha}$, valid for any simply-laced algebra. The corresponding general relation

$$\frac{f_{\alpha,\beta}^{\alpha+\beta}}{(\alpha+\beta)^2} = \frac{f_{\beta,-\alpha-\beta}^{-\alpha}}{\alpha^2} = \frac{f_{-\alpha-\beta,\alpha}^{-\beta}}{\beta^2} \quad (3.78)$$

valid for all simple groups, follow from the Jacobi identities. We find the following property of the Cartan matrix for $SL(n)$ very useful

$$(\alpha_i^\nu, \alpha_{kl}) = 0 \quad , \quad k < i < l \quad (3.79)$$

In this section we use the shorthand notations

$$D_\alpha = \partial_{\gamma^\alpha} \quad , \quad \partial_\alpha = \partial_{x^\alpha} \quad , \quad \partial_{ij} = \partial_{x^{ij}} \quad (3.80)$$

A natural generalisation of the primary fields of $SL(2)$ and $SL(3)$ is

$$\phi(w, x) \sim \prod_{j=1}^r \left(R^j(\gamma(w), x) \right)^{\lambda_j} V_\lambda(w) \quad (3.81)$$

where $R^j(\gamma, x)$ is a polynomial in γ and x . We find that (3.72) reduces to

$$V_{\alpha_i}^\beta(\gamma) D_\beta R^j(\gamma, x) = V_{-\alpha_i}^\beta(x) \partial_\beta R^j(\gamma, x) + x^j \delta_i^j R^j(\gamma, x) \quad (3.82)$$

However, as it is stated in [113], this is *not* a primary field. It is for the non-existence of solutions in the cases $2 \leq j \leq r-1$ that we will now argue using that $R^j(\gamma, x)$ can be expanded in powers of $x^\theta = x^{1r}$

$$R^j(\gamma, x) = \sum_{l=0}^n \left(x^{1r} \right)^l R_{(l)}^j(\gamma, x) \quad (3.83)$$

where $R_{(l)}^j(\gamma, x)$ is independent of x^{1r} . In the proof of this 'no-go theorem' we let j be an arbitrary but fixed integer in the interval given. Insertion of the expansion in (3.82) leads to the following set of equations

$$\begin{aligned} V_{\alpha_i}^\alpha(\gamma) \sum_{l=0}^n \left(x^{1r} \right)^l D_\alpha R_{(l)}^j(\gamma, x) &= V_{-\alpha_i}^{\alpha_{1r}}(x) \sum_{l=0}^n l \left(x^{1r} \right)^{l-1} R_{(l)}^j(\gamma, x) \\ &+ V_{-\alpha_i}^\beta(x) \sum_{l=0}^n \left(x^{1r} \right)^l \partial_\beta R_{(l)}^j(\gamma, x) \\ &+ x^j \delta_i^j \sum_{l=0}^n \left(x^{1r} \right)^l R_{(l)}^j(\gamma, x) \end{aligned} \quad (3.84)$$

Using (3.32) to explore the dependence on x^{1r} of $V_{-\alpha_i}^\beta$ we consider the powers N of x^{1r} higher or equal to n and find

$i = 1; N = n + 1:$

$$0 = (-x^{1r})(x^{1r})^n \partial_{2r} R_{(n)}^j \quad (3.85)$$

$i = r; N = n + 1:$

$$0 = x^{1r}(x^{1r})^n \partial_{1,r-1} R_{(n)}^j \quad (3.86)$$

$i = 1; N = n:$

$$V_{\alpha_1}^\alpha(\gamma)(x^{1r})^n D_\alpha R_{(n)}^j = -\frac{1}{2} x^1 x^{1r} n (x^{1r})^{n-1} R_{(n)}^j + V_{-\alpha_1}^\beta(x)(x^{1r})^n \partial_\beta R_{(n)}^j \quad (3.87)$$

$i = j; N = n:$

$$V_{\alpha_j}^\alpha(\gamma)(x^{1r})^n D_\alpha R_{(n)}^j = V_{-\alpha_j}^\beta(x)(x^{1r})^n \partial_\beta R_{(n)}^j + x^j (x^{1r})^n R_{(n)}^j \quad (3.88)$$

$i = r; N = n:$

$$V_{\alpha_r}^\alpha(\gamma)(x^{1r})^n D_\alpha R_{(n)}^j = -\frac{1}{2} x^r x^{1r} n (x^{1r})^{n-1} R_{(n)}^j + V_{-\alpha_r}^\beta(x)(x^{1r})^n \partial_\beta R_{(n)}^j \quad (3.89)$$

$i \neq 1, j, r; N = n:$

$$V_{\alpha_i}^\alpha(\gamma)(x^{1r})^n D_\alpha R_{(n)}^j = V_{-\alpha_i}^\beta(x)(x^{1r})^n \partial_\beta R_{(n)}^j \quad (3.90)$$

The last equation only exists for $r \geq 4$. To indicate the strategy in the most general case of $r \geq 4$, let us go through the case of $r = 3$ (where the only possibility is $j = 2$) and begin by summarising

$$\begin{aligned} \partial_{13} R_{(n)}^2 &= \partial_{23} R_{(n)}^2 = \partial_{12} R_{(n)}^2 = 0 \\ V_{\alpha_1}^\alpha(\gamma) D_\alpha R_{(n)}^2 &= -\frac{1}{2} n x^1 R_{(n)}^2 + V_{-\alpha_1}^\beta(x) \partial_\beta R_{(n)}^2 \\ V_{\alpha_2}^\alpha(\gamma) D_\alpha R_{(n)}^2 &= V_{-\alpha_2}^\beta(x) \partial_\beta R_{(n)}^2 + x^2 R_{(n)}^2 \\ V_{\alpha_3}^\alpha(\gamma) D_\alpha R_{(n)}^2 &= -\frac{1}{2} n x^3 R_{(n)}^2 + V_{-\alpha_3}^\beta(x) \partial_\beta R_{(n)}^2 \end{aligned} \quad (3.91)$$

Now we ask when $V_{-\alpha_1}^\beta$ for $\beta \neq \alpha_{13}, \alpha_{12}, \alpha_{23}$ includes x^{12} and find that it is only through the term

$$x^{12} f_{\alpha_{12}, -\alpha_1}^{\alpha_2} \delta_{\alpha_2}^\beta \quad (3.92)$$

Using this fact in (3.91) we find

$$\partial_2 R_{(n)}^2 = 0 \quad (3.93)$$

Similarly we ask when $V_{-\alpha_3}^\beta$ for $\beta \neq \alpha_{13}, \alpha_{12}, \alpha_{23}$ includes x^{23} and find the contribution

$$x^{23} f_{\alpha_{23}, -\alpha_3}^{\alpha_2} \delta_{\alpha_2}^\beta \quad (3.94)$$

so we may once again conclude that $\partial_2 R_{(n)}^2 = 0$. To see that $\partial_1 R_{(n)}^2 = \partial_3 R_{(n)}^2 = 0$ we consider the case $i = 2 (= j)$ which can be reduced to

$$\begin{aligned} V_{\alpha_2}^\alpha(\gamma) D_\alpha R_{(n)}^2 &= V_{-\alpha_2}^{\alpha_1}(x) \partial_1 R_{(n)}^2 + V_{-\alpha_2}^{\alpha_3}(x) \partial_3 R_{(n)}^2 + x^2 R_{(n)}^2 \\ &= \left(x^{12} f_{\alpha_{12}, -\alpha_2}^{\alpha_1} - \frac{1}{2} x^2 x^1 (\alpha_2^\nu, \alpha_1) \right) \partial_1 R_{(n)}^2 \\ &+ \left(x^{23} f_{\alpha_{23}, -\alpha_2}^{\alpha_3} - \frac{1}{2} x^2 x^3 (\alpha_2^\nu, \alpha_3) \right) \partial_3 R_{(n)}^2 \\ &+ x^2 R_{(n)}^2 \end{aligned} \quad (3.95)$$

and we immediately read off

$$\partial_1 R_{(n)}^2 = \partial_3 R_{(n)}^2 = 0 \quad (3.96)$$

We can therefore conclude that $R_{(n)}^2$ only depends on the γ 's but then according to (3.95)

$$V_{\alpha_2}^\alpha(\gamma) D_\alpha R_{(n)}^2(\gamma) = x^2 R_{(n)}^2(\gamma) \quad (3.97)$$

which means that

$$R_{(n)}^2(\gamma, x) = 0 \quad (3.98)$$

Since this holds for the highest power, n , of x^{1r} it must hold for all positive integers (or zero) less or equal to n and we have finally proven that

$$R^2(\gamma, x) = 0 \quad (3.99)$$

This means that for $\lambda_2 \neq 0$ (3.81) is not a primary field.

In the general case $r \geq 4$ we have already seen that

$$\partial_{1r} R_{(n)}^j = \partial_{2r} R_{(n)}^j = \partial_{1,r-1} R_{(n)}^j = 0 \quad (3.100)$$

For $2 \leq i < j$ (if any; j might be 2) we consider the appearance of x^{ir} in $V_{-\alpha_i}^\beta$ starting from 'the left' $i = 2, \dots, j-1$ and find in the first step

$$x^{2r} f_{\alpha_{2r}, -\alpha_2}^{\alpha_{3r}} \delta_{\alpha_{3r}}^\beta \quad (3.101)$$

so

$$\partial_{3r} R_{(n)}^j = 0 \quad (3.102)$$

We continue and conclude

$$\partial_{ir} R_{(n)}^j = 0 \quad , \quad 1 \leq i \leq j \quad (3.103)$$

where we have included the facts $\partial_{1r} R_{(n)}^j = \partial_{2r} R_{(n)}^j = 0$. From the 'other end' $i = r-1, \dots, j+1$ we consider the appearance of x^{1i} and conclude

$$\partial_{1i} R_{(n)}^j = 0 \quad , \quad j \leq i \leq r \quad (3.104)$$

Next we consider the appearance of $x^{1,r-1}$ in $V_{-\alpha_1}^\beta$:

$$x^{1,r-1} f_{\alpha_{1,r-1}, -\alpha_1}^{\alpha_{2,r-1}} \delta_{\alpha_{2,r-1}}^\beta \quad (3.105)$$

and we find

$$\partial_{2,r-1} R_{(n)}^j = 0 \quad (3.106)$$

We continue by putting $i = 2, \dots, j-1$ in $V_{-\alpha_i}^\beta$ and may then conclude that

$$\partial_{i,r-1} R_{(n)}^j = 0 \quad , \quad 1 \leq i \leq j \quad (3.107)$$

From the 'other end' $i = r, \dots, j+1$ we consider the appearance of x^{2r} and conclude

$$\partial_{2i} R_{(n)}^j = 0 \quad , \quad j \leq i \leq r \quad (3.108)$$

Then we consider $x^{1,r-2}$ in $V_{-\alpha_i}^\beta$ for $i = 1, \dots, j-1$ and x^{3r} in $V_{-\alpha_i}^\beta$ for $i = r, \dots, j+1$, and conclude

$$\partial_{i,r-2} R_{(n)}^j = 0 \quad , \quad 1 \leq i \leq j \quad (3.109)$$

and

$$\partial_{3i} R_{(n)}^j = 0 \quad , \quad j \leq i \leq r \quad (3.110)$$

Following this procedure it allows one to eliminate the dependence of $R_{(n)}^j$ on all x^α , $\alpha \geq \alpha_j$.

The procedure eliminating the remaining x 's may start by a consideration of $i = j$ and the appearance of x^{1j} . Even though $V_{-\alpha_j}^\beta$ might include x^{1j} in many terms, it will only be the term

$$x^{1j} f_{\alpha_{1j}, -\alpha_j}^{\alpha_{1,j-1}} \delta_{\alpha_{1,j-1}}^\beta \quad (3.111)$$

which will be of interest because the other terms will be for $\beta > \alpha_j$. This consideration excludes $x^{1,j-1}$ in $R_{(n)}^j$. We continue by putting $i = j-1$ and consider $x^{1,j-1}$, and so on. We conclude

$$\partial_{1i} R_{(n)}^j = 0 \quad , \quad 1 \leq i \leq j \quad (3.112)$$

Then we look at x^{2j} for $i = j$; then at $x^{2,j-1}$ for $i = j-1$ etc. We may hereby eliminate dependence on all x^α , α being to 'the left' of α_j . Similarly for the roots on 'the right' of α_j , starting by considering x^{jr} for $i = j$ etc. Finally we find that $R_{(n)}^j$ only depends on the γ 's

$$R_{(n)}^j(\gamma, x) = R_{(n)}^j(\gamma) \quad (3.113)$$

But from

$$\begin{aligned} V_{\alpha_j}^\alpha(\gamma) D_\alpha R_{(n)}^j(\gamma) &= V_{-\alpha_j}^\beta(x) \partial_\beta R_{(n)}^j(\gamma) + x^j R_{(n)}^j(\gamma) \\ &= x^j R_{(n)}^j(\gamma) \end{aligned} \quad (3.114)$$

we conclude as for the case $SL(3)$ that

$$R^j(\gamma, x) = 0 \quad , \quad 2 \leq j \leq r-1 \quad (3.115)$$

This concludes the proof of the 'no-go theorem'.

We *do* have a partial result in the general case of $SL(r+1)$, namely the proposition

Proposition

For $\lambda_2 = \dots = \lambda_{r-1} = 0$ the primary field is given by

$$\phi(w, x) = \left(1 + \sum_{\alpha > 0} P_{-\alpha}^1(\gamma(w)) P_{-\alpha}^1(x)\right)^{\lambda_1} \left(1 + \sum_{\alpha > 0} P_{-\alpha}^r(\gamma(w)) P_{-\alpha}^r(x)\right)^{\lambda_r} V_\lambda(w) \quad (3.116)$$

Proof

Firstly we note that the proposition comply with the negative result above and with the cases of $SL(2)$ and $SL(3)$. Secondly we only need to verify (3.82) for $j = 1, r$ and we find that it reduces to

$$\begin{aligned} \sum_{\alpha > 0} V_{\alpha_i}^\beta(\gamma) D_\beta P_{-\alpha}^j(\gamma) P_{-\alpha}^j(x) &= \sum_{\alpha > 0} P_{-\alpha}^j(\gamma) V_{-\alpha_i}^\beta(x) \partial_\beta P_{-\alpha}^j(x) \\ &+ x^j \delta_i^j \left(1 + \sum_{\alpha > 0} P_{-\alpha}^j(\gamma) P_{-\alpha}^j(x)\right) \end{aligned} \quad (3.117)$$

Using the classical identities (3.39) we find the sufficient conditions

$$\sum_{\alpha > 0} P_{-\alpha}^j(\gamma) \left(V_{-\alpha}^{\alpha_j}(x) + x^j P_{-\alpha}^j(x) \right) \delta_i^j = 0 \quad , \quad j = 1, r \quad (3.118)$$

which obviously are respected for $i \neq j$. Due to the observation

$$P_{-\alpha_{2,l}}^1(\gamma) = P_{-\alpha_{i,r-1}}^r(\gamma) = 0 \quad (3.119)$$

the following are sufficient conditions

$$\begin{aligned} V_{-\alpha_{1l}}^{\alpha_1}(x) &= -x^1 P_{-\alpha_{1l}}^1(x) \quad , \quad 1 \leq l \leq r \\ V_{-\alpha_{ir}}^{\alpha_r}(x) &= -x^r P_{-\alpha_{ir}}^r(x) \quad , \quad 1 \leq i \leq r \end{aligned} \quad (3.120)$$

These can be verified by induction using the recursion relations (3.39) in $V_{-\alpha}^\beta$ and $P_{-\alpha}^j$.

□

3.5 A Quantum Group Structure

In this section we shall discuss a quantum group structure in the braiding-commutation algebra of the screening currents. The original work of this kind is due to Gomez and Sierra [75] for the minimal models and was soon after carried out for affine algebras [120], but that work was only based on screening currents of the *first* kind. As we will demonstrate, using the known results [75, 120, 76] it is a simple matter to derive the quantum group structure based on *both* kinds of screening currents. For much more detailed accounts on quantum groups the reader is referred to the books [63, 32, 92, 100, 76] and references therein.

3.5.1 Hopf Algebra

A Hopf algebra A is a vector space (over the field F) endowed with the following five operations

$$\begin{aligned} M : A \times A &\rightarrow A && \text{(multiplication)} \\ \eta : F &\rightarrow A && \text{(unit map)} \\ \Delta : A &\rightarrow A \times A && \text{(co-multiplication)} \\ \epsilon : A &\rightarrow F && \text{(co-unit map)} \\ \gamma : A &\rightarrow A && \text{(antipode)} \end{aligned} \quad (3.121)$$

which possess the following five properties

$$\begin{aligned} M \circ (I \times M) &= M \circ (M \times I) && \text{(associativity)} \\ M \circ (I \times \eta) &= I = M \circ (\eta \times I) && \text{(existence of unit)} \\ (I \times \Delta) \circ \Delta &= (\Delta \times I) \circ \Delta && \text{(co-associativity)} \\ (\epsilon \times I) \circ \Delta &= I = (I \times \epsilon) \circ \Delta && \text{(existence of co-unit)} \\ M \circ (I \times \gamma) \circ \Delta &= \eta \circ \epsilon = M \circ (\gamma \times I) \circ \Delta && \text{(existence of antipode)} \end{aligned} \quad (3.122)$$

The permutation map π from $A \times A$ to $A \times A$ is introduced as

$$\pi(a \otimes b) = b \otimes a \quad (3.123)$$

Using this we define the co-multiplication Δ' by

$$\Delta' = \pi \circ \Delta \quad (3.124)$$

A Hopf algebra is called quasi-triangular if the co-multiplications Δ and Δ' are conjugate

$$\Delta(a) = R\Delta'(a)R^{-1} \quad \forall a \in A \quad (3.125)$$

where $R \in A \otimes A$ is invertible and

$$\begin{aligned} (I \times \Delta)(R) &= R_{13}R_{12} \\ (\Delta \times I)(R) &= R_{13}R_{23} \\ (\gamma \times I)(R) &= R^{-1} \end{aligned} \quad (3.126)$$

Here we mean that when we write $R = \sum_i A_i \otimes B_i$ then $R_{12} = \sum_i A_i \otimes B_i \otimes I$, $R_{13}R_{23} = \sum_{i,j} A_i \otimes A_j \otimes B_i B_j$ etc. The notion of quantum groups is usually meant to cover the set of (non-co-commutative) quasi-triangular Hopf algebras. From the physicists point of view theses are interesting for example because they provide solutions to the Yang-Baxter equation of integrable models.

3.5.2 Screening Current Realization

In [76] Gomez and Sierra have outlined the derivation of the quantum group structure in the general setting of

$$\begin{aligned} S_i(z)S_j(w) &= e^{i\pi\Omega_{ij}}S_j(w)S_i(z) \\ S_i(z)V_\lambda(w) &= e^{i\pi\Omega_{i\lambda}}V_\lambda(w)S_i(z) \\ V_\lambda(z)V_{\lambda'}(w) &= e^{i\pi\Omega_{\lambda\lambda'}}V_{\lambda'}(w)V_\lambda(z) \end{aligned} \quad (3.127)$$

where $\{S_i\}_{i=1,\dots,2r}$ denote the set of screening currents and $\{V_\lambda\}$ denote the set of vertex operators. We use the notation $S_i = s_i$ and $S_{i+r} = \tilde{s}_i$ for $1 \leq i \leq r$. According to [76] there exists a quantum group structure when (3.127) is satisfied for Ω_{ij} and $\Omega_{\lambda\lambda'}$ being symmetric. Since there is no relative braiding between β and γ we get $\beta(z)^a\gamma(w)^b = \gamma(w)^b\beta(z)^a$, hence the braiding-commutation of the screening currents will be independent of the β, γ part. Therefore in our case, one may easily deduce that

$$\begin{aligned} \Omega_{\lambda\lambda'} &= \frac{1}{t}\lambda \cdot \lambda' \\ \Omega_{i\lambda} &= \begin{cases} -\alpha_i \cdot \lambda/t & , \quad i \leq r \\ \alpha_{i-r}^\nu \cdot \lambda & , \quad i > r \end{cases} \\ \Omega_{ij} &= \begin{cases} \alpha_i \cdot \alpha_j/t & , \quad i, j \leq r \\ -\alpha_i \cdot \alpha_{j-r}^\nu & , \quad i \leq r < j \\ -\alpha_{i-r}^\nu \cdot \alpha_j & , \quad j \leq r < i \\ \alpha_{i-r}^\nu \cdot \alpha_{j-r}^\nu t & , \quad r < i, j \end{cases} \end{aligned} \quad (3.128)$$

The construction of the quantum group heavily relies on contour manipulations analogous to the ones employed in chapters 4 and 5. Let us here confine ourselves to outline the strategy [76]. First one introduces (for every z) a vector space $\mathcal{V}_{\lambda,z}$ spanned by screened vertex operators $U_{\lambda,I}(z)$, where $I = \{i_1, \dots, i_n\}$ is an ordered set of labels of screening currents

$$U_{\lambda,I}(z) = \int_{C_1} dw_1 S_{i_1}(w_1) \dots \int_{C_n} dw_n S_{i_n}(w_n) V_{\lambda}(z) \quad (3.129)$$

The contours all start at infinity, encircle counter-clockwise the branch point z once, and return to infinity. They do not intersect, nor do they cross the branch cut from z to infinity. The ordering may be chosen such that C_i lies inside C_j for $i > j$. Let us denote such contours as standard contours around z . The vertex operator itself belongs to $\mathcal{V}_{\lambda}$, namely $V_{\lambda} = U_{\lambda,\emptyset}$. It is called the 'highest weight vector', distinguished by the absence of contour structure.

For every screening current S_i we shall define a triplet of operators F_i, K_i and E_i . The former is defined as the contour creating operator

$$F_i(U_{\lambda,I}(z)) = \int_C dw S_i(w) U_{\lambda,I}(z) \quad (3.130)$$

where C is a standard contour around z and encloses the ones in $U_{\lambda,I}(z)$. Now choosing the branch cuts inherent in $\mathcal{V}_{\lambda_1}(z_1)$ and $\mathcal{V}_{\lambda_2}(z_2)$ non-intersecting, we may define the co-multiplication $\Delta(F_i)$ on $\mathcal{V}_{\lambda_1}(z_1) \otimes \mathcal{V}_{\lambda_2}(z_2)$ by

$$\begin{aligned} \Delta(F_i)(U_{\lambda_1,I_1}(z_1) \otimes U_{\lambda_2,I_2}(z_2)) &= \int_C dw S_i(w) U_{\lambda_1,I_1}(z_1) U_{\lambda_2,I_2}(z_2) \\ &= F_i(U_{\lambda_1,I_1}(z_1)) U_{\lambda_2,I_2}(z_2) \\ &+ K_i^{-1}(U_{\lambda_1,I_1}(z_1)) F_i(U_{\lambda_2,I_2}(z_2)) \end{aligned} \quad (3.131)$$

where we have introduced the operator

$$K_i(U_{\lambda,I}(z)) = e^{-i\pi \left(\sum_{i_l \in I} \Omega_{ii_l} + \Omega_{i\lambda} \right)} U_{\lambda,I}(z) \quad (3.132)$$

The contour C is of the standard type wrt the *pair* z_1 and z_2 . By an obvious contour deformation it may be written as a sum $C = C_1 + C_2$ where C_1 (C_2) is a standard contour around z_1 (z_2). Thereby we performed the rewriting in (3.131) which resulted in

$$\Delta(F_i) = F_i \otimes 1 + K_i^{-1} \otimes F_i \quad (3.133)$$

The co-multiplication of K_i is trivial

$$\Delta(K_i) = K_i \otimes K_i \quad (3.134)$$

Next we want to introduce the raising operator E_i which should be a contour annihilating operator, in some sense dual to F_i . Recall the fundamental property of the screening currents

$$\left[L_{-1}, \int_C dw S(w) \right] = \int_C dw \partial_w S(w) \quad (3.135)$$

leaving only boundary terms. Thus, let us define implicitly the operator $\hat{E}_i$ by

$$\begin{aligned} L_{-1}U_{\lambda,I}(z) &= \left[L_{-1}, \prod_{i_l \in I} \int_{C_l} dw_l S_{i_l}(w_l) \right] V_\lambda(z) + \prod_{i_l \in I} \int_{C_l} dw_l S_{i_l}(w_l) L_{-1}V_\lambda(z) \\ &= - \sum_{j \in J} (q_j - q_j^{-1}) S_j(\infty) \hat{E}_j U_{\lambda,I}(z) + \prod_{i_l \in I} F_{i_l} L_{-1}V_\lambda(z) \end{aligned} \quad (3.136)$$

The last term is a screened descendant of V_λ and J denotes the total set of different screening currents. The factor $(q_i - q_i^{-1})$ is for later convenience. Here we have introduced the deformation parameters

$$q_i = e^{i\pi\Omega_{ii}/2} \quad (3.137)$$

Let us here work out the explicit expression for the action of $\hat{E}_i$ on $U_{\lambda,I}(z)$. This computation is carried out in [76] only for all screenings being identical. We find ($n = \dim I$)

$$\begin{aligned} \left[L_{-1}, \prod_{i_l \in I} \int_{C_l} dw_l S_{i_l}(w_l) \right] V_\lambda(z) &= - \sum_{i_l \in I} (q_{i_l} - q_{i_l}^{-1}) S_{i_l}(\infty) \frac{1 - e^{2\pi i (\sum_{j>i_l} \Omega_{ij} + \Omega_{i_l\lambda})}}{q_{i_l} - q_{i_l}^{-1}} \\ &\quad \cdot e^{i\pi \sum_{j<i_l} \Omega_{ij}} \prod_{i_{l'} \in I \setminus \{i_l\}} \int_{C_{l'}} dw_{l'} S_{i_{l'}}(w_{l'}) V_\lambda(z) \end{aligned} \quad (3.138)$$

from which we obtain

$$\hat{E}_i U_{\lambda,I} = \sum_{i_l \in I, i_l \sim i} \frac{1 - e^{2\pi i (\sum_{j>i_l} \Omega_{ij} + \Omega_{i_l\lambda})}}{q_i - q_i^{-1}} e^{i\pi \sum_{j<i_l} \Omega_{ij}} U_{\lambda,I \setminus \{i_l\}} \quad (3.139)$$

Here we have used that $U_{\lambda,I \setminus \{i\}} = 0$ for $i \notin I$. The restriction in the summation means $i_l \sim i$ if $S_{i_l} = S_i$. In the special case of only one type of screening current, (3.139) reduces to ($U_{\lambda;n} = U_{\lambda,I}$)

$$\hat{E}_i U_{\lambda;n} = \frac{1 - q_i^{2(n-1)} e^{2\pi i \Omega_{i\lambda}}}{q_i - q_i^{-1}} [n]_{q_i} U_{\lambda;n-1} \quad (3.140)$$

This is in agreement with [76]. Here we have introduced

$$[a]_q = \frac{1 - q^a}{1 - q} \quad (3.141)$$

From the trivial co-multiplication of L_{-1} it is not difficult to work out that

$$\Delta(\hat{E}_i) = \hat{E}_i \otimes 1 + K_i^{-1} \otimes \hat{E}_i \quad (3.142)$$

After redefining the raising operator by $E_i = K_i \hat{E}_i$ the co-multiplication reads

$$\Delta(E_i) = E_i \otimes K_i + 1 \otimes E_i \quad (3.143)$$

With the following definitions of an anti-pode γ

$$\gamma(E_i) = -E_i K_i^{-1} \quad , \quad \gamma(K_i) = K_i^{-1} \quad , \quad \gamma(F_i) = -K_i F_i \quad (3.144)$$

and a co-unit ϵ

$$\epsilon(E_i) = 0 \quad , \quad \epsilon(K_i) = 1 \quad , \quad \epsilon(F_i) = 0 \quad (3.145)$$

it is straightforward to verify that the above is indeed a Hopf algebra. The final task is to implement quasi-triangularity in the Hopf algebra A . One defines $R = R_{\lambda_1, \lambda_2}$ to be the braiding matrix of two screened vertex operators

$$U_{\lambda_1, I_1}(z_1)U_{\lambda_2, I_2}(z_2) = \sum_{I'_1, I'_2} [R_{\lambda_1, \lambda_2}]_{I_1, I_2}^{I'_1, I'_2} U_{\lambda_2, I'_2}(z_2)U_{\lambda_1, I'_1}(z_1) \quad (3.146)$$

We refer to [76] for details.

Let us summarise the commutation relations, which are not difficult to determine

$$\begin{aligned} K_i K_j &= K_j K_i \\ K_i E_j &= e^{i\pi\Omega_{ij}} E_j K_i \\ K_i F_j &= e^{-i\pi\Omega_{ij}} F_j K_i \\ [E_i, F_j] &= \delta_{ij} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}} \end{aligned} \quad (3.147)$$

and review the defining (Chevalley) commutation relations for the q deformed enveloping algebra $U_q(\mathfrak{g})$ of a simple Lie algebras $\mathfrak{g}$

$$\begin{aligned} k_i k_j &= k_j k_i \\ k_i e_j &= q_i^{A_{ij}} e_j k_i \\ k_i f_j &= q_i^{-A_{ij}} f_j k_i \\ [e_i, f_j] &= \delta_{ij} \frac{k_i - k_i^{-1}}{q_i - q_i^{-1}} \end{aligned} \quad (3.148)$$

where $q_i = q^{D_i}$. $D = \text{diag}(D_i)$ is a diagonal matrix symmetrising the Cartan matrix, $D_i A_{ij} = D_j A_{ji}$. In addition there are generalised (or quantum) Serre relations which will not concern us here.

In our case we find the deformation parameters (3.137) to be

$$q_j = e^{i\pi\Omega_{jj}/2} = \begin{cases} e^{i\pi\alpha_j^2/2t} & , \quad j \leq r \\ e^{i\pi(\alpha_{j-r}^\vee)^2 t/2} & , \quad j > r \end{cases} \quad (3.149)$$

We immediately see that each screening current S_i , $i = 1, \dots, 2r$, gives rise to a subalgebra $\mathcal{U}_{q_i}(sl(2))$ generated by $\{E_i, K_i, F_i\}$ with q_i given by (3.149). Thus, the total quantum group A may be viewed as a semi-direct sum of these $2r$ subalgebras. These are not the only subalgebras. Both *kinds* of screening currents also give rise to a subalgebra each. One observes that with

$$q = e^{i\pi/t} \quad , \quad D_i = \alpha_i^2/2 \quad (3.150)$$

the screening currents of the *first* kind (S_i , $i \leq r$) give rise to the subalgebra $\mathcal{U}_q(\mathfrak{g})$, due to

$$(q^{D_i})^{\pm A_{ij}} = e^{\pm i\pi\Omega_{ij}} \quad (3.151)$$

Similarly, the screening currents of the *second* kind (S_i , $i > r$) with

$$\tilde{q} = e^{i\pi t} \quad , \quad \tilde{D}_i = \left(\alpha_{i-r}^\nu \right)^2 / 2 = 2 / \alpha_{i-r}^2 \quad (3.152)$$

give rise to the subalgebra $\mathcal{U}_{\tilde{q}}(\mathbf{g}^t)$, due to

$$\left(\tilde{q}^{\tilde{D}_i} \right)^{\pm A_{j-r, i-r}} = e^{\pm i\pi \Omega_{ij}} \quad (3.153)$$

Here $\mathbf{g}^t$ is the dual Lie algebra to $\mathbf{g}$, obtained by transposing the Cartan matrix $A_{ij} \rightarrow A_{ij}^t = A_{ji}$. Hence, alternatively one may view A as the semi-direct sum

$$A = \mathcal{U}_q(\mathbf{g}) \oplus_{\text{semi}} \mathcal{U}_{\tilde{q}}(\mathbf{g}^t) \quad (3.154)$$

That the sum is only semi-direct is due to the fact that Ω_{ij} is not a block matrix. For admissible representations (see Chapter 4) t is rational and the deformation parameters q and $\tilde{q}$ are roots of unity.

In the discussion on primary fields it was argued that they are products of vertex operators and functions of γ . This means that one may generalise the construction above from vertex operators of weight λ to primary fields labelled by λ .

Chapter 4

Chiral Blocks in $SL(2)$ Current Algebra

N -point correlators of two dimensional conformal WZNW theories based on affine $SL(2)$ current algebra have been much studied already. They are typically constructed either by applying the free field realization of Wakimoto [139], from which results have been given for example in [22, 11, 64, 41], or by solving the Knizhnik-Zamolodchikov (KZ) equations [97], from which results have been given for example in [97, 143, 6, 34, 127, 64, 54]. Recently the structure of solutions of the KZ equations on higher genus Riemann surfaces has been examined [57]. The results given in these various pieces of works are quite complete as far as unitary, integrable representations [69] are concerned, but appear surprisingly incomplete for the general case, including admissible representations [89], see below. In general the WZNW theory (for $\widehat{sl}(2)_k$) is characterised by the level k or equivalently by $t = k + 2$. Then degenerate primary fields exist for representations characterised by spins $j_{r,s}$ given by [87, 101]

$$2j_{r,s} + 1 = r - st \tag{4.1}$$

with r, s integers. However, previous applications of the free field Wakimoto realization are complete only for the case $s = 0$ which is the full case only for integrable representations. The reason for this restriction is fairly natural, since the screening charge usually employed in the free field realization is capable of screening just such primary fields. In fact, a possible second screening operator capable of screening the general case was proposed by Bershadsky and Ooguri [24], but since it involved fractional powers of the free ghost fields, discussions on its interpretation have been only partly successful [41, 64].

In Chapter 2 we outlined how the techniques of fractional calculus [96, 102, 124, 126] naturally provide a solution. As a result we are able to render the free field Wakimoto realization applicable in a straightforward way, and to present how it leads to general integral formulas for the N -point chiral blocks on the sphere. In the process we introduce auxiliary integration variables. We shall check that the blocks provide exact solutions to the KZ equations. This is merely a check of the procedure. The real merit, we believe, is that we manage to render the free field realization straightforward in cases where at first sight it appears very difficult to use. It appears from comparing with known solutions in the mathematics literature [127, 54], that our formulas represent fairly powerful ways of dealing with such solutions. We briefly explain why the solutions to the KZ equations considered in [54] constitute a class having only little overlap with ours and discuss a possible hybrid treatment. Furthermore, we discuss the slightly non-trivial way in which projective invariances of our correlators are established.

Utilising a freedom in the choice of numbers of screenings needed, we introduce the notion of over-screening. In the case of 3-point functions based on our free field realization, we are then able to re-derive the fusion rules for admissible representations. In [12] the fusion rules are determined from considering decoupling of singular vectors as they are described in [101]. Shortly after the same two sets of fusion rules were obtained by cohomological methods [55]. Only the first set corresponds to a direct generalisation of the situation in integrable representations. Our derivation of the fusion rules brings in the question of choosing contours for the 3-point functions, and we find that a peculiar mix of known contours is needed. The discussion of choosing contours for 4-point blocks is postponed to Chapter 5.

The relation between the $SL(2)$ current algebra and the Virasoro algebra via Hamiltonian reduction is well-known [20, 117]. In particular Bershadsky and Ooguri [24] have used the powerful BRST formalism for the reduction to establish equivalence between on the one hand $\widehat{sl}(2)_k$ WZNW theory after reduction, and on the other hand conformal minimal theory labelled by (p, q) , provided $k + 2 = p/q$. Here admissible representations are used in the WZNW case, see also the work [50] by Feigin and Frenkel. This equivalence is discussed in those references at the level of the algebra and of the BRST cohomology of physical states. A particularly simple and remarkable realization of these ideas has been discussed by Furlan, Ganchev, Paunov and Petkova [64] at the level of N -point blocks on the sphere. Prior to our work [110, 111], the most detailed analysis of N -point functions has not been in terms of the free field Wakimoto realization. Rather it has been in terms of studying solutions to the KZ equations. Thus Furlan, Ganchev, Paunov and Petkova [64] presented a systematic approach whereby one makes use of representations of primary $SL(2)$ fields based on two variables (z, x) as described in the previous chapter, and utilised in our approach too. The aforementioned interesting proposal checked in many explicit examples [64], is that minimal model conformal blocks are obtained from the affine $SL(2)$ blocks by a simple substitution, one of identifying the x_i variables related to the i 'th $SL(2)$ representation, with the Koba-Nielsen variables z_i . Based on this, solutions of the KZ equations are written as power series of $(x_i - z_i)$ [64], and by construction those solutions are selected for which the boundary conditions are, that $x_i - z_i = 0$ reproduces the corresponding minimal model correlator. The expansion coefficients are given in terms of recurrence relations¹. To make sure that such a solution of the KZ equations really generates the WZNW correlator (up to normalisation), a study is performed in [64] of the null vector decoupling that follows from that solution, and whether that is as expected for a WZNW correlator. Although this has been checked in many examples [64] no explicit general proof has been provided. The relation between null vector decoupling in WZNW correlators and minimal model Virasoro correlators has been discussed for example in [66].

Using our free field realization, the issue of whether the $x_i \rightarrow z_i$ limit gives rise to correlators in minimal models may be addressed. We shall provide a simple direct proof of this formulation of Hamiltonian reduction and evaluate a normalisation constant which occurs. We find that the result is obtained in general just by requiring that the x_i 's are *proportional* to the z_i 's, independent of the common factor of proportionality. This is reasonable because such a proportionality constant would depend on normalisations of the fields. Furthermore, we shall verify that our correlators based on free field realizations indeed respect the proposed formulation of Hamiltonian reduction. Finally we compare with more standard forms of Hamiltonian

¹ In fact, it appears that the sums may be explicitly performed, see the very recent work [65].

reduction.

4.1 Dual States

Before considering correlators let us define our notations and choices as far as dual states are concerned [60, 41, 110]. We are only going to consider the group $SL(2)$. To comply with the notation in [110, 111, 112] we will not use the conventions of Chapter 3 on free fields but make the substitutions

$$\begin{aligned}
J^+ &= E \\
J^3 &= \frac{1}{2}H \\
J^- &= F \\
k &= k^\nu \\
\varphi &= -\sqrt{\frac{\theta^2}{4}}\varphi_1 \\
t &= \frac{2}{\theta^2}t \\
j &= \frac{1}{2}\lambda_1
\end{aligned} \tag{4.2}$$

where the rhs refers to Chapter 3. This results in the following realizations supplemented by the introduction of the following notation

$$\begin{aligned}
\sigma(J^+)(x) &= D_x^+ = -x^2\partial_x + 2xj \\
\sigma(J^3)(x) &= D_x^3 = -x\partial_x + j \\
\sigma(J^-)(x) &= D_x^- = \partial_x \\
J^+(z) &= \beta(z) \\
J^3(z) &= -:\gamma\beta:(z) - \sqrt{t/2}\partial\varphi(z) \\
J^-(z) &= -:\gamma^2\beta:(z) + k\partial\gamma(z) - \sqrt{2t}\gamma\partial\varphi(z) \\
T(z) &= :\beta\partial\gamma:(z) + \frac{1}{2}:\partial\varphi\partial\varphi:(z) + \frac{1}{\sqrt{2t}}\partial^2\varphi(z) \\
c &= \frac{3k}{k+2} \\
s(z) &= \beta(z)e^{\sqrt{2/t}\varphi(z)} \\
\tilde{s}(z) &= \beta(z)^{-t}e^{-\sqrt{2t}\varphi(z)} \\
\phi_j(z) &= (1 + \gamma(z)x)^{2j} : e^{-j\sqrt{2/t}\varphi(z)} :
\end{aligned} \tag{4.3}$$

where ϕ_j is a primary field with spin j and (β, γ) is a bosonic ghost pair of conformal weights $(1,0)$. Also the phases of the screening currents have been changed. Furthermore we have the OPE's

$$\varphi(z)\varphi(w) = \ln(z-w)$$

$$\begin{aligned}
\beta(z)\gamma(w) &= \frac{1}{z-w} \\
J^+(z)J^-(w) &= \frac{k}{(z-w)^2} + \frac{2}{z-w}J^3(w) \\
J^3(z)J^\pm(w) &= \pm \frac{1}{z-w}J^\pm(w) \\
J^3(z)J^3(w) &= \frac{k/2}{(z-w)^2}
\end{aligned} \tag{4.4}$$

The ghost number current $j = - : \beta\gamma :$ may be realized as $j(z) = \partial\phi(z)$ where $\phi(z)\phi(w) = -\ln(z-w)$. Notice that $\phi \neq \varphi$. We use the following mode expansions

$$\begin{aligned}
j(z) &= \sum_{n \in \mathbb{Z}} j_n z^{-n-1} \\
\varphi(z) &= q_\varphi + a_0 \ln z + \sum_{n \neq 0} \frac{a_n}{-n} z^{-n} \\
\phi(z) &= q_\phi + j_0 \ln z + \sum_{n \neq 0} \frac{j_n}{-n} z^{-n}
\end{aligned} \tag{4.5}$$

with hermiticity properties (see (4.10))

$$a_n^\dagger = a_{-n} - \sqrt{\frac{2}{t}} \delta_{n,0} \quad , \quad j_n^\dagger = -j_{-n} + \delta_{n,0} \tag{4.6}$$

The non-trivial commutation relations read

$$\begin{aligned}
[j_0, q_\phi] &= -1 \\
[j_n, j_m] &= -n\delta_{n+m,0} \\
[a_0, q_\varphi] &= 1 \\
[a_n, a_m] &= n\delta_{n+m,0}
\end{aligned} \tag{4.7}$$

The ket-vacuum, invariant under both conformal and loop projective $SL(2)$, is $|0\rangle$ satisfying

$$\begin{aligned}
\beta_n|0\rangle &= 0 = \gamma_n|0\rangle = a_n|0\rangle = j_n|0\rangle, \quad n > 0 \\
\beta_0|0\rangle &= 0 = a_0|0\rangle = j_0|0\rangle \\
\gamma_0|0\rangle &\neq 0 \\
L_n|0\rangle &= 0 \quad , \quad n \geq -1
\end{aligned} \tag{4.8}$$

with

$$L_n = \sum_{m \in \mathbb{Z}} (-m : \beta_{n-m} \gamma_m : + \frac{1}{2} : a_{n-m} a_m :) - (n+1) \sqrt{\frac{1}{2t}} a_n \tag{4.9}$$

The normal ordering of modes places annihilation operators to the right of creation operators. Correspondingly the $SL(2)$ invariant bra-vacuum $\langle sl_2|$ satisfies

$$\langle sl_2| L_n = 0 \quad , \quad n \leq 1$$

$$\begin{aligned}
\langle sl_2 | \beta_n &= 0 & , & & n \leq 0 \\
\langle sl_2 | \gamma_n = \langle sl_2 | j_n = \langle sl_2 | a_n &= 0 & , & & n \leq -1 \\
\langle sl_2 | j_0 &= \langle sl_2 | \\
\langle sl_2 | a_0 &= \sqrt{\frac{2}{t}} \langle sl_2 | \\
\langle sl_2 | 0 \rangle &= 0
\end{aligned} \tag{4.10}$$

The second to last equality follows from $\langle sl_2 | L_0 = 0$. The last equality is due to the fact that the bra-vacuum defined above carries different charges comparing to the ket-vacuum. In what follows we shall define another bra-vacuum with all the charges at infinity screened. This will be the dual vacuum we are mostly going to use in calculating the correlators.

We define the *dual vacuum state* $\langle 0 |$ in the WZNW free field realization as

$$\langle 0 | = \langle sl_2 | e^{-q_\phi} e^{\sqrt{2/t} q_\phi} \tag{4.11}$$

It satisfies

$$\begin{aligned}
\langle 0 | 0 \rangle &= 1 \\
\langle 0 | \gamma_0 &= 0 \\
\langle 0 | \beta_0 &\neq 0 \\
\langle 0 | a_0 &= 0 \\
\langle 0 | j_0 &= 0 \\
\langle 0 | \beta(z) \gamma(w) | 0 \rangle &= \frac{1}{z - w}
\end{aligned} \tag{4.12}$$

From the dual vacuum we construct dual bra-states of lowest $SL(2)$ weight

$$\langle j | = \langle 0 | e^{j \sqrt{2/t} q_\phi} \tag{4.13}$$

This state indeed satisfies the conditions for being a *lowest weight state* of the affine algebra

$$\begin{aligned}
\langle j | J_0^3 &= j \langle j | \\
\langle j | J_n^- &= 0 & , & & n \leq 0 \\
\langle j | J_n^3 &= 0 & , & & n < 0 \\
\langle j | J_n^+ &= 0 & , & & n < 0
\end{aligned} \tag{4.14}$$

For the corresponding ket-states

$$|j\rangle = e^{-j \sqrt{2/t} q_\phi} |0\rangle \tag{4.15}$$

we have

$$\langle j | j \rangle = 1 \tag{4.16}$$

and this ket-state is similarly a *highest weight state* of the affine algebra. We notice that

$$\langle 0 | J_0^+ = \langle 0 | \beta_0 \neq 0 \tag{4.17}$$

Thus we are performing all calculations with an $SL(2)$ non-invariant bra-vacuum. This gives rise to some complications when we wish to prove projective and global (loop projective) $SL(2)$ invariance of our correlators. In a subsequent section we shall explicitly demonstrate that the above state, $\langle 0|\beta_0$, while not being zero is in fact a BRST [19] exact state in the sense of Felder [56], and that therefore it must be expected to decouple from all correlators. This decoupling we then verify.

4.1.1 Admissible Representations

Here we introduce the notion of admissible representations [89]. Most of the constructions in this chapter and in Chapter 5 pertain to such representations.

Degenerate highest weight representations [87] may be parameterised as

$$\begin{aligned} 2j^+ + 1 &= r - st & , & & (r, s) \geq (1, 0) \\ 2j^- + 1 &= -r + st & , & & (r, s) \geq (1, 1) \end{aligned} \quad (4.18)$$

Among these the admissible ones [89] are characterised by having rational t (or rational level $k = t - 2$)

$$t = \pm \frac{p}{q} \quad , \quad (p, q) = 1 \quad (4.19)$$

where $(p, q) = 1$ means that p and q are co-prime. In the upper case ($t = +\frac{p}{q}$) it is possible to recover the minimal models by Hamiltonian reduction (see later this chapter) while the lower case corresponds to the Liouville series with $c > 25$ which is dual to the minimal models. Here we concentrate on the first case. Any $j_{r,s}^-$ may be written

$$j_{r,s}^- = -j_{r,s}^+ - 1 \quad (4.20)$$

and in the case of admissible representations we have the translation symmetries

$$\begin{aligned} j_{r,s}^- &= \frac{p - r - 1}{2} - \frac{q - s}{2}t = j_{p-r, q-s}^+ \\ j_{r,s}^\pm &= j_{r+np, s+nq}^\pm \end{aligned} \quad (4.21)$$

One notices the special case of $s = 0$ (or the series $s = nq$ for admissible representations) which is only present in the plus region. Because of the translation symmetry we may choose to work with j^+ parametrisations solely.

Among the admissible representations we find the better known integrable representations where $2j$ is integer, so they simply correspond to the case $s = 0$. In the following sections we shall discuss how one constructs correlators using screening currents in the case of admissible representations, generalising the Dotsenko-Fateev construction [42] of minimal models. In order to be able to screen both integer (the r part) and fractional (the s part) spins, we need both kinds of screening currents. For integrable representations the first kind suffices, and it is exactly the introduction of the second kind of screening current that courses many of the complications due to the fractional power of the β ghost field (4.3).

4.2 3-point Functions

Let us now consider the evaluation of the (chiral) 3-point function

$$\langle j_3 | \phi_{j_2}(z, x) | j_1 \rangle \quad (4.22)$$

where the dual bra $\langle j_3 |$ and the ket $| j_1 \rangle$ were defined in the preceding section. Using the free field realizations of $\phi_{j_2}(z, x)$, the 3-point function may be evaluated only provided the charges may be screened away in a way similar to the one employed in the case of minimal models [42], see above. Correspondingly $\phi_{j_2}(z, x)$ is replaced by the intertwining field $[\phi_{j_2}(z, x)]_{j_1}^{j_3}$ [56, 22, 26], which maps a j_1 highest weight module into a j_3 highest weight module. In this case [110] we are led to consider

$$\begin{aligned} [\phi_{j_2}(z, x)]_{j_1}^{j_3} &= \int \prod_{j=1}^s dv_j \prod_{i=1}^r du_i \phi_{j_2}(z, x) P(u_1, \dots, u_r; v_1, \dots, v_s) \\ P(u_1, \dots, u_r; v_1, \dots, v_s) &= \prod_{j=1}^s \tilde{s}(v_j) \prod_{i=1}^r s(u_i) \\ &= \prod_{j=1}^s \beta^{-t}(v_j) : e^{-\sqrt{2t}\varphi(v_j)} : \prod_{i=1}^r \beta(u_i) : e^{\sqrt{2/t}\varphi(u_i)} : \end{aligned} \quad (4.23)$$

The a_0 charge conservation reads

$$j_1 + j_2 - j_3 = r - st \quad (4.24)$$

with r and s non-negative integers. Let us notice that the definition (3.65) of the primary field ensures that it transforms covariantly under both conformal transformations and loop projective transformations, namely as an $\Delta(\phi_j)$ tensor field for the former, and a $-j$ tensor field for the latter

$$\begin{aligned} z &\rightarrow f(z) \\ x &\rightarrow \frac{a(z)x + b(z)}{c(z)x + d(z)} \quad , \quad a(z)d(z) - b(z)c(z) = 1 \end{aligned} \quad (4.25)$$

The last statement follows from the fact that the solution to

$$T_a J^a(x) \phi_j(z, x) = \delta_\epsilon \phi_j(z, x) = T_a \epsilon^a(x) \partial_x \phi_j(z, x) - j T_a \partial_x \epsilon^a(x) \phi_j(z, x) \quad (4.26)$$

is a second order polynomial in x which is indeed an infinitesimal (loop) projective transformation. Invariance under (loop) projective transformations, notably the conservation of the (J_0^3) L_0 charge

$$\langle j_3 | L_0 [\phi_{j_2}(z, x)]_{j_1}^{j_3} | j_1 \rangle = \langle j_3 | [L_0, [\phi_{j_2}(z, x)]_{j_1}^{j_3}] | j_1 \rangle + \langle j_3 | [\phi_{j_2}(z, x)]_{j_1}^{j_3} L_0 | j_1 \rangle \quad (4.27)$$

determines the singular z -(x -)behaviour

$$\langle j_3 | [\phi_{j_2}(z, x)]_{j_1}^{j_3} | j_1 \rangle = C_{j_1 j_2 j_3} z^{-\Delta_1 - \Delta_2 + \Delta_3} x^{j_1 + j_2 - j_3} (1 + \mathcal{O}(z, x)) \quad (4.28)$$

It is obvious that this fact simply follows from putting $z_3, x_3 = \infty$, $z_2, x_2 = z, x$ and $z_1, x_1 = 0$ (taking into account powers of z, x as in (2.50)) in the general expression for the 3-point function

$$\langle \phi_{j_3}(z_3, x_3) \phi_{j_2}(z_2, x_2) \phi_{j_1}(z_1, x_1) \rangle \propto \frac{(x_2 - x_1)^{j_2+j_1-j_3} (x_3 - x_2)^{j_3+j_2-j_1} (x_3 - x_1)^{j_3+j_1-j_2}}{(z_2 - z_1)^{\Delta_2+\Delta_1-\Delta_3} (z_3 - z_2)^{\Delta_3+\Delta_2-\Delta_1} (z_3 - z_1)^{\Delta_3+\Delta_1-\Delta_2}} \quad (4.29)$$

which is indeed found using the projective invariances (see also (2.56)). In Chapter 5 there will be a discussion on the structure constants $C_{j_1 j_2 j_3}$ appearing as the proportionality constants when including the anti-holomorphic sector..

The techniques to perform the φ -part of the Wick contractions were described in the section on minimal models and we find

$$\begin{aligned} W_3^\varphi &= \int \prod_{i=1}^r du_i \prod_{j=1}^s dv_j \prod_{i=1}^r u_i^{-2j_1/t} (1 - u_i)^{-2j_2/t} \prod_{i_1 < i_2} (u_{i_1} - u_{i_2})^{2/t} \\ &\cdot \prod_{j=1}^s v_j^{2j_1} (1 - v_j)^{2j_2} \prod_{j_1 < j_2} (v_{j_1} - v_{j_2})^{2t} \prod_{i,j} (u_i - v_j)^{-2} \end{aligned} \quad (4.30)$$

Here we concentrate on explaining how to perform the β, γ -part. First we have to determine the asymptotic expansion in γ within $\phi_{j_2}(z, x)$. By projective invariance (4.28) x could be fractionally powered when s (4.24) is non-zero, hence we should expand asymptotically

$$\begin{aligned} \phi_{j_2}(z, x) &= (1 + \gamma(z)x)_{(-st)}^{2j_2} : e^{-j_2 \sqrt{2/t} \varphi(z)} : \\ &= \sum_n \binom{2j_2}{n - st} (\gamma(z)x)^{n-st} : e^{-j_2 \sqrt{2/t} \varphi(z)} : \end{aligned} \quad (4.31)$$

and as in (2.97)

$$\begin{aligned} &\beta(w)^{-t} (1 + \gamma(z)x)_{(-st)}^{2j_2} \\ &= : (\beta(w) + \frac{1}{w-z} \partial_{\gamma(z)})^{-t} \sum_m \binom{2j_2}{m - st} (\gamma(z)x)^{m-st} : \\ &= \sum_n \binom{-t}{n} : \beta^n(w) (w-z)^{t+n} \partial_{\gamma(z)}^{-t-n} \sum_m \binom{2j_2}{m - st} (\gamma(z)x)^{m-st} : \\ &= \sum_n \binom{-t}{n} : \beta^n(w) (w-z)^{t+n} \frac{\Gamma(2j_2+1)}{\Gamma(2j_2+t+n+1)} (1 + \gamma(z)x)_{((1-s)t)}^{2j_2+t+n} : x^{-t-n} \end{aligned} \quad (4.32)$$

Similarly we deduce

$$\begin{aligned} &\prod_{i=1}^s \beta(w_i)^{-t} (1 + \gamma(z)x)_{(-st)}^{2j_2} \\ &= : \prod_{i=1}^s (\beta(w_i) + \frac{1}{w_i-z} \partial_{\gamma(z)})^{-t} \sum_m \binom{2j_2}{m - st} (\gamma(z)x)^{m-st} : \\ &= : \prod_{i=1}^s \sum_{n_i} \binom{-t}{n_i} \beta^{n_i}(w_i) (w_i-z)^{t+n_i} \partial_{\gamma(z)}^{-t-n_i} \sum_m \binom{2j_2}{m - st} (\gamma(z)x)^{m-st} : \\ &= : \prod_{i=1}^s \sum_{n_i} \binom{-t}{n_i} \beta^{n_i}(w_i) (w_i-z)^{t+n_i} \frac{\Gamma(2j_2+1)}{\Gamma(2j_2+st+\sum_i n_i+1)} \\ &\cdot (1 + \gamma(z)x)^{2j_2+st+\sum_i n_i} : x^{-st-\sum_i n_i} \end{aligned} \quad (4.33)$$

Notice that in these equations β and γ appear within normal ordering signs with integral powers. (4.32) and (4.33) suggest that we consider some kind of generating function, which looks like the exponential function

$$F(u) = \sum_n \frac{1}{\Gamma(n-a+1)} (1 + \gamma(z)x)_{(\alpha)}^{n-a} u^{-n+a} \quad (4.34)$$

In the simple case of 3-point functions we do not gain much by this introduction but for more general N -point function it turns out to be of great value. We shall find it useful to use the following rather trivial identity

$$(1 + \gamma(z)x)^{2j} = \Gamma(2j+1) \oint_0 \frac{du}{2\pi i} \frac{1}{u} (u^{-1}D)^{-2j} \exp[(1 + \gamma(z)x)/u] \quad (4.35)$$

where D converts the exponential function into the derivative of that function, in particular it acts on and only on the entire argument of that function. We now prove the following

Lemma

$$\beta^a(w) \exp[(1 + \gamma(z)x)/u] =: (\beta(w) + \frac{x/u}{w-z})^a D^a \exp[(1 + \gamma(z)x)/u] : \quad (4.36)$$

Proof

$$\begin{aligned} & \beta^a(w) \exp[(1 + \gamma(z)x)/u] \\ &= \sum_{k,m,n} \binom{a}{m} : \beta^m(w) (w-z)^{m-a} \partial_{\gamma(z)}^{a-m} \frac{1}{\Gamma(n+1)} \binom{n}{k} (\gamma(z)x)^{n-k} : u^{-n} \\ &= \sum_{k,m,n} \binom{a}{m} : \beta^m(w) (w-z)^{m-a} \gamma^{n-k-a+m}(z) : \\ & \quad \cdot \frac{1}{\Gamma(k+1)} \frac{1}{\Gamma(n-k-a+m+1)} x^{n-k-a+m} x^{a-m} u^{-n} \\ &= \sum_{k,m,N} \binom{a}{m} : \beta^m(w) (w-z)^{m-a} x^{a-m} u^{m-a} \gamma^{-a+N-k}(z) : \\ & \quad \cdot \binom{-a+N}{k} \frac{1}{\Gamma(-a+N+1)} x^{-a+N-k} u^{a-N} \\ &= : (\beta(w) + \frac{x/u}{w-z})^a D^a \exp[(1 + \gamma(z)x)/u] : \end{aligned} \quad (4.37)$$

□

We may now calculate the $\beta\gamma$ contractions in the 3-point function

$$\begin{aligned} & \prod_{i=1}^r \beta(u_i) \prod_{j=1}^s \beta^{-t}(v_j) (1 + \gamma(z)x)^{2j} \\ &= \oint_0 \frac{du}{2\pi i} \frac{u^{2j}}{u} : \prod_{i=1}^r [\beta(u_i) + \frac{x/u}{u_i-z}] \prod_{j=1}^s [\beta(v_j) + \frac{x/u}{v_j-z}]^{-t} \\ & \quad \cdot D^{-2j+r-st} \exp\{\frac{1 + \gamma(z)x}{u}\} : \Gamma(2j+1) \end{aligned} \quad (4.38)$$

When inserted between $\langle j_3|$ and $|j_1\rangle$ to produce $W_3^{\beta\gamma}$ (the β, γ part of the 3-point function) we effectively put $\beta = 0 = \gamma$ (according to j_0 charge conservation) whereupon the u integration becomes trivial, and we find the result

$$W_3^{\beta\gamma} = \frac{\Gamma(2j_2 + 1)}{\Gamma(2j_2 - r + st + 1)} x^{r-st} \int \prod_{i=1}^r du_i (u_i - z)^{-1} \prod_{j=1}^s dv_j (v_j - z)^t \quad (4.39)$$

Combining this with the φ -part (4.30) and using projective invariances to put $z = 1 = x$ we obtain

$$\begin{aligned} W_3 = & \frac{\Gamma(2j_2 + 1)}{\Gamma(2j_2 - r + st + 1)} \int \prod_{i=1}^r du_i \prod_{j=1}^s dv_j \prod_{i=1}^r u_i^{-2j_1/t} (1 - u_i)^{-2j_2/t-1} \prod_{i_1 < i_2} (u_{i_1} - u_{i_2})^{2/t} \\ & \cdot \prod_{j=1}^s v_j^{2j_1} (1 - v_j)^{2j_2+t} \prod_{j_1 < j_2} (v_{j_1} - v_{j_2})^{2t} \prod_{i,j} (u_i - v_j)^{-2} \end{aligned} \quad (4.40)$$

In the derivation we did not pay any attention to the phases. Nevertheless, in the sequel we will consider (4.40) as the defining expression for the 3-point functions.

The final expression for the 3-point function is obviously depending on the choice of contours. However, let us postpone a discussion of these to a subsequent section.

Due to projective invariance

$$W(j_1, j_2, j_3) \propto W(P(j_1, j_2, j_3)) \quad (4.41)$$

for any permutation P where the proportionality constant is non-vanishing and finite. There exists an involution

$$\begin{aligned} J_n^+ & \rightarrow J_n'^+ = J_{1+n}^- \\ J_n^3 & \rightarrow J_n'^3 = -J_n^3 + \frac{t-2}{2} \delta_{n,0} \\ J_n^- & \rightarrow J_n'^- = J_{-1+n}^+ \end{aligned} \quad (4.42)$$

Using this one must expect that

$$\langle j_3 | \phi_{j_2}(z, x) | j_1 \rangle \propto \langle -j_3 - 1 + t/2 | \phi'_{j_2}(z', x') | -j_1 - 1 + t/2 \rangle \quad (4.43)$$

again with a non-vanishing and finite proportionality constant and where

$$\phi'_j(z', x') = \phi_j(z, x) z^j x^{-2j} \quad z' = z, x' = z/x \quad (4.44)$$

is seen to be a primary field in the primed notation in the same sense that $\phi_j(z, x)$ is primary in the unprimed system

There is a considerable freedom in choosing the numbers of screenings subject to the charge conservation (4.24) since $p - qt = 0$. However, it turns out [112] that we only need two different sets of screenings

Standard Screening

$$\begin{aligned} 2r &= r_1 + r_2 - r_3 - 1 \\ 2s &= s_1 + s_2 - s_3 \end{aligned} \quad (4.45)$$

Over-screening

$$\begin{aligned} 2r &= r_1 + r_2 - r_3 - 1 + p \\ 2s &= s_1 + s_2 - s_3 + q \end{aligned} \quad (4.46)$$

This is denoted over-screening due to the addition of p, q . We will make use of both of these when determining the fusion rules below.

We conclude this section by making a comment on the possible $SL(2)$ representations carried by the intertwining field $[\phi_{j_2}(z, x)]_{j_1}^{j_3}$. On the ket-vacuum (see (4.143)),

$$\begin{aligned} [\phi_{j_2}(z, x)]_{j_1}^{j_3} |0\rangle &= [\phi_{j_2}(z, x)]_0^{j_2} |0\rangle \delta_{j_1, 0} \delta_{j_3, j_2} \\ &= e^{xJ_0^-} e^{zL-1} |j_2\rangle \delta_{j_1, 0} \delta_{j_3, j_2} \end{aligned} \quad (4.47)$$

is in a highest weight representation of the $SL(2)$ current algebra with the highest weight state $|j_2\rangle$. Similarly on the dual vacuum state, $\langle 0|[\phi_{j_2}(z, x)]_{j_1}^{j_3}$ is in a lowest weight representation of the $SL(2)$ current algebra with the lowest weight state $\langle j_2|$. However, when sandwiched in the middle of the correlator, the intertwining field $[\phi_{j_2}(z, x)]_{j_1}^{j_3}$ might carry representations belonging to the continuous series of the $SL(2)$ algebra. A representation belong to this series when neither a highest nor a lowest weight state exists. In our case the intertwining field belong to such a representation when both $j_1 + j_2 - j_3$ and $j_1 - j_2 - j_3$ are non-integers. To understand this let us notice that the monodromy of the field forces an expansion of it to be offset the integers by $\alpha = r - st = j_1 + j_2 - j_3$

$$[\phi_{j_2}(z, x)]_{j_1}^{j_3} = \sum_n \phi_{j_2, j_2 - \alpha - n}(z, x) \quad , \quad \phi_{j_2, \lambda}(z, x) = x^{j_2 - \lambda} \hat{\phi}_{j_2, \lambda}(z) \quad (4.48)$$

This is a J_0^3 eigenstate decomposition

$$\begin{aligned} [J_0^3, \phi_{j_2, \lambda}(z, x)] &= \lambda \phi_{j_2, \lambda}(z, x) \\ [J_0^+, \phi_{j_2, \lambda}(z, x)] &= (j_2 + \lambda) x \phi_{j_2, \lambda}(z, x) \\ [J_0^-, \phi_{j_2, \lambda}(z, x)] &= (j_2 - \lambda) x^{-1} \phi_{j_2, \lambda}(z, x) \end{aligned} \quad (4.49)$$

We see that the condition for $[\phi_{j_2}(z, x)]_{j_1}^{j_3}$ to belong to the continuous series may equivalently be characterised by $\lambda \notin \mathbb{Z} \pm j_2$. Although in that case $[\phi_{j_2}(z, x)]_{j_1}^{j_3}$ does not correspond to a highest weight representation, it maps a j_1 highest weight representation to a j_3 highest weight representation.

4.3 Fusion Rules

4.3.1 Decoupling of Singular States

In [12] the fusion rules are determined from considering decoupling of singular vectors as they are described in [101]. Here we review this derivation of the fusion rules.

Let M_j denote the Verma module generated by the highest weight vector $|j\rangle$ (2.194), notably $J_0^3 |j\rangle = j |j\rangle$. An affine singular vector $|\chi\rangle \in M_j$ of grade (or at level) N and charge Q is defined

by being a highest weight state itself, of weight $j + Q$ and by satisfying $L_0|\chi\rangle = (\Delta_j + N)|\chi\rangle$, where $\Delta_j = j(j+1)/t$. According to [87, 101] there exists a unique singular vector $|\chi^\pm\rangle$ of grade $N = rs$ and charge $Q^\pm = \mp r$ when one uses the parametrisation of (4.18), and it is given by [101]

$$\begin{aligned} |\chi_{r,s}^+\rangle &= (F_0)^{r+st}(E_{-1})^{r+(s-1)t} \dots (E_{-1})^{r-(s-1)t} (F_0)^{r-st} |j_{r,s}^+\rangle \\ |\chi_{r,s}^-\rangle &= (E_{-1})^{r+(s-1)t} (F_0)^{r+(s-2)t} \dots (F_0)^{r-(s-2)t} (E_{-1})^{r-(s-1)t} |j_{r,s}^-\rangle \end{aligned} \quad (4.50)$$

Note the consistency $N = (j^\pm \mp r)(j^\pm \mp 1)/t - j^\pm(j^\pm + 1)/t = rs$. For general $t \in \mathbb{C} \setminus \{0\}$ the exponents in (4.50) are complex numbers, nevertheless it is shown in [101] that they make sense by analytic continuation. This may be done by rearranging the terms and then show that the product of generators lie in the enveloping algebra, thereby rendering the above standard singular states. Further discussions and relations to the Virasoro singular vectors can be found in [17, 94].

The analogue of (4.28) but involving a singular state is

$$\langle j_3 | [\phi_{j_2}(z, x)]_{j_1^\pm}^{j_3} |\chi_{r,s}^\pm\rangle = C_{j_1 j_2 j_3} z^{-\Delta_1 - \Delta_2 + \Delta_3 - rs} x^{j_1^\pm + j_2 - j_3 \mp r} f_{r,s}^\pm(j_1^\pm, j_2, j_3) (1 + \mathcal{O}(z, x)) \quad (4.51)$$

Using (4.3) ($F_0 = \partial_x$, $E_{-1} = z^{-1}(-x^2\partial_x + 2xj)$) and the rule (2.109) for fractional derivation one finds

$$\begin{aligned} f_{r,s}^+ &= \prod_{n=0}^{r-1} \prod_{m=0}^s (j_1^+ + j_2 - j_3 - n + mt) \prod_{n=1}^r \prod_{m=1}^s (-j_1^+ + j_2 + j_3 + n - mt) \\ f_{r,s}^- &= \prod_{n=0}^{r-1} \prod_{m=0}^{s-1} (-j_1^- + j_2 + j_3 - n + mt) \prod_{n=1}^r \prod_{m=1}^{s-1} (j_1^- + j_2 - j_3 + n - mt) \end{aligned} \quad (4.52)$$

For a non-vanishing structure constant $C_{j_1 j_2 j_3}$, the decoupling of the singular vector $|\chi_{r,s}^\pm\rangle$ is equivalent to $f_{r,s}^\pm = 0$. For admissible representations where $j_{r,s}^- = j_{p-r,q-s}^+$ (4.21), we have the two independent singular vectors $|\chi_{r,s}^+\rangle$ and $|\chi_{p-r,q-s}^+\rangle$. Their decouplings result in (now $j_1 = j_1^+$)

$$\begin{aligned} 0 &= \prod_{n=0}^{r-1} \prod_{m=0}^s (j_1 + j_2 - j_3 - n + mt) \prod_{n=1}^r \prod_{m=1}^s (-j_1 + j_2 + j_3 + n - mt) \\ &= \prod_{n=0}^{p-r-1} \prod_{m=0}^{q-s-1} (-j_1 + j_2 + j_3 - n + mt) \prod_{n=1}^{p-r} \prod_{m=1}^{q-s-1} (j_1 + j_2 - j_3 + n - mt) \end{aligned} \quad (4.53)$$

Now the fusion rules are readily obtained using that if

$$\begin{aligned} x &= n + m + c_x & , & & a_n \leq n \leq b_n \\ y &= n - m + c_y & , & & a_m \leq m \leq b_m \end{aligned} \quad (4.54)$$

then

$$\begin{aligned} -x + c_x + c_y + 2a_n &\leq y \leq -x + c_x + c_y + 2b_n \\ x - c_x + c_y - 2b_m &\leq y \leq x - c_x + c_y - 2a_m \end{aligned} \quad (4.55)$$

In conclusion, the fusion rules are

Fusion Rule I

$$\begin{aligned} 1 + |r_1 - r_2| &\leq r_3 \leq p - 1 - |r_1 + r_2 - p| \\ |s_1 - s_2| &\leq s_3 \leq q - 1 - |s_1 + s_2 - q + 1| \end{aligned} \quad (4.56)$$

Fusion Rule II

$$\begin{aligned} 1 + |p - r_1 - r_2| &\leq r_3 \leq p - 1 - |r_1 - r_2| \\ 1 + |q - s_1 - s_2 - 1| &\leq s_3 \leq q - 2 - |s_1 - s_2| \end{aligned} \quad (4.57)$$

In both cases, r_3 and s_3 jump in steps of 2. It is easily checked that both sets of fusion rules cannot be satisfied simultaneously.

In [55] the same set of fusion rules are obtained by cohomological methods.

4.3.2 Standard and Over-screening of 3-point Functions

In this section we will discuss how the fusion rules arise from the 3-point function (4.40) by appropriate choices of contours and numbers of screening currents [110, 112].

Our considerations will rely on the famous Dotsenko-Fateev (DF) integral (last paper in Ref. [42], appendix A, here with minor misprints corrected)

$$\begin{aligned} \mathcal{J}_{nm}(a, b; \rho) &= T \int_0^1 \prod_{i=1}^n du_i \prod_{j=1}^m dv_j u_i^{a'} (1 - u_i)^{b'} \prod_{i_1 < i_2} (u_{i_1} - u_{i_2})^{2\rho'} \\ &\cdot v_j^a (1 - v_j)^b \prod_{j_1 < j_2} (v_{j_1} - v_{j_2})^{2\rho} \prod_{i,j}^{n,m} (u_i - v_j)^{-2} \\ &= \rho^{2nm} \prod_{i=1}^n \frac{\Gamma(i\rho')}{\Gamma(\rho')} \prod_{i=1}^m \frac{\Gamma(i\rho - n)}{\Gamma(\rho)} \\ &\cdot \prod_{i=0}^{n-1} \frac{\Gamma(1 + a' + i\rho') \Gamma(1 + b' + i\rho')}{\Gamma(2 - 2m + a' + b' + (n - 1 + i)\rho')} \\ &\cdot \prod_{i=0}^{m-1} \frac{\Gamma(1 - n + a + i\rho) \Gamma(1 - n + b + i\rho)}{\Gamma(2 - n + a + b + (m - 1 + i)\rho)} \end{aligned} \quad (4.58)$$

where the parameters are subject to the relations

$$a' = -\rho' a, \quad b' = -\rho' b, \quad \rho' = \rho^{-1} \quad (4.59)$$

The integral is real and time ordered (2.166). In Chapter 5 we shall also need to consider another version of this integral, namely

$$\begin{aligned} \tilde{\mathcal{J}}_{nm}(a, b; \rho) &= T \int_1^\infty \prod_{i=1}^n du_i \prod_{j=1}^m dv_j u_i^{a'} (u_i - 1)^{b'} \prod_{i_1 > i_2} (u_{i_1} - u_{i_2})^{2\rho'} \\ &\cdot v_j^a (v_j - 1)^b \prod_{j_1 > j_2} (v_{j_1} - v_{j_2})^{2\rho} \prod_{i,j}^{n,m} (u_i - v_j)^{-2} \\ &= \mathcal{J}_{nm}(-a - b + 2(n - 1) - 2(m - 1)\rho, b; \rho) \end{aligned} \quad (4.60)$$

The last equality follows from changing variables $u_i \rightarrow 1/u_i$, $v_j \rightarrow 1/v_j$ and using (4.59). It follows that the 3-point function (4.40) may be evaluated using the DF integration. However, we need complex contours, both of the Felder [56] type (a set of closed circle-like contours passing through a common point which is the only intersection point) and of the Dotsenko-Fateev [42] type (a set of open contours with common end points which are the only intersection points). The choice of contours give rise to certain pre-factors multiplying the DF integral. Due to the relations (4.59) one may interchange or pull through each other a DF contour of a u variable and a DF contour of a v variable [42].

Let us be more specific and consider (4.40) in the case of admissible representations. In the case that all contours are Felder contours (labelling by $1, \dots, r$ and $1, \dots, s$ the 'circles' from in-most to out-most) one finds [112]

$$W_{FF}^{r,s} = \chi_r^{(1)}(r_1; 1/t) \chi_s^{(2)}(s_1; t) \frac{\Gamma(2j_2 + 1)}{\Gamma(2j_2 - r + st + 1)} \mathcal{J}_{rs}(2j_1, 2j_2 + t; t) \quad (4.61)$$

where the χ -functions [112]

$$\begin{aligned} \chi_r^{(1)}(r_1; 1/t) &= e^{i\pi r(r+1-2r_1)/t} \prod_{j=1}^r \frac{(1 - e^{2\pi i(r_1-j)/t})(1 - e^{2\pi i j/t})}{1 - e^{2\pi i/t}} \\ &= (2i)^r e^{i\pi r(r-r_1)/t} \prod_{j=1}^r \frac{s((j-r_1)/t)s(j/t)}{s(1/t)} \end{aligned} \quad (4.62)$$

$$\begin{aligned} \chi_s^{(2)}(s_1; t) &= e^{i\pi t s(s-1-2s_1)} \prod_{j=1}^s \frac{(1 - e^{2\pi i t(s_1+1-j)})(1 - e^{2\pi i t j})}{1 - e^{2\pi i t}} \\ &= (2i)^s e^{i\pi t s(s-1-s_1)} \prod_{j=1}^s \frac{s((j-s_1-1)t)s(jt)}{s(t)} \end{aligned} \quad (4.63)$$

(with $s(x) = \sin(\pi x)$) arise when performing contour deformations². It turns out that the Felder contours alone cannot produce a well-defined and non-vanishing 3-point function corresponding to fusion rule II (see below). We need the combination that the r screening variables of the first kind are integrated along DF contours, while the s screenings of the second kind are taken along Felder contours (or vice versa). Let the convention for DF contours be that the labelling puts contours with lower indices below contours with higher indices. Now contour manipulations give

$$W_{DF}^{r,s} = \lambda_r(1/t) \chi_s^{(2)}(s_1; t) \frac{\Gamma(2j_2 + 1)}{\Gamma(2j_2 - r + st + 1)} \mathcal{J}_{rs}(2j_1, 2j_2 + t; t) \quad (4.64)$$

where we have introduced the functions [42, 112]

$$\lambda_r(1/t) = \prod_{j=1}^r e^{-i\pi(j-1)/t} \frac{s(j/t)}{s(1/t)} \quad (4.65)$$

²It is not necessary to introduce two χ -functions since $\chi_n^{(1)}(a; t) = \chi_n^{(2)}(a-1; t)$. However, it helps clarifying the origins of the expressions.

Using the alternative combination where the r screening variables of the first kind are integrated along Felder contours while the s screening variables are taken along DF contours, results in the pre-factor $\chi_r^{(1)}(r_1; 1/t)\lambda_s(t)$. Similarly one may choose all contours of the DF type and then obtain the pre-factor $\lambda_r(1/t)\lambda_s(t)$.

The fusion rules determine when a 3-point function is well-defined and non-vanishing. According to (4.41) and (4.43) the fusion rules should be invariant under certain transformations. Because of the translation symmetry (4.21) we may choose to work with j^+ parameterisations whereby we choose the fusion rules to be expressed in terms of such parameterisations. Hence the transformations are

$$(r_1, r_2, r_3; s_1, s_2, s_3) \rightarrow (P(r_1, r_2, r_3); P(s_1, s_2, s_3)) \quad (4.66)$$

and

$$\begin{aligned} r_1 &\rightarrow p - r_1 \\ r_2 &\rightarrow r_2 \\ r_3 &\rightarrow p - r_3 \\ s_1 &\rightarrow q - s_1 - 1 \\ s_2 &\rightarrow s_2 \\ s_3 &\rightarrow q - s_3 - 1 \end{aligned} \quad (4.67)$$

Of course these may be invoked in any order. The exact form of (4.67) follows from (4.43) since $j_i \rightarrow -j_i - 1 + t/2$ corresponds to $(r_i, s_i) \rightarrow (p - r_i, q - s_i - 1)$.

Even before going into a detailed evaluation of the gamma- and sine-functions in the 3-point function we can derive the fusion rules as *necessary* conditions for a 3-point function to be well-defined. Let us consider the case of over-screening (4.46). Assume that $r \geq p$ which means that $r_1 + r_2 - r_3 - 1 + p \geq 2p$ and therefore $r_3 \leq -p - 1 + r_1 + r_2$. If this inequality is allowed by a set of fusion rules then we may impose permutation invariance (4.66) leading to $p + 1 + |r_1 - r_2| \leq r_3 \leq -p - 1 + r_1 + r_2$. Without loss of generality we let $r_1 \geq r_2$ and find $p + 1 + r_1 - r_2 \leq -p - 1 + r_1 + r_2$ or equivalently $p + 1 \leq r_2$ which takes us out of the standard region, so we conclude that $r < p$. This leads to $2r \leq 2p - 2$ and $-p + 1 + r_1 + r_2 \leq r_3$. Imposing permutation invariance (4.66) yields $-p + 1 + r_1 + r_2 \leq r_3 \leq p - 1 - |r_1 - r_2|$. Finally we impose the involutory invariance (4.67) and obtain

$$1 + |p - r_1 - r_2| \leq r_3 \leq p - 1 - |r_1 - r_2| \quad (4.68)$$

Similarly the assumption $s \geq q$ will take us out of the standard region by $q \leq s_2 \leq s_1$. On the other hand $s < q$ leads to $-q + 2 + s_1 + s_2 \leq s_3$ and as before we impose permutation invariance (4.66) yielding $-q + 2 + s_1 + s_2 \leq s_3 \leq q - 2 - |s_1 - s_2|$ and finally the involution (4.67) gives

$$1 + |q - 1 - s_1 - s_2| \leq s_3 \leq q - 2 - |s_1 - s_2| \quad (4.69)$$

and we have obtained fusion rule II. Fusion rule I follows as necessary conditions simply from imposing the invariances on the conditions $r, s \geq 0$.

Now we want to establish that the 3-point function (4.40) with Felder contours (4.61) is well-defined and non-vanishing exactly when fusion rule I is imposed. Furthermore, we want to

establish that (4.40), with the combination of the r screening variables of the first kind being integrated along DF contours and the s screening variables taken along Felder contours (4.64), is well-defined and non-vanishing exactly when fusion rule II is imposed. In the verification we allow cancellations of the form $\Gamma(0)/\Gamma(0) = 1$. Since we have already found that the fusion rules are necessary conditions we only need to show that indeed they are sufficient conditions. Thus, we simply count the numbers of $\Gamma(0)$ and $1/\Gamma(0)$ in $\mathcal{J}_{rs}(2j_1, 2j_2 + t; t)$ and in the pre-factors when imposing the fusion rules. The sufficient conditions are then equivalent to equality of these numbers. Notice that $s(0) \sim 1/\Gamma(0)$. We find that for fusion rule I all terms are regular

$$\lambda_r(1/t) \sim \lambda_s(t) \sim \chi_r^{(1)}(r_1; 1/t) \sim \chi_s^{(2)}(s_1; t) \sim \frac{\Gamma(2j_2 + 1)}{\Gamma(2j_2 - r + st + 1)} \sim \mathcal{J}_{rs}(2j_1, 2j_2 + t; t) \sim 1 \quad (4.70)$$

while for fusion rule II we find

$$\begin{aligned} \lambda_r(1/t) \sim \lambda_s(t) &\sim 1 \\ \chi_r^{(1)}(r_1; 1/t) \sim \chi_s^{(2)}(s_1; t) &\sim 1/\Gamma(0) \\ \frac{\Gamma(2j_2 + 1)}{\Gamma(2j_2 - r + st + 1)} &\sim 1 \\ \mathcal{J}_{rs}(2j_1, 2j_2 + t; t) &\sim (\Gamma(0))^2 / \Gamma(0) \end{aligned} \quad (4.71)$$

This concludes the derivation of the fusion rules from the free field realization of the 3-point function.

The above analysis shows that for fusion rule I, (4.40) is well-defined and non-vanishing for all 4 combinations of contours ($FF, FDF, DFF, DFDF$), while for fusion rule II we need a combination of Felder and DF contours (DF, FDF). Hence, for a given set of fusion rules a choice of contours among these allowed possibilities only affects the normalisation of the 3-point function.

4.3.3 Over-screening in Minimal Models

The fusion rules for minimal models have been known in many years [21, 56]. However, the technique of over-screening might reveal the existence of an unknown second set of fusion rules. This turns out not to be the case as one should expect, since in minimal models the primary fields are solely specified by their conformal weights, contrary to the primary fields in $SL(2)$ current which are characterised by spins as well. Nevertheless, we will present an explicit analysis of this lack of a second set of fusion rules.

Consider the minimal model 3-point function (in the limits $z_3, z_2, z_1 \rightarrow \infty, 1, 0$)

$$\begin{aligned} W_3 \sim & \int \prod_{k=1}^N dv_k v_k^{\alpha_{n_1, m_1} \alpha_+} (1 - v_k)^{\alpha_{n_2, m_2} \alpha_+} \prod_{k < k'} (v_k - v_{k'})^{\alpha_+^2} \\ & \cdot \prod_{i=1}^M dw_i w_i^{\alpha_{n_1, m_1} \alpha_-} (1 - w_i)^{\alpha_{n_2, m_2} \alpha_-} \prod_{i < i'} (w_i - w_{i'})^{\alpha_-^2} \prod_{k, i}^{N, M} (v_k - w_i)^{-2} \end{aligned} \quad (4.72)$$

where the charge conservation reads

$$\alpha_{n_1, m_1} + \alpha_{n_2, m_2} - \alpha_{n_3, m_3} + N\alpha_+ + M\alpha_- = 0 \quad (4.73)$$

Using the rationality condition $\alpha_-/\alpha_+ = -p/q = -t$ (where $\alpha_-^2/2 = t$, $\alpha_+^2/2 = 1/t$) we consider the following screenings

Standard Screening

$$N = \frac{1}{2}(n_1 + n_2 - n_3 - 1) \quad , \quad M = \frac{1}{2}(m_1 + m_2 - m_3 - 1) \quad (4.74)$$

Over-screening

$$N = \frac{1}{2}(n_1 + n_2 - n_3 - 1 + p) \quad , \quad M = \frac{1}{2}(m_1 + m_2 - m_3 - 1 + q) \quad (4.75)$$

Choosing all contours to be Felder contours and considering standard screening, yields

$$W_{FF} = \chi_N^{(1)}(n_1; 1/t) \chi_M^{(2)}(m_1 - 1; t) \mathcal{J}_{NM}(n_1 - 1 - (m_1 - 1)t, n_2 - 1 - (m_2 - 1)t; t) \quad (4.76)$$

This is seen to be non-vanishing and well-defined exactly when one employs the fusion rules

$$\begin{aligned} 1 + |n_1 - n_2| &\leq n_3 \leq p - 1 - |p - n_1 - n_2| \\ 1 + |m_1 - m_2| &\leq m_3 \leq q - 1 - |q - m_1 - m_2| \end{aligned} \quad (4.77)$$

where n_3 and m_3 jump in steps of 2. If we consider over-screening and choose the N contours as DF contours and the M contours as Felder contours, we find

$$W_{DF} = \lambda_N(1/t) \chi_M^{(2)}(m_1 - 1; t) \mathcal{J}_{NM}(n_1 - 1 - (m_1 - 1)t, n_2 - 1 - (m_2 - 1)t; t) \quad (4.78)$$

An analysis of this along the lines indicated above, one finds the cancellation $(\Gamma(0)/\Gamma(0))^2$ exactly when

$$\begin{aligned} 1 + |p - n_1 - n_2| &\leq n_3 \leq p - 1 - |n_1 - n_2| \\ 1 + |q - m_1 - m_2| &\leq m_3 \leq q - 1 - |m_1 - m_2| \end{aligned} \quad (4.79)$$

However, the point is that this set of fusion rules is obtained from (4.77) simply by transforming $n_3, m_3 \rightarrow p - n_3, q - m_3$, while the conformal weights enjoy the symmetry

$$\Delta_{n,m} = \Delta_{p-n,q-m} \quad (4.80)$$

In conclusion, (4.79) is equivalent to (4.77), and the act of over-screening in minimal models merely reflects the freedom (4.80) in parametrising the primary fields.

4.4 N -point Functions

We wish to evaluate the chiral block

$$W_N = \langle j_N | [\phi_{j_{N-1}}(z_{N-1}, x_{N-1})]_{\iota_{N-2}}^{j_N} \cdots [\phi_{j_n}(z_n, x_n)]_{\iota_{n-1}}^{\iota_n} \cdots [\phi_{j_2}(z_2, x_2)]_{j_1}^{\iota_2} | j_1 \rangle \quad (4.81)$$

Thus we have primary fields at points

$$z_1 = 0, z_2, \dots, z_{N-1}, z_N = \infty \quad (4.82)$$

having x values

$$x_1 = 0, x_2, \dots, x_{N-1}, x_N = \infty \quad (4.83)$$

From the pictorial version

one reads off the following screening conditions

$$\begin{aligned} j_1 + j_2 - l_2 &= \rho_2 - \sigma_2 t \\ l_2 + j_3 - l_3 &= \rho_3 - \sigma_3 t \\ &\vdots \\ l_{n-1} + j_n - l_n &= \rho_n - \sigma_n t \\ &\vdots \\ l_{N-2} + j_{N-1} - j_N &= \rho_{N-1} - \sigma_{N-1} t \\ 2j_i + 1 &= r_i - s_i t \end{aligned} \quad (4.85)$$

with σ_n, ρ_n non-negative integers, while the last line is the usual parametrisation of the weights. We then get for the β, γ part of the correlator, denoting by $w(n, i)$ and $v(n, k)$ the positions of the i 'th and the k 'th screening currents of the first and second kinds respectively around the n 'th primary field

$$\begin{aligned} W_N^{\beta\gamma} &= \langle j_N | \prod_{n=2}^{N-1} (1 + x_n \gamma(z_n))^{2j_n} \prod_{k=1}^{\sigma_n} [\beta(v(n, k))]^{-t} \prod_{i=1}^{\rho_n} \beta(w(n, i)) | j_1 \rangle \\ &= \oint_0 \prod_{n=2}^{N-1} \frac{du_n}{2\pi i} \prod_{i=1}^{\rho_n} \prod_{k=1}^{\sigma_n} \\ &\quad \langle j_N | : \left(\beta(w(n, i)) + \frac{x_n/u_n}{w(n, i) - z_n} \right) \left(\beta(v(n, k)) + \frac{x_n/u_n}{v(n, k) - z_n} \right)^{-t} \\ &\quad \cdot u_n^{2j_n} D_n^{-2j_n + \rho_n - \sigma_n t} \exp \left(\frac{1 + x_n \gamma(z_n)}{u_n} \right) \Gamma(2j_n + 1) \frac{1}{u_n} : | j_1 \rangle \\ &= \oint_0 \prod_{n=2}^{N-1} \frac{du_n}{2\pi i} \prod_{i=1}^{\rho_n} \left(\sum_{\ell=2}^{N-1} \frac{x_\ell/u_\ell}{w(n, i) - z_\ell} \right) \prod_{k=1}^{\sigma_n} \left(\sum_{\ell=2}^{N-1} \frac{x_\ell/u_\ell}{v(n, k) - z_\ell} \right)^{-t} \\ &\quad \cdot u_n^{2j_n-1} D_n^{-2j_n + \rho_n - \sigma_n t} e^{(1/u_n)} \Gamma(2j_n + 1) \end{aligned} \quad (4.86)$$

where we have used the techniques already developed for the 3-point function. In particular, in the second equality we applied the lemma of the previous section, and in the last equality we kept doing that until normal ordering signs surrounded all operators, at which point the calculation was completed by putting β 's and γ 's under normal ordering signs equal to zero. Conforming with the discussion in the previous section we may also throw away all derivatives

on the exponential (they are with respect to the full argument of the exponential), though their presence serve to remind us in some cases, what representations would conveniently be used. In the sequel we drop these derivatives. It is straightforward to write down the contribution from the φ -part of the free field realization. It is

$$\begin{aligned}
W_N^\varphi &= \prod_{1 \leq m < n \leq N-1} (z_m - z_n)^{2j_m j_n / t} \prod_{n=2}^{N-1} \prod_{i=1}^{\rho_n} \prod_{m=1}^{N-1} (w(n, i) - z_m)^{-2j_m / t} \\
&\cdot \prod_{n=2}^{N-1} \prod_{k=1}^{\sigma_n} \prod_{m=1}^{N-1} (v(n, k) - z_m)^{2j_m} \prod_{(n, i) < (n', i')} (w(n, i) - w(n', i'))^{2/t} \\
&\cdot \prod_{(n, k) < (n', k')} (v(n, k) - v(n', k'))^{2t} \prod_{(n, i), (n', k)} (w(n, i) - v(n', k))^{-2} \quad (4.87)
\end{aligned}$$

Here we have introduced a rather arbitrary ordering of indices, for example as

$$(n, i) < (n', i') \quad (4.88)$$

if either $n < n'$ or $n = n'$, $i < i'$.

Let us summarise our findings in a more compact notation: Let

$$\begin{aligned}
M &= \sum_{m=2}^{N-1} (\rho_m + \sigma_m) \\
w_i & \quad i = 1, \dots, M \quad (4.89)
\end{aligned}$$

collectively denote the position of all screening charges

$$\{w_i\} = \{w(n, i), v(n, k)\} \quad (4.90)$$

Further let

$$\begin{aligned}
k_i &= \begin{cases} -1 & i = 1, \dots, \sum_m \rho_m \\ t & i = \sum_m \rho_m + 1, \dots, M \end{cases} \\
B(w_i) &= \sum_{\ell=1}^{N-1} \frac{x_\ell / u_\ell}{w_i - z_\ell} \quad (4.91)
\end{aligned}$$

(here $x_1 = 0$). Then the *integrand* of the N -point function is given by (we use the same letters for the integrated expressions, we hope this will not cause confusion)

$$W_N = W_B W_N^\varphi F \quad (4.92)$$

with

$$\begin{aligned}
W_N^{\beta\gamma} &= W_B F \\
W_B &= \prod_{i=1}^M B(w_i)^{-k_i} \\
W_N^\varphi &= \prod_{m < n} (z_m - z_n)^{2j_m j_n / t} \prod_{i=1}^M \prod_{m=1}^{N-1} (w_i - z_m)^{2k_i j_m / t} \prod_{i < j} (w_i - w_j)^{2k_i k_j / t} \\
F &= \prod_{m=2}^{N-1} \Gamma(2j_m + 1) u_m^{2j_m - 1} e^{1/u_m} \quad (4.93)
\end{aligned}$$

We believe the above general closed expression for integral representation of the N -point function to be useful for further development, in particular integrations over the auxiliary variables, u_ℓ , $\ell = 2, \dots, N-1$ seem tractable as they stand. If for some reason, one needs to get rid of these integrations, it is not too difficult. As an example, we provide an explicit form [110] for the result for integrable representations. First we define the following notation

$$\begin{aligned} J_N &= \{2, 3, \dots, N-1\} \\ I_N &= \{(n, i) | n = 2, \dots, N-1, i = 1, 2, \dots, \rho_n\} \\ \mathcal{F}_N &= \{\text{maps, } f, \text{ from } I_N \text{ to } J_N\} \end{aligned} \quad (4.94)$$

For $t = k + 2$ integer, all σ 's are $= 0$. In this case we may then write

$$\begin{aligned} W_N^{\beta\gamma} &= \oint_0 \prod_{n=2}^{N-1} \frac{du_n}{2\pi i} u_n^{2j_n} \frac{e^{1/u_n}}{u_n} (2j_n)! \prod_{(n,i) \in I_N} \left(\sum_{\ell=2}^{N-1} \frac{x_\ell/u_\ell}{w(n,i) - z_\ell} \right) \\ &= \oint_0 \prod_{n=2}^{N-1} \frac{du_n}{2\pi i} u_n^{2j_n} \frac{e^{1/u_n}}{u_n} (2j_n)! \sum_{f \in \mathcal{F}_N} \prod_{(n,i) \in I_N} \left(\frac{x_{f(n,i)}/u_{f(n,i)}}{w(n,i) - z_{f(n,i)}} \right) \\ &= \oint_0 \prod_{n=2}^{N-1} \frac{du_n}{2\pi i} u_n^{2j_n} \frac{e^{1/u_n}}{u_n} (2j_n)! \\ &\quad \sum_{f \in \mathcal{F}_N} \prod_{\ell \in J_N} (x_\ell/u_\ell)^{|f^{-1}(\ell)|} \prod_{(n,i) \in f^{-1}(\ell)} (w(n,i) - z_\ell)^{-1} \\ &= \sum_{f \in \mathcal{F}_N} \prod_{\ell \in J_N} x_\ell^{|f^{-1}(\ell)|} \frac{(2j_\ell)!}{(2j_\ell - |f^{-1}(\ell)|)!} \prod_{(n,i) \in f^{-1}(\ell)} (w(n,i) - z_\ell)^{-1} \end{aligned} \quad (4.95)$$

A similar but even more complicated sum formula obtains in the general case.

4.4.1 Different Realizations

We now have two ways of calculating the chiral block corresponding to N primary fields. The way so far described is by using (part of) projective invariance (to be discussed further later) and global $SL(2)$ invariance to work it out as

$$\begin{aligned} W_N^{(I)}(z_N = \infty, x_N = \infty, z_{N-1}, x_{N-1}, \dots, z_2, x_2, z_1 = 0, x_1 = 0) &= \\ \langle j_N | [\phi_{j_{N-1}}(z_{N-1}, x_{N-1})]_{\ell_{N-2}}^{j_N} \dots [\phi_{j_2}(z_2, x_2)]_{j_1}^{\ell_2} | j_1 \rangle \end{aligned} \quad (4.96)$$

However, obviously, we may also use our techniques to evaluate the same N -point chiral block as

$$\begin{aligned} &W_N^{(II)}(z_N, x_N, z_{N-1}, x_{N-1}, \dots, z_2, x_2, z_1, x_1) \\ &= \langle 0 | [\phi_{j_N}(z_N, x_N)]_{j_N}^0 [\phi_{j_{N-1}}(z_{N-1}, x_{N-1})]_{\ell_{N-2}}^{j_N} \dots [\phi_{j_2}(z_2, x_2)]_{j_1}^{\ell_2} [\phi_{j_1}(z_1, x_1)]_0^{j_1} | 0 \rangle \end{aligned} \quad (4.97)$$

We now want to demonstrate [110] that up to normalisation these expressions are equivalent in the appropriate limits. Notice that the second form involves more screening charges around the last field than the first one. We shall see that these extra screenings give rise to a constant contribution in the limit $z_N, x_N \rightarrow \infty$. But for finite z_N and x_N , unlike in the case of the

conformal minimal models, there does not seem to be any simple way of getting a conjugate field, which would get rid of the extra screening charges.

It is clear that in the limit $z_1, x_1 \rightarrow 0$, the second formulation coincides with the first, regarding the ket-part. In particular, the second formulation involves no extra screening operators. Thus we shall concentrate on the limit $z_N, x_N \rightarrow \infty$.

As before, we let w_i denote the position of screening operators *in the first case*, $W_N^{(I)}$, and we let i run over the same set as in the first case. Further, we let z_n, x_n denote the arguments as in the first case and $n = 1, 2, \dots, N-1$ runs over the same set as in the first case. The new feature in the second case is:

(i) the appearance of

$$j_N + j_N - 0 = r_N - 1 - s_N t = \rho_N - \sigma_N t$$

extra screening operators, the positions of which we denote by

$$w_{i_N}^N, \quad i_N = 1, \dots, \rho_N + \sigma_N$$

$$k_{i_N}^N = \begin{cases} -1 & , \quad i_N = 1, \dots, \rho_N \\ t & , \quad i_N = \rho_N + 1, \dots, \rho_N + \sigma_N \end{cases}$$

and

(ii) an extra u integration over a variable we call u_N . Now we want to consider the limit as $z_N, x_N \rightarrow \infty$ (letting $W_N^{(I)}$ stand for the integrand in an appropriate way) of

$$\begin{aligned} & z_N^{\frac{2j_N(j_N+1)}{t}} x_N^{-2j_N} W_N^{(II)} \\ = & z_N^{\frac{2j_N(j_N+1)}{t}} x_N^{-2j_N} \Gamma(2j_N + 1) \prod_{i_N} B^{-k_{i_N}^N}(w_{i_N}^N) \prod_{i_N < j_N} (w_{i_N}^N - w_{j_N}^N)^{2k_{i_N}^N k_{j_N}^N/t} \\ & \cdot \prod_{i, i_N} (w_{i_N}^N - w_i)^{2k_{i_N}^N k_i/t} \prod_{i_N} (z_N - w_{i_N}^N)^{2k_{i_N}^N j_N/t} \prod_{i_N, n} (w_{i_N}^N - z_n)^{2k_{i_N}^N j_n/t} \\ & \cdot \prod_n (z_N - z_n)^{2j_N j_n/t} \prod_i (z_N - w_i)^{2j_N k_i/t} u_N^{2j_N-1} e^{1/u_N} \frac{du_N}{2\pi i} W_N^{(I)} \prod_{i_N} dw_{i_N}^N \end{aligned} \quad (4.98)$$

where the function $B(w)$ is defined with one more term than for case (I), cf. (4.91). We now use that

$$\begin{aligned} - \sum_{i_N} k_{i_N}^N &= \rho_N - \sigma_N t = 2j_N \\ \sum_i k_i &= - \sum_n j_n + j_N \end{aligned} \quad (4.99)$$

In the limit $z_N \rightarrow \infty, x_N \rightarrow \infty$ we find

$$\begin{aligned} w_{i_N}^N / z_N &= \tilde{w}_{i_N}^N \\ w_i / z_N &\rightarrow 0 \\ z_n / z_N &\rightarrow 0 \end{aligned} \quad (4.100)$$

with $\tilde{w}_{i_N}^N$ finite and

$$B(w_{i_N}^N) \sim \frac{x_N/u_N}{(1 - \tilde{w}_{i_N}^N)z_N} \quad (4.101)$$

Hence, we have

$$\begin{aligned} & \lim_{z_N, x_N \rightarrow \infty} z_N^{\frac{2j_N(j_N+1)}{t}} x_N^{-2j_N} W_N^{(II)} \\ & \sim z_N^{\frac{2j_N(j_N+1)}{t}} x_N^{-2j_N} x_N^{-\sum_{i_N} k_{i_N}^N} u_N^{\sum_{i_N} k_{i_N}^N} z_N^{\sum_{i_N} k_{i_N}^N} \prod_{i_N} (1 - \tilde{w}_{i_N}^N)^{k_{i_N}^N} \\ & \cdot \prod_{i_N < j_N} (\tilde{w}_{i_N}^N - \tilde{w}_{j_N}^N)^{2k_{i_N}^N k_{j_N}^N / t} z_N^{\sum_{i_N < j_N} 2k_{i_N}^N k_{j_N}^N / t} \\ & \cdot \prod_{i_N, i} (\tilde{w}_{i_N}^N)^{2k_{i_N}^N k_i / t} z_N^{\sum_{i_N, i} 2k_{i_N}^N k_i / t} \prod_{i_N} (1 - \tilde{w}_{i_N}^N)^{2k_{i_N}^N j_N / t} z_N^{\sum_{i_N} 2k_{i_N}^N j_N / t} \\ & \cdot \prod_{i_N, n} (\tilde{w}_{i_N}^N)^{2k_{i_N}^N j_n / t} z_N^{\sum_{i_N, n} 2k_{i_N}^N j_n / t} z_N^{\sum_n 2j_N j_n / t} z_N^{\sum_i 2j_N k_i / t} \\ & \cdot u_N^{2j_N-1} e^{1/u_N} \Gamma(2j_N + 1) \frac{du_N}{2\pi i} W_N^{(I)} \prod_{i_N} d\tilde{w}_{i_N}^N z_N^{\rho_N + \sigma_N} \end{aligned} \quad (4.102)$$

(with minor misprint corrected, here and in (4.98), compared to [110]). We may evaluate the total power of z_N as

$$\begin{aligned} & \frac{2}{t} j_N(j_N + 1) - 2j_N + \frac{1}{t} [(\sum_{i_N} k_{i_N}^N)^2 - \sum_{i_N} (k_{i_N}^N)^2] + \frac{2}{t} (-2j_N)(-\sum_n j_n + j_N) \\ & + \frac{2}{t} j_N(-2j_N) + \frac{2}{t} (-2j_N) \sum_n j_n + \frac{2}{t} j_N \sum_n j_n + \frac{2}{t} j_N(-\sum_n j_n + j_N) + \rho_N + \sigma_N \\ & = 0 \end{aligned} \quad (4.103)$$

which merely shows that the intertwining field, $[\phi_{j_N}(z_N, x_N)]_{j_N}^0$ indeed has the right scaling dimension in this formalism. Similarly the total power of zero for x_N says that we treat the field with correct global $SL(2)$ properties.

Furthermore, the u_N -integrand becomes trivial, involving only

$$u_N^{-1} e^{1/u_N} \quad (4.104)$$

while the dependence on $\tilde{w}_{i_N}^N$ becomes

$$\prod_{i_N} (\tilde{w}_{i_N}^N)^{2k_{i_N}^N j_N / t} \prod_{i_N} (1 - \tilde{w}_{i_N}^N)^{k_{i_N}^N (2j_N + t) / t} \prod_{i_N < j_N} (\tilde{w}_{i_N}^N - \tilde{w}_{j_N}^N)^{2k_{i_N}^N k_{j_N}^N / t} \quad (4.105)$$

This yields an integral over the $\tilde{w}_{i_N}^N$'s which is independent of the remaining parameters of the correlator, except j_N , and thus merely contributes to the normalisation of the state $\langle j_N |$. In conclusion we note that the alternative description (4.97) (in the appropriate limits) with the initial and final states being vacua, is more symmetric and does not involve dual states (besides the bra-vacuum), but in general it involves a larger number of screening integrations.

4.4.2 The Knizhnik-Zamolodchikov Equations

One may wonder whether the rules for contractions we have put forward, really reproduce the structure of the conformal field theory. In order to settle this question in the affirmative we provide in this section an explicit proof [110] that the N -point functions satisfy the Knizhnik-Zamolodchikov equations. In this proof we should not, therefore, make any use of the rules of contractions.

The KZ equation (2.204) corresponding to the primary field at position z_{m_0} may be written as

$$\left(t\partial_{z_{m_0}} + \sum_{m \neq m_0} \frac{D_{x_{m_0}}^+ D_{x_m}^- + 2D_{x_{m_0}}^3 D_{x_m}^3 + D_{x_{m_0}}^- D_{x_m}^+}{z_m - z_{m_0}} \right) W_N = 0 \quad (4.106)$$

where W_N is the N -point function after requisite integrals have been performed.

The structure of the proof is as follows. For the selected position z_{m_0} of the primary field at that position, we shall define a function

$$G(w) = \frac{1}{w - z_{m_0}} \left(D_{x_{m_0}}^+ G^-(w) + 2D_{x_{m_0}}^3 G^3(w) + D_{x_{m_0}}^- G^+(w) \right) \quad (4.107)$$

where the $G^a(w)$'s are functions to be defined and will turn out *a posteriori* to be

$$G^a(w) = \langle J^a(w) \mathcal{O} \rangle \quad (4.108)$$

where $\mathcal{O}$ is the collection of free field realizations of all the primary fields and screening charges. Indeed we shall evaluate the $G^a(w)$'s using our contraction rules from that idea. However, the point about the proof is that the function $G(w)$ eventually written down (4.110) will only have *pole* singularities as a function of w , and will behave as $\mathcal{O}(w^{-2})$ for $w \rightarrow \infty$, and thus the sum of residues will vanish. What we shall show explicitly is that the vanishing condition for this sum of residues is precisely the KZ equation on the N -point function. This should come as no surprise since this is merely the standard technique for proving the KZ equations. The point is that in the standard proof one makes use of associativity properties of the operators, and the purpose of the proof is exactly to establish that the rules for contractions in fact conform to those.

Now it is clear how we build the functions $G^a(w)$. To use the contraction rules with the free field realizations (4.3), is very easy in our correlator, since one may establish the rules

$$\begin{aligned} \beta(w) &\rightarrow B(w) \\ \gamma(w) &\rightarrow -\sum_{i=1}^M \frac{D_{B_i}}{w - w_i} \\ -\sqrt{t/2} \partial \varphi(w) &\rightarrow \sum_{i=1}^M \frac{k_i}{w - w_i} + \sum_{m=1}^{N-1} \frac{j_m}{w - z_m} \end{aligned} \quad (4.109)$$

where $D_{B_i} = \frac{\partial}{\partial B(w_i)}$ and k_i is defined as before. Then we have (the notation here does not distinguish between integrands and integrated expressions)

$$G^+(w) = B(w) W_B W_N^\varphi F$$

$$\begin{aligned}
G^3(w) &= \left(B(w) \sum_{i=1}^M \frac{D_{B_i}}{w - w_i} + \sum_{i=1}^M \frac{k_i}{w - w_i} + \sum_{m=1}^{N-1} \frac{j_m}{w - z_m} \right) W_B W_N^\varphi F \\
G^-(w) &= \left(-B(w) \sum_{i,j=1}^M \frac{D_{B_i} D_{B_j}}{(w - w_i)(w - w_j)} + (t-2) \sum_{i=1}^M \frac{D_{B_i}}{(w - w_i)^2} \right. \\
&\quad \left. - 2 \sum_{i,j=1}^M \frac{k_i D_{B_j}}{(w - w_i)(w - w_j)} - 2 \sum_{m,j=1}^{N-1,M} \frac{j_m D_{B_j}}{(w - z_m)(w - w_j)} \right) W_B W_N^\varphi F \quad (4.110)
\end{aligned}$$

These expressions *define* the function $G(w)$ and from now on we may completely forget that they came from applying the contraction rules to certain correlators.

In the following calculations the structure of the auxiliary u integrations turns out to be very crucial. In fact, any dependence on an x_m is via the combination x_m/u_m so that we may write

$$x_m \partial_{x_m} = -u_m \partial_{u_m} \quad (4.111)$$

and subsequently do a partial integration in u_m , writing effectively

$$x_m \partial_{x_m} W_B W_N^\varphi F \sim W_B W_N^\varphi \partial_{u_m} (u_m F) \quad (4.112)$$

Let

$$D_{x_{m_0}}^- G_{z_{m_0}}^+ = D_{x_{m_0}}^- \oint_{z_{m_0}} \frac{dw}{2\pi i} \frac{1}{w - z_{m_0}} G^+(w) \quad (4.113)$$

denote the contribution to the pole residue in $G(w)$ at $w = z_{m_0}$ coming from the term $D_{x_{m_0}}^- G^+(w)/(w - z_{m_0})$, etc. Then we find after some calculations for the pole at z_{m_0}

$$\begin{aligned}
D_{x_{m_0}}^- G_{z_{m_0}}^+ &= \sum_{\ell \neq m_0} \frac{x_\ell/u_\ell}{z_{m_0} - z_\ell} \partial_{x_{m_0}} W_B \cdot W_N^\varphi F \\
2D_{x_{m_0}}^3 G_{z_{m_0}}^3 &= 2 \left(- \sum_{\ell \neq m_0} \frac{x_\ell/u_\ell}{z_{m_0} - z_\ell} u_{m_0} \partial_{x_{m_0}} W_B \cdot W_N^\varphi \right. \\
&\quad \left. - \partial_{z_{m_0}} W_B \cdot W_N^\varphi + \frac{t}{2j_{m_0}} W_B \cdot \partial_{z_{m_0}} W_N^\varphi \right) (-j_{m_0} + u_{m_0}^{-1}) F \\
D_{x_{m_0}}^+ G_{z_{m_0}}^- &= \left(- \sum_{\ell \neq m_0} \frac{x_\ell/u_\ell}{z_{m_0} - z_\ell} \partial_{x_{m_0}} W_B \cdot W_N^\varphi (2j_{m_0} u_{m_0}^{2j_{m_0}} - u_{m_0}^{2j_{m_0}-1}) \right. \\
&\quad - 2\partial_{z_{m_0}} W_B \cdot W_N^\varphi ((2j_{m_0} - 1)u_{m_0}^{2j_{m_0}-1} - u_{m_0}^{2j_{m_0}-2}) \\
&\quad + (t - 2 + 2j_{m_0}) \partial_{z_{m_0}} W_B \cdot W_N^\varphi u_{m_0}^{2j_{m_0}-1} \\
&\quad + \frac{t}{j_{m_0}} W_B \cdot \partial_{z_{m_0}} W_N^\varphi (2j_{m_0} u_{m_0}^{2j_{m_0}-1} - u_{m_0}^{2j_{m_0}-2}) \Big) \\
&\quad \cdot e^{1/u_{m_0}} \Gamma(2j_{m_0} + 1) \prod_{m \neq m_0} \Gamma(2j_m + 1) u_m^{2j_m-1} e^{1/u_m} \quad (4.114)
\end{aligned}$$

This sums up to become

$$t \partial_{z_{m_0}} W_N \quad (4.115)$$

which is the first term in the KZ equation.

In a similar fashion the pole residue at $w = w_j$ may be calculated. However, here we shall be more general and show that substituting $1/(w - z_{m_0})$ in the definition (4.107) of $G(w)$ by any function $f(w)$ (free to depend on e.g. z_m but independent of w_j), leads to

$$\begin{aligned} G_f(w) &= f(w) \left(D_{x_{m_0}}^+ G^-(w) + 2D_{x_{m_0}}^3 G^3(w) + D_{x_{m_0}}^- G^+(w) \right) \\ \oint_{w_j} \frac{dw}{2\pi i} G_f(w) &= D_{x_{m_0}}^+ \oint_{w_j} \frac{dw}{2\pi i} f(w) G^-(w) \\ &= t \partial_{w_j} \left(D_{x_{m_0}}^+ \left(f(w_j) \frac{W_B W_N^\varphi F}{B(w_j)} \right) \right) \end{aligned} \quad (4.116)$$

The first equality in the computation of the residue is easily verified, while the second goes as follows

$$\begin{aligned} \oint_{w_j} \frac{dw}{2\pi i} f(w) G^-(w) &= \left(-2B(w_i) f(w_j) \sum_{i \neq j} \frac{D_{B_i} D_{B_j}}{(w_j - w_i)} - \partial_w [f(w) B(w)]_{w_j} D_{B_j}^2 \right. \\ &\quad + (t - 2) f'(w_j) D_{B_j} - 2 \sum_{i \neq j} \frac{k_i D_{B_j} + k_j D_{B_i}}{w_j - w_i} f(w_j) - 2 f'(w_j) k_j D_{B_j} \\ &\quad \left. - 2 \sum_m \frac{j_m D_{B_j}}{w_j - z_m} f(w_j) \right) W_B W_N^\varphi F \\ &= t \partial_{w_j} \left(f(w_j) \frac{W_B W_N^\varphi F}{B(w_j)} \right) \end{aligned} \quad (4.117)$$

where we have used that

$$\begin{aligned} \partial_{w_j} W_N^\varphi &= \left(\sum_m \frac{2j_m k_j / t}{w_j - z_m} + \sum_{i \neq j} \frac{2k_i k_j / t}{w_j - w_i} \right) W_N^\varphi \\ \partial_{w_j} W_B &= -\frac{k_j}{B(w_j)} B(w_{j'}) W_B \end{aligned} \quad (4.118)$$

and that for both $k_j = -1$ and $k_j = t$

$$k_j(k_j + 1) - t k_j = t \quad (4.119)$$

Putting $f(w) = 1/(w - z_{m_0})$, we may now conclude that upon integration over the positions w_j , the contribution from the poles at w_j vanishes.

Finally, the pole residues at the points $w = z_m \neq z_{m_0}$ give the remaining terms in the KZ equation (4.106). Since there are no other singularities in w , the KZ equation has been proven.

The above proof assumes that the integration contours of the screening charges are closed. When parameters are such that this effectively is not the case, it is no longer true that the contribution from the total derivatives met at $w = w_j$ may be thrown away. However, in that case the derivative after z_{m_0} contains an extra term coming from varying the end point of the integration contour, and it is possible to check that this extra contribution exactly cancels the contribution from the total derivative term, so that the validity of the KZ equations remains true in those cases as well.

4.4.3 Projective Invariances

In [34] it is shown, that solutions to the KZ equations are projectively invariant provided the primary fields can add up to a singlet. In our case this is the requirement of global (loop projective) $SL(2)$ invariance. Let us restrict to the case where the initial and final bra and ket carry just the vacuum and the dual vacuum: $j_1 = 0 = j_N$, i.e. we are really looking at an $(N - 2)$ -point function. Then global $SL(2)$ invariance is the statement that

$$\sum_{m=2}^{N-1} D_{x_m}^a W_{N-2} = 0 \quad (4.120)$$

This is equivalent to the statement that in

$$G^a(w) = \langle J^a(w) \mathcal{O} \rangle \quad (4.121)$$

the leading behaviour as $w \rightarrow \infty$ is $\mathcal{O}(w^{-2})$ rather than $\mathcal{O}(w^{-1})$. From the expressions above, that is trivial for $G^-(w)$, and for $G^3(w)$ it follows from the fact that

$$\sum_{i=1}^M k_i = - \sum_{m=1}^{N-1} (\rho_m - t\sigma_m) = - \sum_{m=1}^{N-1} j_m \quad (4.122)$$

for $j_N = 0$. For $G^+(w)$ it is more complicated to see. As previously discussed this is related to the fact that we are not using the projectively invariant vacuum $\langle sl_2 |$ in our calculations, rather we are using the dual vacuum $\langle 0 |$ for which $\langle 0 | J_0^+ = \langle 0 | \beta_0 \neq 0$.

What we are going to show [110] is that the state

$$\langle 0 | \beta_0 \quad (4.123)$$

even though it is non-vanishing, is BRST exact in the sense of Felder [56], and that (hence) it decouples from correlators of BRST invariant operators.

First let us argue at the operator level, and subsequently at the level of our correlators. We write

$$\langle 0 | \beta_0 = \langle 0 | e^{-\sqrt{2/t}\varphi} e^{\sqrt{2/t}\varphi} \beta_0 = \langle -1 | \oint \frac{dz}{2\pi i} : e^{\sqrt{2/t}\varphi(z)} : \beta(z) \quad (4.124)$$

where the bra state $\langle -1 |$ is the lowest weight state $\langle j = -1 |$ (4.13). Now this integral is in fact the appropriate BRST operator in Felder's formulation. To see this, recall, that acting on the Fock space pertaining to $j_{r,s}$ and labelled $F_{r,s}$, the relevant BRST operator [56] (for which the BRST current is single valued) is

$$Q_r \sim \oint \frac{dv_0}{2\pi i} \dots \frac{dv_{r-1}}{2\pi i} S_1(v_0) \dots S_1(v_{r-1}) \quad (4.125)$$

with

$$Q_r : F_{r,s} \mapsto F_{-r,s} \quad (4.126)$$

For $j_N = 0$, this is the Fock space with $r = 1, s = 0$ ($2j_N + 1 = 1 - 0 \cdot t$). Therefore the relevant BRST operator on this space is Q_1 which is just the one we obtained (4.124).

Next let us see how the argument works at the level of the correlator. We are going to show that inserting the operator

$$\beta_0 = \oint_{\infty} \frac{dw}{2\pi i} \beta(w) \quad (4.127)$$

furthest to the left in a correlator with $j_N = 0$, is equivalent to inserting the BRST charge operator

$$\oint_{\infty} \frac{dw}{2\pi i} \beta(w) : e^{\sqrt{2/t}\varphi(w)} : \quad (4.128)$$

furthest to the left of the operators and to the right of the bra $\langle j_N = -1|$. Indeed in the first case, using the rules for building correlators, we obtain (up to normalisation)

$$\begin{aligned} & \oint_{\infty} \frac{dw}{2\pi i} B(w) \prod_i B(w_i)^{-k_i} \prod_{i < j} (w_i - w_j)^{2k_i k_j / t} \\ & \cdot \prod_{n < m} (z_n - z_m)^{2j_n j_m / t} \prod_{i, m} (w_i - z_m)^{2k_i j_m / t} \prod_m u_m^{2j_m - 1} e^{1/u_m} \end{aligned} \quad (4.129)$$

In the second case, we notice that the state labelled $\langle -1|$ is exactly the lowest weight state $\langle j_N = -1|$ and we are formally looking at an $(N - 1)$ -point function with $j_N = -1$. Since we have

$$\iota_{N-2} + j_{N-1} - j_N = \rho_{N-1} - \sigma_{N-1} t \quad (4.130)$$

we see that the value of ρ_{N-1} will be one unit larger in the case $j_N = -1$ than in the case $j_N = 0$. This means we have one extra screening charge of the first kind in that case, which we may 'lift off' the intertwining field furthest to the left, in other words that calculation will just be the one we seek to carry out. Using the rules developed we find in that case, letting w denote the position of the extra screening operator compared to the previous case (so in the formulas below, the index i runs over exactly the same set as before)

$$\begin{aligned} & \oint_{\infty} \frac{dw}{2\pi i} B(w) \prod_i B(w_i)^{-k_i} \prod_i (w - w_i)^{-2k_i / t} \prod_n (w - z_n)^{-2j_n / t} \prod_{i < j} (w_i - w_j)^{2k_i k_j / t} \\ & \cdot \prod_{n < m} (z_n - z_m)^{2j_n j_m / t} \prod_{i, n} (w_i - z_n)^{2k_i j_n / t} \prod_n u_n^{2j_n - 1} e^{1/u_n} \\ & = \oint_{\infty} \frac{dw}{2\pi i} B(w) \prod_i B(w_i)^{-k_i} \prod_i (1 - w_i/w)^{-2k_i / t} \prod_n (1 - z_n/w)^{-2j_n / t} w^{-2(\sum_i k_i + \sum_n j_n)/t} \\ & \cdot \prod_{i < j} (w_i - w_j)^{2k_i k_j / t} \prod_{n < m} (z_n - z_m)^{2j_n j_m / t} \prod_{i, n} (w_i - z_n)^{2k_i j_n / t} \prod_n u_n^{2j_n - 1} e^{1/u_n} \end{aligned} \quad (4.131)$$

We now use that

$$-\sum_i k_i = \sum_n \rho_n - t \sum_n \sigma_n = \sum_n j_n - j_N \quad (4.132)$$

Since we use the notation that sums over i and n are pertaining to the case with $j_N = 0$, we see that the extra power of w becomes zero, and for a very large integration contour for w , all w dependence drops out except for the one in $B(w)$ so that we exactly prove the identity of the two cases also at the level of our correlators.

Having come this far, we may move the Felder type BRST operator through all intertwining fields in exactly the same way as for minimal models [56], until in the end it hits the ket-vacuum. Since we are only using screening operators of the first kind in the BRST operators, the procedure will work just as for the minimal models.

In addition it is rather easy to verify that the functions $G^a(w)$ have the expected pole residues for $\langle J^a(w)\mathcal{O} \rangle$ at points $w = z_m$. Further one verifies that there are no pole residues in $G^+(w)$ and $G^3(w)$ for $w = w_i$, the position of a screening charge. For $G^-(w)$ that residue is proportional to the total derivative

$$\partial_{w_i}(D_{B_i}W_B \cdot W_N^\varphi) \quad (4.133)$$

All of those remarks establish that the N -point chiral blocks have the correct projective and global $SL(2)$ invariance properties.

4.4.4 Possible Hybrid

This section is devoted to a very brief discussion of a generalisation [110] of a work by Malikov *et al* [54] in which a class of solutions to the KZ equations are considered.

Without going into the precise mathematical language employed in [54] it is easy enough to describe the approach by these authors in terms of ours. Given some number of primary fields characterised by spins $j_1, j_2, \dots, j_{N-1}$ they begin by supplementing this set by an auxiliary primary field of spin

$$j_N^{(0)} = \sum_{i=1}^{N-1} j_i \quad (4.134)$$

carefully chosen so that no screening charges are required in the free field approach. In general, writing

$$\rho - \sigma t = j_N^{(0)} - j_N \quad (4.135)$$

our approach is applicable whenever ρ and σ are non-negative integers, in which case they represent the number of screening charges needed of the two kinds. Obviously now, for $j_N = j_N^{(0)}$ no screening charges are needed, then the correlators are independent of the x_i 's and may be trivially written down. This expression is the starting point in [54]. From this a larger and more interesting class of solutions to the KZ equations is obtained as follows. From the state $|j_{r,s}\rangle$ it is well-known (4.50) how to build the singular vectors in the Verma module generated by this highest weight state, simply by multiplying $|j_{r,s}\rangle$ by certain products of generators. Thus when this operator is inserted into a correlator the state decouples and the correlator vanishes. The trick in [54] is to only make use of a subset of the operators, starting from the right and truncating the expansion at some point to the left. Each factor performs a Weyl reflection and it may easily be shown that any subset generates a formally singular state, although of course not one that lies in the Verma module of $|j_{r,s}\rangle$. Hence, inserting such a reduced set of operators will not lead to decoupling, but instead to a new solution of the KZ equations [54].

What we shall argue now [110], however, is that this procedure can never exhaust the possibilities we wish to consider in conformal field theory. The point is that when this procedure is applied to the state $\langle j_N^{(0)} |$ the new spin value for the state may only be modified relative to $j_N^{(0)}$ in a way which is insufficient for producing all the spins needed by the fusion rules in a

theory based on admissible representations. Indeed keeping m operators results in a new spin $j_N^{(m)}$ given by

$$\begin{aligned} j_N^{(2n)} &= j_N^{(0)} + nt \\ j_N^{(2n+1)} &= j_N^{(0)} - r_N^{(0)} + (s_N^{(0)} - n)t \end{aligned} \quad (4.136)$$

for m even and odd respectively. We see that even though j_N is not required to be equal to $j_N^{(0)}$ the possibilities are too limited to allow considerations of all the relevant spin values in an admissible representation: relative to $j_N^{(0)}$ only the s value can be changed at will, the r value is fixed either at its value in $j_N^{(0)}$ or at minus that value. The above spins may be treated also in the formalism based on free fields and screening operators. Indeed the number of screening operators are given by

$$\begin{aligned} \rho_{2n} &= 0 \\ \sigma_{2n} &= n \\ \rho_{2n+1} &= r_N^{(0)} \\ \sigma_{2n+1} &= s_N^{(0)} - n \end{aligned} \quad (4.137)$$

This is the number of integrations in the integral formulas (in addition to the simpler ones having to do with u integrations).

A possibly more interesting hybrid treatment based on the free field realization and [54] might be possible. In fact, if instead of starting with correlators needing no screening charges, one starts with correlators needing only the first kind of screening charge for which extensive studies exist in the literature (see e.g. [22, 41]), it would seem that the techniques of [54] could provide a way alternative to the one discussed in the preceding sections to circumventing the use of the second screening operator. The result would be a strange mixture of free field integral expressions and the ones considered in [54], but it should agree with the ones given earlier up to normalisation. It is not clear that such hybrid integral representations have advantages over what has otherwise been presented here, but at least they would also be complete, contrary to the ones in [54].

4.5 Hamiltonian Reduction

Here we want to discuss [111] how conformal blocks in $SL(2)$ current algebra reduce to corresponding ones in minimal models.

A general conformal block (on the sphere) in the affine theory is given by

$$W_N = \langle j_N | \phi_{j_{N-1}}(z_{N-1}, x_{N-1}) \dots \phi_{j_n}(z_n, x_n) \dots \phi_{j_2}(z_2, x_2) \prod_i \int dw_i S_{k_i}(w_i) | j_1 \rangle \quad (4.138)$$

Different choices of integration contours for the screening charges define different intertwining chiral vertex operators and different conformal blocks. The two screening charges are

$$\begin{aligned} S_{k_{\pm}}(w) &= \beta^{-k_{\pm}}(w) S_{k_{\pm}}^{\varphi}(w) \\ S_{k_{\pm}}^{\varphi}(w) &= : e^{-k_{\pm} \sqrt{2/t} \varphi(w)} : \\ k_+ &= -1 \\ k_- &= t \end{aligned} \quad (4.139)$$

The relation to minimal models is obtained by writing [24]

$$\begin{aligned}
2j_{r,s} + 1 &= r - st \\
t &= k + 2 = p/q \\
\alpha_+ &= \sqrt{\frac{2}{t}} = -2/\alpha_- \\
\alpha_{r,s+1} &= -j_{r,s} \sqrt{\frac{2}{t}} = \frac{1}{2}((1-r)\alpha_+ - s\alpha_-) \\
2\alpha_0 &= \alpha_+ + \alpha_- \\
\Delta_{r,s+1} &= \frac{j_{r,s}(j_{r,s} + 1)}{t} - j_{r,s} = \frac{1}{2}\alpha_{r,s+1}(\alpha_{r,s+1} - 2\alpha_0) \\
\phi_{r,s+1}(z) &= : e^{\alpha_{r,s+1}\varphi(z)} := \phi_{j_{r,s}}(z) \\
V_{\alpha_{\pm}}(w) &= : e^{\alpha_{\pm}\varphi(w)} := S_{k_{\pm}}^{\varphi}(w)
\end{aligned} \tag{4.140}$$

It is now clear that if one truncates the β dependence of the screening currents and the γ dependent factor in the primary fields, then the minimal model correlators are obtained [64]. This is true despite the fact that the two theories, the WZNW model and the minimal model, have different background charges for the φ field: namely $-\alpha_+ = -\sqrt{\frac{2}{t}}$ (see (4.6)) for the WZNW model and $-2\alpha_0 = -\sqrt{\frac{2}{t}} + \sqrt{2t}$ for the minimal models (see Chapter 2). However, this difference is of no consequence in the practical evaluation of the free field correlators since in both cases suitable dual bra-states are used to absorb those background charges. In the following, we want to discuss how the Hamiltonian reduction works at the level of correlators. Thereby we prove [111] the statement in [64] that the correlators in $SL(2)$ current algebra reduce to corresponding ones in a particular minimal model in the limit where all x 's are put equal to the corresponding z 's.

4.5.1 At the Level of Correlators

The primary field (4.3) may be written as

$$\begin{aligned}
\phi_j(z, x) &= e^{x\partial_y} \phi_j(z, y)|_{y=0} \\
&= e^{xD_y^-} \phi_j(z, y)|_{y=0} \\
&= e^{xJ_0^-} \phi_j(z, 0) e^{-xJ_0^-} \\
&= e^{xJ_0^-} : e^{-j\sqrt{2/t}\varphi(z)} : e^{-xJ_0^-}
\end{aligned} \tag{4.141}$$

Here there is a subtlety in that the way the exponential function $e^{xJ_0^-}$ should be expanded, must also respect the monodromy conditions in the x variables. We shall come back to this subtlety presently. We may further write

$$\begin{aligned}
\phi_j(z, xz) &= e^{zxJ_0^-} : e^{-j\sqrt{2/t}\varphi(z)} : e^{-zxJ_0^-} \\
&= e^{xzD_y^-} \phi_j(z, y)|_{y=0} \\
&= e^{xJ_1^-} : e^{-j\sqrt{2/t}\varphi(z)} : e^{-xJ_1^-}
\end{aligned} \tag{4.142}$$

and

$$\phi_j(z, x) = e^{xJ_0^-} e^{zL_{-1}} : e^{-j\sqrt{2/t}\varphi(0)} : e^{-xJ_0^-} e^{-zL_{-1}} \quad (4.143)$$

Consider the following conformal blocks

$$W_N = \langle j_N | \phi_{j_{N-1}}(z_{N-1}, xz_{N-1}) \dots \phi_{j_n}(z_n, xz_n) \dots \phi_{j_2}(z_2, xz_2) \prod_i \int dw_i S_{k_i}(w_i) | j_1 \rangle \quad (4.144)$$

Substituting (4.142) we may rewrite this as

$$W_N = \langle j_N | e^{xJ_1^-} \phi_{j_{N-1}}(z_{N-1}, 0) \dots \phi_{j_n}(z_n, 0) \dots \phi_{j_2}(z_2, 0) \prod_i \int dw_i S_{k_i}(w_i) | j_1 \rangle \quad (4.145)$$

since $J_1^- | j_1 \rangle = 0$. At this point however, there is a subtlety as to how adjacent exponentials $e^{\pm xJ_1^-}$ should be removed, and we should examine how these exponentials are defined. Indeed as discussed earlier the expansions of exponentials and other functions involving the β and γ fields, depend on which monodromy the problem at hand requires one to select. All these subtleties are dealt with using the following two lemmas [111]

Lemma 1

If the fractional part in powers of x is α , then we can expand the last expression in (4.141)

$$e^{xJ_0^-} : e^{-j\sqrt{2/t}\varphi(z)} : e^{-xJ_0^-} = \sum_{n \in \mathbb{Z}} \frac{(xJ_0^-)^{\alpha-\beta+n}}{(\alpha-\beta+n)!} : e^{-j\sqrt{2/t}\varphi(z)} : \sum_{m \in \mathbb{Z}} \frac{(-xJ_0^-)^{\beta+m}}{(\beta+m)!} \quad (4.146)$$

for arbitrary complex number β .

Lemma 2

$$1 = \sum_{n \in \mathbb{Z}} \frac{(xJ_0^-)^{\alpha+n}}{(\alpha+n)!} \sum_{m \in \mathbb{Z}} \frac{(-xJ_0^-)^{-\alpha+m}}{(-\alpha+m)!} = e^{xJ_0^-} e^{-xJ_0^-} \quad (4.147)$$

Before proving these lemmas we make the following remarks. Define

$$\begin{aligned} \phi_j^{[n]}(z, 0) &= [J_0^-, \phi_j^{[n-1]}(z, 0)] \\ \phi_j^{[0]}(z, 0) &= \phi_j(z, 0) \end{aligned} \quad (4.148)$$

When x is integrally powered, it is clear that we can expand $\phi_j(z, x)$ as

$$\begin{aligned} e^{xJ_0^-} : e^{-j\sqrt{2/t}\varphi(z)} : e^{-xJ_0^-} &= \sum_{n \geq 0} \frac{\phi_j^{[n]}(z, 0) x^n}{n!} \\ &= \sum_{n \geq 0} \frac{(D_y^-)^n \phi_j(z, y) x^n}{n!} \Big|_{y=0} \\ &= e^{xD_y^-} \phi_j(z, y) \Big|_{y=0} \\ &= \phi_j(z, x) \end{aligned} \quad (4.149)$$

However, when x is fractionally powered, we can no longer Taylor expand $\phi_j(z, x)$, and the definition for both $\phi_j^{[n]}(z, 0)$ in (4.148) and $(D_y^-)^n \phi_j(z, y) \Big|_{y=0}$ in (4.149) requires specification.

It is possible to generalise (4.148) and (4.149) by defining

$$\frac{\phi_j^{[N+\alpha+\beta]}(z, 0)}{(N+\alpha+\beta)!} = \sum_{\substack{n+m=N \\ n, m \in \mathbb{Z}}} \frac{(J_0^-)^{\alpha+n}}{(\alpha+n)!} : e^{-j\sqrt{2/t}\varphi(z)} : \frac{(-J_0^-)^{\beta+m}}{(\beta+m)!} \quad (4.150)$$

Indeed, this is seen to satisfy the recursion relation in (4.148). Although it looks like that the rhs of (4.150) depends on both α and β , lemma 1 essentially means that it only depends on the combination $\alpha+\beta$. The fractional derivatives at the origin may also be considered as analytical continuations of their integral counterparts. Now $\phi_j(z, x) = (1 + \gamma(z)x)^{2j} : e^{-j\sqrt{2/t}\varphi(z)} :$, so for non-negative integer n we have

$$\frac{(D_y^-)^n}{n!} \phi_j(z, y)|_{y=0} = \binom{2j}{n} \gamma^n(z) : e^{-j\sqrt{2/t}\varphi(z)} : \quad (4.151)$$

We can analytically continue the variable n in the above equation from integers to complex numbers. Therefore n could be any fractional number and we have

$$\frac{(D_y^-)^{n+\alpha}}{(n+\alpha)!} \phi_j(z, y)|_{y=0} = \binom{2j}{n+\alpha} \gamma^{n+\alpha}(z) : e^{-j\sqrt{2/t}\varphi(z)} : \quad (4.152)$$

Proof of Lemma 1

$$\begin{aligned} & \sum_{n \in \mathbb{Z}} \frac{(xJ_0^-)^{\alpha-\beta+n}}{(\alpha-\beta+n)!} : e^{-j\sqrt{2/t}\varphi(z)} : \sum_{m \in \mathbb{Z}} \frac{(-xJ_0^-)^{\beta+m}}{(\beta+m)!} \\ &= \sum_{n \in \mathbb{Z}} \frac{(xJ_0^-)^{\alpha-\beta+n}}{(\alpha-\beta+n)!} \sum_{m \in \mathbb{Z}} \frac{(-xJ_0^- + xD_y^-)^{\beta+m}}{(\beta+m)!} \phi(z, y)|_{y=0} \\ &= \sum_{N \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \frac{(xJ_0^-)^{\alpha-\beta+N-m} (-xJ_0^- + xD_y^-)^{\beta+m}}{(\alpha-\beta+N-m)! (\beta+m)!} \phi(z, y)|_{y=0} \\ &= \sum_{N \in \mathbb{Z}} \frac{(xD_y^-)^{\alpha+N}}{(\alpha+N)!} \phi(z, y)|_{y=0} \\ &= \sum_{N \in \mathbb{Z}} \binom{2j}{N+\alpha} (\gamma(z)x)^{N+\alpha} : e^{-j\sqrt{2/t}\varphi(z)} : \\ &= \phi_j(z, x) \end{aligned} \quad (4.153)$$

□

Proof of Lemma 2

$$\begin{aligned} e^{xJ_0^-} e^{-xJ_0^-} &= \sum_{n \in \mathbb{Z}} \frac{(xJ_0^-)^{\alpha+n}}{(\alpha+n)!} \sum_{m \in \mathbb{Z}} \frac{(-xJ_0^-)^{-\alpha+m}}{(-\alpha+m)!} \\ &= \sum_{N \in \mathbb{Z}} \sum_{n+m=N} \frac{(xJ_0^-)^N}{(\alpha+n)! (-\alpha+m)!} (-1)^{-\alpha+m} \\ &= \sum_{N \in \mathbb{Z}} (xJ_0^-)^N \delta_{N,0} \\ &= 1 \end{aligned} \quad (4.154)$$

□

Thus the manipulations leading to (4.145) are justified. In the proofs of the lemmas we have used versions of (2.112).

We may now go back to (4.145). We see from that expression that the exponential

$$e^{xJ_1^-} \quad (4.155)$$

has to be expanded in powers offset from the integers by the amount

$$-\sum k_i = r - st = \alpha \quad (4.156)$$

with r and s the number of screening charges of the first and second kind respectively, so this is the combined power of the β factors. Since the γ factors from the primary fields decouple for $x_i = 0$, this particular expansion of the exponential is required. One may then work out that [111] (here corrected for minor misprint)

$$\langle j_N | e^{xJ_1^-} = \langle j_N | (1 - x\gamma_1)^{k-2j_N} \frac{s(k-2j_N)}{s(k-2j_N-\alpha)} (-1)^{-\alpha} \quad (4.157)$$

Notice that for α integer the ratio of sine-factors along with the phase disappear. The result (4.157) is obtained by writing (for $\alpha = r - st$)

$$e^{xJ_1^-} = \sum_{n \in \mathbb{Z}} \frac{(xJ_1^-)^{n+\alpha}}{(n+\alpha)!} \quad (4.158)$$

and observing that for any power

$$\langle j_N | (J_1^-)^{n+\alpha} = \langle j_N | \gamma_1^{n+\alpha} \frac{\Gamma(2j_N - k + n + \alpha)}{\Gamma(2j_N - k)} \quad (4.159)$$

The proportionality to a power of γ_1 follows from the free field realization and the properties of the vacuum. The value of the constant is obtained by consistency between $n + \alpha$ and $n + \alpha + 1$ and by normalising with the result for $n + \alpha = 1$.

We may now continue the calculation and obtain

$$\begin{aligned} W_N &= \langle j_N | (1 - x\gamma_1)^{k-2j_N} \frac{s(k-2j_N)}{s(k-2j_N-\alpha)} (-1)^{-\alpha} \\ &\quad \cdot \phi_{j_{N-1}}(z_{N-1}, 0) \dots \phi_{j_n}(z_n, 0) \dots \phi_{j_2}(z_2, 0) \prod_i \int dw_i S_{k_i}(w_i) |j_1\rangle \\ &= C_N(\{j_m\}, x) \langle j_N | \phi_{j_{N-1}}(z_{N-1}, 0) \dots \phi_{j_n}(z_n, 0) \dots \phi_{j_2}(z_2, 0) \prod_i \int dw_i S_{k_i}^\varphi(w_i) |j_1\rangle \\ &= C_N(\{j_m\}, x) W_N^\varphi \end{aligned} \quad (4.160)$$

where $C_N(\{j_m\}, x)$ is the normalisation constant, and W_N^φ is exactly the free field expression for the minimal model correlator

$$W_N^\varphi = \langle \Delta_{r_N, s_N+1} | \phi_{r_{N-1}, s_{N-1}+1}(z_{N-1}) \dots \phi_{r_2, s_2}(z_2) \prod_i \int dw_i V_{\alpha_i}(w_i) | \Delta_{r_1, s_1+1} \rangle \quad (4.161)$$

The point is that since (4.157) now contains the only γ dependence of the correlator, the β dependence is effectively removed from the screening charges since γ_1 interacts only with β_{-1} which is the *constant* (w_i independent) mode. Thus [111]

$$\begin{aligned} C_N(\{j_m\}, x) &= \frac{\Gamma(k - 2j_N + 1)}{\Gamma(k - 2j_N - r + st + 1)} \frac{s(k - 2j_N)}{s(k - 2j_N - r + st)} (-1)^{-r+st} x^{r-st} \\ &= \frac{\Gamma(2j_N - k + r - st)}{\Gamma(2j_N - k)} (-1)^{-r+st} x^{r-st} \end{aligned} \quad (4.162)$$

Of course, this may also be obtained without using (4.157) but inserting (4.158) and (4.159) directly into the correlator. In either way we use that

$$\begin{aligned} \langle 0 | [\gamma_1^\alpha, \beta_{-1}^\alpha] \dots | 0 \rangle &= (-1)^{-\alpha} \Gamma(\alpha + 1) \langle 0 | \dots | 0 \rangle \\ \langle 0 | [\beta_{-1}^\alpha, \gamma_1^\alpha] \dots | 0 \rangle &= (-1)^{-\alpha+1} \Gamma(\alpha + 1) \langle 0 | \dots | 0 \rangle \end{aligned} \quad (4.163)$$

where $\dots$ represents β, γ independent insertions. (4.163) is justified by

$$\begin{aligned} [\gamma_1, \beta_{-1}^\alpha] &= -\alpha \beta_{-1}^{\alpha-1}, \quad [\beta_{-1}, \gamma_1^\alpha] = \alpha \gamma_1^{\alpha-1} \\ \langle 0 | [\gamma_1^n, \beta_{-1}^m] \dots | 0 \rangle &= (-1)^{-n} \frac{\Gamma(m+1)}{\Gamma(m-n+1)} \langle 0 | \beta_{-1}^{m-n} \dots | 0 \rangle = (-1)^{-n} \Gamma(n+1) \delta_{nm} \end{aligned} \quad (4.164)$$

This concludes the simple proof of the statement that correlators in $SL(2)$ current algebra reduce to corresponding ones in a particular minimal model in the limit where all x 's are put equal to the corresponding z 's (4.138).

For $x = 1$ one may check from the 3-point function (4.61) that (4.162) is indeed the relative constant to the 3-point function of minimal models (4.76), up to an irrelevant phase (which may be taken care of by reverting to the phase convention in Chapter 3 for the screening currents, cf. the comment following (4.3)). This is due to the following rewriting of (4.162)

$$C_3(j_1, j_2, j_3; x = 1) = \frac{\Gamma(2j_2 + 1)}{\Gamma(2j_2 - r + st + 1)} \frac{\mathcal{J}_{rs}(2j_1, 2j_2 + t; t)}{\mathcal{J}_{rs}(2j_1, 2j_2; t)} (-1)^{-r+st} \quad (4.165)$$

and the fact that the pre-factors from the contour deformations are identical in the two 3-point functions. This is also true for the other choices of contours.

4.5.2 Analysis of Explicit Correlators

Here we want to verify explicitly that the conformal blocks for the WZNW model evaluated in the preceding sections satisfy the results above. To this end we consider the 'interpolating' correlator

$$\langle j_N | e^{-x_N J_1^-} \phi_{j_{N-1}}(z_{N-1}, x_{N-1}) \dots \phi_{j_2}(z_2, x_2) \prod_{i=1}^M \int dw_i S_{k_i}(w_i) | j_1 \rangle = \langle j_N | \mathcal{O} | j_1 \rangle \quad (4.166)$$

with

$$\begin{aligned} x_\ell &= z_\ell x, \quad \ell = 1, \dots, N-1 \\ x_N &= x - 1 \end{aligned} \quad (4.167)$$

Thus for $x = 1$ we get the WZNW model with all x_i 's put equal to the z_i 's. For $x = 0$ we should get the minimal model correlator up to normalisation. We wish to show that this interpolating correlator is *independent* of x . Using notation of preceding sections, we find

$$\mathcal{O} = W_B W_N^\varphi F \quad (4.168)$$

where

$$\begin{aligned} B(w) &= \sum_{\ell=1}^{N-1} \frac{x_\ell/u_\ell}{w - z_\ell} - x_N/u_N \\ W_B &= \prod_{i=1}^M B(w_i)^{-k_i} \\ F &= \Gamma(k - 2j_N + 1) u_N^{k-2j_N-1} e^{1/u_N} \prod_{\ell=2}^{N-1} \Gamma(2j_\ell + 1) u_\ell^{2j_\ell-1} e^{1/u_\ell} \\ W_N^\varphi &= \prod_{m < n} (z_m - z_n)^{2j_m j_n/t} \prod_{i=1}^M \prod_{m=1}^{N-1} (w_i - z_m)^{2k_i j_m/t} \prod_{i < j} (w_i - w_j)^{2k_i k_j/t} \end{aligned} \quad (4.169)$$

Here we used that

$$\begin{aligned} \langle j_N | e^{-x_N J_1^-} &= \langle j_N | (1 + x_N \gamma_1)^{k-2j_N} \\ &= \lim_{z' \rightarrow \infty} \langle j_N | (1 + x_N z' \gamma(z'))^{k-2j_N} \end{aligned} \quad (4.170)$$

so $x_N z'$ plays the role of the x in the additional term $(1 + x_N z' \gamma(z'))^{k-2j_N}$ of the type of the γ part in a primary field with spin $2j = k - 2j_N$. This certainly gives the contribution

$$\lim_{z' \rightarrow \infty} \frac{x_N z'/u_N}{w - z'} = -x_N/u_N \quad (4.171)$$

in $B(w)$. The integrations over the auxiliary variables u_ℓ are understood in (4.168).

The above expression (4.170) may seem in contradiction to the expansion (4.157), since the ratio of sine-functions we had there is absent now. So let us explain the reason for this subtlety. The point is that for $x \neq 0$ the fractional powers of the β 's in the correlator is balanced by the fractional powers of the γ 's in the primary fields. Thus the exponential in the present case has to be expanded in *integral* powers. This is in contrast to the situation in (4.157) where the γ dependence of the primary fields were suppressed since they involved the case $x_i = 0$. One may then ask how it is that this ratio of sine-factors is recovered in the present context. Indeed, naively putting $x = 0$ in the expression for the B factors renders the u integrals trivial and we recover erroneously the factor $C_N(\{j_m\})$ without the ratio of sine-factors. However, a more careful analysis of the cut structure and of the integration contours shows that in fact this ratio arises again when care is exercised. To see this [111], it is convenient to change variable from u_N to $U_N = 1/u_N$. As a function of U_N the integrand has branch point singularities at $U_N = 0, \infty, U_N(x, w_i)$ with

$$U_N(x, w_i) = \sum_{\ell=1}^{N-1} \frac{x z_\ell/u_\ell}{(w_i - z_\ell)(x - 1)} \quad (4.172)$$

Now we should remember that all the u integrals are to be taken along small circles surrounding the origin. The U_N contour is therefore some *large* circle. However, for x close to 1, we see that $U_N(x, w_i)$ will lie *outside* this circle, which we therefore may deform along a contour running above and below the negative real axis. We should choose the cut structure from the branch points $U_N(x, w_i)$ in a way not interfering with the integration contour for U_N or the deformations of it. As x is decreased from 1, the branch points at $U_N(x, w_i)$ move closer to the original integration circle for U_N and eventually cross it. Therefore, by analytic continuation this circle has to be deformed so as to keep the singularities always outside for $x > 0$. A convenient way of doing that is precisely by deforming it to immediately wrap around the negative axis. But then the contributions from above and below the axis will have different phases depending on the power of U_N (which is $2j_N - k - 1$), and it is not difficult to see that in the limit $x = 0$ these exactly reproduce the seemingly missing ratio of sine-factors, and we are left with (4.162). Here one uses that the remaining u integrations become trivial

$$\Gamma(2j_\ell + 1) \oint \frac{du}{2\pi i} u_\ell^{2j_\ell - 1} e^{1/u_\ell} = 1 \quad (4.173)$$

Consider now [111] the function $G(w)$

$$\begin{aligned} G(w) &= G^-(w)w \\ G^-(w) &= \langle J^-(w)\mathcal{O} \rangle \\ &= \left\{ -\sum_{i,j} B(w) \frac{D_{B_i} D_{B_j}}{(w - w_i)(w - w_j)} + (t - 2) \sum_i \frac{D_{B_i}}{(w - w_i)^2} \right. \\ &\quad \left. - 2 \sum_{i,j} \frac{k_i D_{B_j}}{(w - w_i)(w - w_j)} - 2 \sum_{m,j} \frac{j_m D_{B_j}}{(w - z_m)(w - w_j)} \right\} W_B W_N^\varphi F \end{aligned} \quad (4.174)$$

The function $G(w)$ has simple poles as a function of w . It is a rather simple matter to evaluate the pole residues along the lines described when considering the Knizhnik-Zamolodchikov equations. For $w = z_m, \infty$ the result is

$$\begin{aligned} \oint_{z_m} \frac{dw}{2\pi i} G^-(w)w &= z_m \partial_{x_m} W_B F W_N^\varphi \\ \oint_{\infty} \frac{dw}{2\pi i} G^-(w)w &= \partial_{x_N} W_B F W_N^\varphi \end{aligned} \quad (4.175)$$

These contributions add up to produce the total derivative of the original correlator with respect to x

$$\sum_m z_m \partial_{x_m} \mathcal{O} + \partial_{x_N} \mathcal{O} = \partial_x \mathcal{O} \quad (4.176)$$

The pole residue at $w = w_i$ is (see (4.117))

$$\oint_{w_i} \frac{dw}{2\pi i} G^-(w)w = t \partial_{w_j} \left(w_j \frac{W_N^\varphi W_B F}{B(w_j)} \right) \quad (4.177)$$

After integration over the w_i 's we see that the expression (4.176) will vanish since this merely is the condition that the total sum of pole residues vanishes (when the pole at infinity is included as it is here). Hence, the interpolating correlator is indeed independent of x .

4.5.3 Comparison with Standard Formulations of Hamiltonian Reduction

Having proved the equivalence of the two apparently different kinds of correlators, we now want to understand this equivalence from the point of view of quantum Hamiltonian reduction. We briefly review the background. Setting the affine current

$$J^+(z) = 1 \quad (4.178)$$

in the the equation of motion derived from the $SL(2)$ WZNW theory, one recovers the classical equation of motion for Liouville theory. In order to implement the constraint (4.178) at the quantum level, one introduces a Lagrangian multiplier field $A(z)$ and follows the standard procedure for Hamiltonian reduction [24], where $A(z)$ is treated as a gauge field. The final theory, after gauge fixing, involves Faddeev-Popov ghost fields, which are supposed to cancel out unwanted degrees of freedom in the original WZNW theory. The BRST quantisation has now become a standard approach to constrained Hamiltonian systems. As far as correlation functions on the sphere are concerned, the BRST quantisation is equivalent to imposing the constraint (4.178) on the correlators. Suppose one writes the correlation function on the sphere as an operator insertion

$$\langle 0 | \hat{O} | 0 \rangle \quad (4.179)$$

then for the constrained system satisfying (4.178), we have

$$\langle 0 | \hat{O} (J^+(z) - 1) | 0 \rangle = 0 \quad (4.180)$$

(4.180) is equivalent to the following conditions

$$\begin{aligned} J_n^+ | 0 \rangle &= 0 & n \geq 0 \\ \langle 0 | \hat{O} (J_{-n}^+ - \delta_{n,1}) &= 0 & n \geq 1 \end{aligned} \quad (4.181)$$

In order not to confuse the notations used here, $J^+(z)$ is always considered to be a conformal spin 1 field to fit the WZNW theory, so that it has the expansion

$$J^+(z) = \sum_{n \in \mathbb{Z}} J_n^+ z^{-n-1} \quad (4.182)$$

As usual, to fix $J^+(z)$ to be a constant value would require $J^+(z)$ be a scalar field. In other words, the energy momentum tensor must be improved from the Sugawara construction by adding a term $\partial_z J^3(z)$. In that context, one should rename $J_n^+ \rightarrow J_{n+1}^+$.

(4.181) is called the physical state condition. In BRST quantisation the physical state space is the same as the BRST cohomology space $Ker(Q)/Im(Q)$, where Q is the BRST charge defined by

$$Q = \oint \frac{dw}{2\pi i} (J^+(w) - 1) c(w) \quad (4.183)$$

Here $c(w)$ is a conformal spin 1 fermionic ghost field with respect to the improved energy momentum tensor. Its conjugate field $b(w)$ is the anti-ghost field of spin 0 satisfying

$$b(w)c(z) = \frac{1}{w-z} \quad (4.184)$$

(4.181) is equivalent to the BRST condition, in which one requires that the vacuum states $\langle 0|$ and $|0\rangle$ be physical states, and $\hat{O}$ be a physical operator which maps physical states into physical states (see e.g. [98, 80]). In other words

$$\langle 0|Q = [Q, \hat{O}] = Q|0\rangle = 0 \quad (4.185)$$

Now consider the most general form for a class of conformal blocks in $SL(2)$ WZNW theory, which are proportional to those in the Virasoro minimal models. They can be written in the following form

$$\begin{aligned} & \langle j_{r_N, s_N} | F(J_1^-) \phi_{j_{r_{N-1}, s_{N-1}}}(z_{N-1}, 0) \dots \phi_{j_{r_2, s_2}}(z_2, 0) \prod_i \int dw_i S_{k_i}(w_i) | j_{r_1, s_1} \rangle \\ &= C \langle \Delta_{r_N, s_N+1} | \phi_{r_{N-1}, s_{N-1}+1}(z_{N-1}) \dots \phi_{r_2, s_2}(z_2) \prod_i \int dw_i V_{\alpha_i}(w_i) | \Delta_{r_1, s_1+1} \rangle \end{aligned} \quad (4.186)$$

where the normalisation constant C is found as before to be

$$C = (-1)^{2j_N - \sum_{i=1}^N j_i} \frac{\Gamma(\sum_{i=1}^N j_i - k)}{\Gamma(2j_N - k)} \partial_y^{-2j_N + \sum_{i=1}^N j_i} F(y)|_{y=0} \quad (4.187)$$

where we used that

$$-2j_N + \sum_i j_i = -\sum_i k_i \quad (4.188)$$

In general C depends on t and the j_i 's. For some values of t and j_i 's, C vanishes. Then the conformal blocks in the Virasoro minimal models can only be obtained by dividing out C . In other cases C becomes infinity. Then the conformal blocks for the Virasoro minimal models can be either finite or zero, and in the latter case the relation between the WZNW and the minimal conformal blocks is singular. Strictly speaking, simply taking the limit $x_i \rightarrow z_i$ is not equivalent to quantum Hamiltonian reduction (4.178). Rather it is in accord with the constraint

$$J^+(w) = J_{-1}^+ \quad (4.189)$$

To go to the minimal model we must further impose the condition

$$J_{-1}^+ = 1 \quad (4.190)$$

To see this, let us consider the BRST charge for quantum Hamiltonian reduction (4.189)

$$\tilde{Q} = \oint \frac{dw}{2\pi i} (J^+(w) - J_{-1}^+) c(w) \quad (4.191)$$

The physical state space now becomes the BRST cohomology space $Ker(\tilde{Q})/Im(\tilde{Q})$. It is clear that $\phi_j(z, 0)$ commutes with $\tilde{Q}$, hence maps a physical state into another physical state. Now consider the ket and the bra states. Notice that the ket state $|j_1\rangle$ is a highest weight state and the bra state $\langle j_{r_N, s_N}|$ is a lowest weight state

$$J_n^+ |j_1\rangle = \langle j_{r_N, s_N} | J_{-n-1}^+ = 0, \quad n \geq 0 \quad (4.192)$$

For the b, c ghost fields, we have the following condition

$$c_n |j_1\rangle = b_{n+1} |j_1\rangle = \langle j_{r_N, s_N} | c_{-n-1} = \langle j_{r_N, s_N} | b_{-n} = 0, \quad n \geq 0 \quad (4.193)$$

It can be verified that with respect to the BRST charge $\tilde{Q}$ in (4.191), $|j_1\rangle$ is a physical state, and the bra state $\langle j_{r_N, s_N}|F(J_1^-)$ is a physical state for any arbitrary function $F(J_1^-)$. However, this extra degree of freedom is removed if we further impose the condition (4.190), which would fix the function $F(J_1^-)$ uniquely

$$\langle j_N|F(J_1^-)J_{-1}^+ = \langle j_N|F(J_1^-) \quad , \quad [e^{-\gamma_1}, \beta_{-1}] = e^{-\gamma_1} \quad (4.194)$$

and we recover exactly the conformal blocks in the Virasoro minimal models (here $\alpha = r - st$)

$$\begin{aligned} \langle j_{r_N, s_N}|F(J_1^-) &= \langle j_{r_N, s_N}|e^{-\gamma_1} \\ &= \langle j_{r_N, s_N}| \sum_{n \in \mathbb{Z}} \frac{\Gamma(2j_N - k)}{\Gamma(2j_N - k + n + \alpha)\Gamma(n + \alpha + 1)} (-J_1^-)^{n+\alpha} \end{aligned} \quad (4.195)$$

(here corrected for minor misprint, compared to [111]) where γ_n is conjugate to J_{-n}^+

$$[J_{-n}^+, \gamma_m] = \delta_{n,m} \quad (4.196)$$

If we were to use $\phi_j(z, z)$ to represent a primary field in the Hamiltonian reduced system (strictly speaking, $\phi_j(z, z)$ does not transform as a primary field for the Virasoro algebra in the reduced system), then we should normalise the correlation function by dividing out the normalisation constant C . Then, in the limit C goes to zero, the conformal block in the reduced system would remain finite.

In conclusion, the constraint $J^+(z) = 1$ completely freezes the degrees of freedom of the $J^+(z)$ field. However, we could proceed in two steps in putting the constraint on the correlation functions. First set $J^+(z) = J_{-1}^+$ and then let $J_{-1}^+ = 1$. The first step would result in a class of correlation functions which are proportional to that of the completely constrained system, like the ones considered in the previous sections. However, the remaining degrees of freedom of the J_{-1}^+ mode is reflected by the arbitrariness of the proportionality. If we normalise the correlation function by dividing out the normalisation constant, which is equivalent to setting $J_{-1}^+ = 1$, then we recover the corresponding correlators in the completely reduced system.

Chapter 5

4-point Greens Functions in $SL(2)$ Current Algebra

Once the chiral blocks are obtained the next step in the program is to determine the monodromy invariant Greens functions. These are the ones for which physical applications can be made and they are necessary before for example an application to $2D$ quantum gravity can be made. It is the principal goal in this chapter to obtain these monodromy invariant combinations, and from the ensuing monodromy coefficients to determine the operator algebra coefficients. This is then a generalisation of the celebrated work [42] by Dotsenko and Fateev in conformal minimal models to CFT based on affine $SL(2)$ current algebra, and may be viewed as the completion of the solution of CFT (on the sphere) based on affine $SL(2)$ current algebra for admissible representations.

The problem of determining the Greens functions is conveniently solved by means of the crossing matrix relating the conformal blocks in the s- and t-channels (in fact, just a particular row and column of that matrix). The conformal blocks for 4-point functions may be characterised in terms of couplings to intermediate states. These in turn are determined by the fusion rules of the theory, discussed in the preceding chapter. We shall find that both sets of fusion rules are operating. Thus, the fusion rules provide a neat starting point for giving convenient bases for the conformal blocks in the s- and t-channels. One must then understand how the general integral representations can reproduce these bases. Here we shall use either the ones provided in Chapter 4 (sometimes to be referred to as PRY), or the one for the 4-point function obtained by Andreev [8], and in both cases specify how integration contours must be chosen in order to generate specific members of s- or t-channel bases. In [8] the contours are not specified. In the case of the PRY integral representation, we show how to obtain chiral blocks in the s-channel corresponding to fusion rule I, and how to obtain blocks in the t-channel corresponding to fusion rule II, using contours where the integration of the auxiliary variable u is carried out first. We also show how to obtain blocks in the s-channel corresponding to fusion rule II, and how to obtain blocks in the t-channel corresponding to fusion rule I, using contours where the u integration is done last. In the integral representation of Andreev there is no u variable to worry about and the contours we find are more tractable. That his 4-point blocks are equivalent to ours is a priori rather clear since both he and we have checked that the blocks we write down satisfy the Knizhnik-Zamolodchikov equations. Nevertheless, we find it very instructive to attempt a direct analytic proof of how the equivalence may be obtained.

Substantial evidence will be provided at the level of highly non-trivial integral manipulations.

Having written down the full s- and t-channel bases for conformal blocks and understood the corresponding integration contours, we may go on to calculate the relevant parts of the crossing matrix. Certain ratios of matrix elements determine the monodromy coefficients used in building the monodromy invariant 4-point Greens function.

In the first considerations it will be assumed that both vertices in the 4-point blocks pertain to the same fusion rule (I *or* II). Then the idea of over-screening used for the 3-point function may be employed to obtain additional 4-point blocks in which there are different fusion rules (I *and* II) operating at the two vertices of the block. These new 4-point blocks correspond to different sets of external spins from the ones previously considered, and so they do not intertwine with these under crossing. Based on the calculations in the case with non-mixed fusion rules, it is fairly easy to obtain the new monodromy coefficients. We may finally write down the operator algebra coefficients and in particular, utilising the idea of over-screening allows us to normalise unambiguously the operator algebra coefficients for fusion rule II, contrary to the results in [8]. It turns out that in terms of appropriate parametrisations, the operator algebra coefficients are identical for the two fusion rules up to a power of the parameter t , which is determined as a consistency condition.

5.1 4-point Functions

In the 4-point function

$$\langle \phi_{j_4}(z_4, x_4) \phi_{j_3}(z_3, x_3) \phi_{j_2}(z_2, x_2) \phi_{j_1}(z_1, x_1) \rangle \quad (5.1)$$

we consider as usual the limits

$$\begin{aligned} z_4 &\rightarrow \infty, & x_4 &\rightarrow \infty \\ z_3 &\rightarrow 1, & x_3 &\rightarrow 1 \\ z_2 &\rightarrow z, & x_2 &\rightarrow x \\ z_1 &\rightarrow 0, & x_1 &\rightarrow 0 \end{aligned} \quad (5.2)$$

so that the 4-point conformal blocks will be (in general multi-valued) functions of (z, x) . We label s- and t-channel conformal blocks by tree graphs, the meaning of which is that in the limit $z \rightarrow 0$ followed by $x \rightarrow 0$ the s-channel block corresponding to Fig. 1 has the behaviour following from the OPE's

$$S(z, x) \sim z^{\Delta - \Delta_1 - \Delta_2} (-x)^{j_1 + j_2 - j} (1 + \mathcal{O}(z, -x)) \quad (5.3)$$

whereas for the t-channel block we have in the limit $z \rightarrow 1$ followed by $x \rightarrow 1$

$$T(z, x) \sim (1 - z)^{\Delta - \Delta_2 - \Delta_3} (x - 1)^{j_2 + j_3 - j} (1 + \mathcal{O}(1 - z, x - 1)) \quad (5.4)$$

Here the conformal weights are given by the standard expression $\Delta_i = j_i(j_i + 1)/t$.

A very convenient way to think about the fusion rules consists in the following. Consider the s-channel coupling of j_1, j_2 to a j . When we parametrise $j = j_I$ for fusion rule I as $j_1 + j_2 - j_I = r - st$, the integers r, s are related to the number of screenings of the first and




Figure 5.1: Graphs for s- and t-channel blocks

second kinds around the $j_1 j_2 j_I$ vertex. The singular behaviour of the s-channel block in the limit $z \rightarrow 0, x \rightarrow 0$ is then

$$z^{\Delta-\Delta_1-\Delta_2}(-x)^{j_1+j_2-j_I} = z^{\Delta-\Delta_1-\Delta_2}(-x)^{r-st} \quad (5.5)$$

with $\Delta = j_I(j_I + 1)/t$. For fusion rule II we may then parametrise the internal j as

$$j_{II} = -j_I - 1 \quad (5.6)$$

Of course the conformal dimensions for j_I and j_{II} are the same, but we find the singular behaviour of the s-channel block to be

$$z^{\Delta-\Delta_1-\Delta_2}(-x)^{j_1+j_2-j_{II}} = z^{\Delta-\Delta_1-\Delta_2}(-x)^{2j_1+2j_2-r+st+1} \quad (5.7)$$

All these statements follow by analysing the fusion rules (4.56) and (4.57). By analysing the s-channel 4-point blocks in the limit $z \rightarrow 0, x \rightarrow 0$ we indeed find both of these singular behaviours (see below) and hence verify that the blocks realize both fusion rules I and II. In the t-channel the discussion is analogous, with $j_1 \leftrightarrow j_3$, $z \rightarrow 1 - z$ and $x \rightarrow 1 - x$, so that we consider the limits $z \rightarrow 1$ followed by $x \rightarrow 1$.

Before specifying the contours needed to produce chiral blocks with these singular behaviours, we reconsider the general result for N -point functions in the case of $N = 4$ in order to simplify the integral expression. First of all we want to reduce the number of auxiliary integration variables u from two to one. In due course, let us reconsider the β, γ part of the 4-point function

$$\langle j_4 | [\phi_{j_3}(z_3, x_3)]_j^{j_4} [\phi_{j_2}(z_2, x_2)]_{j_1}^j | j_1 \rangle_{\beta\gamma} \quad (5.8)$$

with the notation

$$\begin{aligned} j_1 + j_2 - j &= \rho_2 - \sigma_2 t \\ j + j_3 - j_4 &= \rho_3 - \sigma_3 t \end{aligned} \quad (5.9)$$

Thus we get

$$W_4 \sim \Gamma(2j_3 + 1)\Gamma(2j_2 + 1) \oint_0 \frac{du_2}{2\pi i} \frac{du_3}{2\pi i} \frac{1}{u_2 u_3}$$

$$\begin{aligned}
& \cdot \prod_{i_3=1}^{\rho_3} \left[\frac{x_2/u}{w(3, i_3) - z_2} + \frac{x_3}{w(3, i_3) - z_3} \right] \prod_{l_3=1}^{\sigma_3} \left[\frac{x_2/u}{v(3, l_3) - z_2} + \frac{x_3}{v(3, l_3) - z_3} \right]^{-t} \\
& \cdot \prod_{i_2=1}^{\rho_2} \left[\frac{x_2/u}{w(2, i_2) - z_2} + \frac{x_3}{w(2, i_2) - z_3} \right] \prod_{l_2=1}^{\sigma_2} \left[\frac{x_2/u}{v(2, l_2) - z_2} + \frac{x_3}{v(2, l_2) - z_3} \right]^{-t} \\
& \cdot u^{\rho_2 - \sigma_2 t} (u_3^{-1} D_3)^{-2j_3 + \rho_3 - \sigma_3 t} \exp(1/u_3) (u_2^{-1} D_2)^{-2j_2 + \rho_2 - \sigma_2 t} \exp(1/u_2)
\end{aligned} \tag{5.10}$$

where we have introduced

$$u = u_2/u_3 \tag{5.11}$$

Again the (somewhat misleading) notation is that the D 's are derivatives with respect to the entire argument of the relevant exponentials.

We want to consider a change of variables from (u_2, u_3) to (u_2, u) . The integration measure is

$$\frac{du_2 du_3}{u_2 u_3} = -\frac{du_2 du}{u_2 u} \tag{5.12}$$

and

$$2j_2 + 2j_3 - \rho_3 + \sigma_3 t - \rho_2 + \sigma_2 t = 2j_2 + 2j_3 - (j_3 + j - j_4) - (j_2 + j_1 - j) = j_2 + j_3 + j_4 - j_1 = J_1 \tag{5.13}$$

Here we have introduced the notation

$$J_i = j_1 + j_2 + j_3 + j_4 - 2j_i \tag{5.14}$$

Using the generalised exponential identity (2.120), we obtain the following u_2 dependence

$$\frac{du du_2}{u u_2} u^{-2j_3 + \rho_2 - \sigma_2 t + \rho_3 - \sigma_3 t} u_2^{J_1} D^{-J_1} \exp\left(\frac{1+u}{u_2}\right) \tag{5.15}$$

Now the integral over u_2 will produce the factor

$$\frac{(1+u)^{J_1}}{\Gamma(J_1 + 1)} \tag{5.16}$$

We are left with the following integral

$$\begin{aligned}
& - \int \frac{du}{2\pi i} \prod_{i_3=1}^{\rho_3} \left[\frac{x_2}{w(3, i_3) - z_2} + \frac{x_3 u}{w(3, i_3) - z_3} \right] \prod_{l_3=1}^{\sigma_3} \left[\frac{x_2}{v(3, l_3) - z_2} + \frac{x_3 u}{v(3, l_3) - z_3} \right]^{-t} \\
& \cdot \prod_{i_2=1}^{\rho_2} \left[\frac{x_2}{w(2, i_2) - z_2} + \frac{x_3 u}{w(2, i_2) - z_3} \right] \prod_{l_2=1}^{\sigma_2} \left[\frac{x_2}{v(2, l_2) - z_2} + \frac{x_3 u}{v(2, l_2) - z_3} \right]^{-t} \\
& \cdot u^{-2j_3-1} \frac{(1+u)^{J_1}}{\Gamma(J_1 + 1)} \Gamma(2j_3 + 1) \Gamma(2j_2 + 1)
\end{aligned} \tag{5.17}$$

To write the final result for the 4-point function in a more compact form, let us collectively denote the positions for both kinds of screening charges as

$$w_i, \quad i = 1, \dots, \rho_2 + \rho_3 + \sigma_2 + \sigma_3 \equiv M \tag{5.18}$$

Then the complete expression for the 4-point function is

$$\begin{aligned}
& \langle j_4 | [\phi_{j_3}(z_3, x_3)]_j^{j_4} [\phi_{j_2}(z_2, x_2)]_{j_1}^j | j_1 \rangle \\
&= - \prod_{m < n}^3 (z_m - z_n)^{2j_m j_n / t} \Gamma(2j_3 + 1) \Gamma(2j_2 + 1) \\
&\cdot \int \frac{du}{2\pi i} \prod_{i=1}^M \frac{dw_i}{2\pi i} \left(\frac{x_2}{w_i - z_2} + \frac{x_3 u}{w_i - z_3} \right)^{-k_i} \prod_{i < j} (w_i - w_j)^{\frac{2k_i k_j}{t}} \\
&\cdot \prod_{i=1}^M \prod_{l=1}^3 (w_i - z_l)^{\frac{2k_i j_l}{t}} u^{-2j_3-1} \frac{(1+u)^{J_1}}{\Gamma(J_1+1)} \tag{5.19}
\end{aligned}$$

where only *one* auxiliary u integration is involved and we have included the φ part. $k_i = -1$ for screenings of the first kind and $k_i = t$ for screenings of the second kind. In the limit (5.2) we obtain, letting $u \rightarrow -u$ and up to normalisation

$$\begin{aligned}
W_4 &= z^{2j_1 j_2 / t} (1-z)^{2j_2 j_3 / t} \int \frac{du}{2\pi i} \prod_{i=1}^M \frac{dw_i}{2\pi i} w_i^{2k_i j_1 / t} (w_i - z)^{2k_i j_2 / t} (w_i - 1)^{2k_i j_3 / t} \\
&\cdot \prod_{i < j} (w_i - w_j)^{2k_i k_j / t} \left(-\frac{u}{w_i - 1} + \frac{x}{w_i - z} \right)^{-k_i} u^{-2j_3-1} (1-u)^{J_1} \tag{5.20}
\end{aligned}$$

5.1.1 Integration Contours

In this section we shall specify the integration contours [112] for the various variables in the integral representation (5.20) of the 4-point function. The first contours we indicate will give rise to a set of s-channel conformal blocks corresponding to the intermediate state j in Fig. 1 being given by fusion rule I, and to a set of conformal blocks in the t-channel corresponding to the intermediate j being given by fusion rule II. In these cases the u integration is carried out first.

We first describe the situation in the s-channel corresponding to fusion rule I. We write for the conformal block

$$\begin{aligned}
W_{(r,s)}^{(R,S)}(j_1, j_2, j_3, j_4; z, x) &= z^{2j_1 j_2 / t} (1-z)^{2j_2 j_3 / t} \oint_{\mathcal{C}_{\mathcal{I}}} \prod_{i \in \mathcal{I}} \frac{dw_i}{2\pi i} \oint_{\mathcal{C}_{\mathcal{O}}} \prod_{j \in \mathcal{O}} \frac{dw_j}{2\pi i} \oint_{\mathcal{C}_u} \frac{du}{2\pi i} \\
&\cdot w_i^{2k_i j_1 / t} (w_i - z)^{2k_i j_2 / t} (w_i - 1)^{2k_i j_3 / t} \prod_{\substack{i, j \in \mathcal{A} \\ i < j}} (w_i - w_j)^{2k_i k_j / t} \\
&\cdot \prod_{i \in \mathcal{A}} \left(-\frac{u}{w_i - 1} + \frac{x}{w_i - z} \right)^{-k_i} u^{-2j_3-1} (1-u)^{2j_2+2j_3-R+St} \tag{5.21}
\end{aligned}$$

Here we are considering an integral representation of the 4-point conformal block with a total of R screening operators of the first kind and a total of S screening operators of the second kind. The w_i 's are the positions of the screening operators. When $i \in \mathcal{O}$ the corresponding w_i is integrated along the contour of Fig. 5.2(a) (whether it is of the first or second kind) corresponding to a screening of the vertex $j_1 j_2 j$. Different $i, i' \in \mathcal{O}$ are taken along slightly different contours in order to avoid the singularity coming from $(w_i - w_{i'})^{2k_i k_{i'} / t}$. Similarly, the




Figure 5.2: The integration contours $\mathcal{C}_\mathcal{O}$ (a) and $\mathcal{C}_\mathcal{I}$ (b) for an s-channel block corresponding to fusion rule I.

w_j 's for $j \in \mathcal{I}$ are integrated along the contour Fig. 5.2(b) corresponding to a screening of the vertex jj_3j_4 . $\mathcal{A} = \mathcal{O} \cup \mathcal{I}$ is simply the combined index set. We denote the numbers of screenings of the first kind at the j_1j_2j and the jj_3j_4 vertices respectively as r and $R - r$. Similarly the corresponding numbers of screenings of the second kinds at the two vertices are denoted s and $S - s$. In the product of factors $(w_i - w_j)^{2k_ik_j/t}$ an arbitrary ordering of the indices is implied.

For fixed w_i 's the integrand has singularities in the u plane at $u = 0, 1, \Delta_i$, where

$$\Delta_i = \frac{w_i - 1}{w_i - z}x \quad (5.22)$$

The integration contour for u is to divide the singularities Δ_i , so that the ones for $i \in \mathcal{O}$ lie outside $\mathcal{C}_u$ and the ones for $i \in \mathcal{I}$ lie inside $\mathcal{C}_u$, and $\mathcal{C}_u$ should pass through $u = 1$. For z and x sufficiently small, we may take $\mathcal{C}_u$ to be the unit circle, Fig. 5.3. Remember that in order to identify the nature of the block and the value of the intermediate j , we are going to investigate the limit $z \rightarrow 0$ followed by $x \rightarrow 0$. The different positions of the singularities Δ_i in u for $i \in \mathcal{O}$ and $i \in \mathcal{I}$ mean that they give rise to different singularities in the corresponding w_i planes after the u integration has been performed. In fact, for $i \in \mathcal{O}$ there occurs a pinching of singularities when Δ_i collides with either 0 or 1 (the additional singularities in u). This happens for w_i equal to 1, and for w_i equal to

$$\delta = \frac{x - z}{x - 1} \quad (5.23)$$

respectively. In particular no extra singularity is generated at $w_i = z$. This is why we may take the contour in w_i to start from z as indicated, since the singularity for w_i is what we term 'pure', meaning that it is of the form

$$(w_i - z)^a(1 + \mathcal{O}(w - z)) \quad (5.24)$$




Figure 5.3: The integration contour $\mathcal{C}_u$ for an s-channel block corresponding to fusion rule I.

One can check that this is enough to ensure that the corresponding block will satisfy the KZ equations, going over the proof presented in chapter 4 [110]. In contrast, the singularity at $w_i = 0$ is 'non-pure': it is a mixture of different powers of w_i . Hence we cannot allow the contour to end in $w_i = 0$, it has to surround that point as indicated.

Turning to the singularities in w_i for $i \in \mathcal{I}$, we see that pinching occurs only when $\Delta_i = 1$, so that there is no extra singularity produced at $w_i = 1$: it remains pure, and we may take the integration contour to start in $w_i = 1$ as indicated. If more convenient, one may take the contour to wrap around the real axis from 1 to ∞ , which is a form closer to the one used by Dotsenko and Fateev [42].

Having established that the choice of contours indicated is allowed in the sense that the conformal block will satisfy the KZ equations, it is a relatively simple matter to find the leading singularity in the limit $z \rightarrow 0$ followed by $x \rightarrow 0$. In fact, we may scale all the w_i 's with $i \in \mathcal{O}$ as

$$w_i \rightarrow zw_i \quad (5.25)$$

In the limit $z \rightarrow 0$ this is easily seen to result in a leading z behaviour of the form

$$W_{(r,s)}^{(R,S)}(z, x) \sim z^{-\Delta(j_1) - \Delta(j_2) + \Delta(j_I)} \quad (5.26)$$

where $\Delta_j = \Delta(j) = j(j+1)/t$ and where

$$j_I = j_1 + j_2 - r + st \quad (5.27)$$

This is not enough to prove that indeed the intermediate state corresponds to a primary field with that value of j , since

$$\Delta(j) = \Delta(-j-1) \quad (5.28)$$

In fact, according to our earlier discussion, the difference between fusion rules I and II is exactly that for fusion rule I we should obtain $j = j_I$ whereas for fusion rule II we should




Figure 5.4: The integration contour $\mathcal{C}_u$ for a t-channel block corresponding to fusion rule II.



Figure 5.5: The integration contours for the w_i 's: $\mathcal{C}_O$ (a) and $\mathcal{C}_I$ (b) for a t-channel block corresponding to fusion rule II.

obtain $j = j_{II} = -j_I - 1$. In other words the z behaviour is precisely unable to distinguish between the two fusion rules. To distinguish we must investigate the leading x behaviour in the limit $x \rightarrow 0$ after we have taken $z \rightarrow 0$. However, it is an easy matter to do so and we find

$$W_{(r,s)}^{(R,S)} \sim z^{-\Delta(j_1) - \Delta(j_2) + \Delta(j_I)} (-x)^{r-st} \quad (5.29)$$

This is the proof that the conformal block we have constructed corresponds to fusion rule I, since $r - st = j_1 + j_2 - j_I$.

We next describe how contours have to be chosen in order to produce a t-channel block corresponding to fusion rule II. This situation can occur only provided there is at least one screening operator of the second kind. We use the same defining equation as in (5.21), but the sets of indices as well as r and s have different meanings. Again we have a total of R and S screenings of the first and second kinds. There are r and s screenings associated with the upper vertex, and the corresponding index set for the w_i 's is $\mathcal{O}$. There are $R - r$ and $S - s$ screening operators of the two kinds associated with the lower vertex and the corresponding index set for the w_i 's is $\mathcal{I}$. The integration contour $\mathcal{C}_u$ is indicated in Fig. 5.4. One checks that in the limit $z \rightarrow 1$ followed by $x \rightarrow 1$ the two sets of singularities, Δ_i for $i \in \mathcal{O}$ and $i \in \mathcal{I}$ respectively, are well separated, so that the contour may be taken to separate them as indicated. The contours $\mathcal{C}_O$ and $\mathcal{C}_I$ for the two sets of w_i variables, are shown in Fig. 5.5 (a) and (b). In all cases one checks as for the s-channel block that the nature of singularities is such that the contours may

be chosen as indicated, and that the block thus defined will satisfy the KZ equations. Then we investigate the combined behaviour $z \rightarrow 1$ followed by $x \rightarrow 1$. To this end we perform the following scalings of the integration variables

$$\begin{aligned} w_i &\rightarrow \frac{w_i - 1}{z - 1}, \quad i \in \mathcal{O} \\ w_i &\rightarrow \frac{w_i}{w_i - 1}, \quad i \in \mathcal{I} \\ u &\rightarrow \frac{u - \Delta_{j_0}}{1 - \Delta_{j_0}} \end{aligned} \quad (5.30)$$

where j_0 is an arbitrary index in the set $\mathcal{I}$, however with the restriction that w_{j_0} is the position of a screening operator of the second kind. It is rather straightforward to check that this gives rise to the combined singular behaviour

$$W_{(r,s)}^{(R,S)}(z, x) \sim (1 - z)^{-\Delta(j_2) - \Delta(j_3) + \Delta(j_{II})} (x - 1)^{2j_2 + 2j_3 - r + st + 1} \quad (5.31)$$

where

$$\begin{aligned} j_{II} &= -j_I - 1 \\ j_I &= j_2 + j_3 - r + st \end{aligned} \quad (5.32)$$

so that (5.31) exactly demonstrates that we have fusion rule II, since $j_2 + j_3 - j_{II} = 2j_2 + 2j_3 - r + st + 1$.

It follows that the conformal blocks defined on the basis of the free field realization elaborated in [110, 112], indeed do give rise to both the fusion rules.

In [112] also realizations of conformal blocks corresponding to fusion rule II in the s-channel and of ones corresponding to fusion rule I in the t-channel, have been found. However, these are not given by quite as simple contours as above. It is the appearance of non-pure singularities which prevents one from finding such simple contours. The new idea is to carry out first the integrations of the screening operators letting the contours depend on u . Then there are only pure singularities in the w_i planes and there will be no problems caused by non-pure singularities. It turns out that it is possible to find contours like that leaving, upon integration, a simple u integral. Let us first consider the conformal blocks in the s-channel corresponding to fusion rule II, where the contours are depicted in Fig. 5.6

$$\begin{aligned} &W_{(r,s)}^{(R,S)}(j_1, j_2, j_3, j_4; z, x) \\ &= z^{2j_1 j_2 / t} (1 - z)^{2j_2 j_3 / t} \oint_{\mathcal{C}_u} \frac{du}{2\pi i} \oint_{\mathcal{C}_w} \frac{dw}{2\pi i} \int_1^\infty \prod_{i \in \mathcal{I}} dw_i \int_0^z \prod_{i \in \mathcal{O}} dw_i \\ &\cdot w_i^{2k_i j_1 / t} (w_i - z)^{2k_i j_2 / t} (w_i - 1)^{2k_i j_3 / t} \prod_{\substack{i, j \in \mathcal{A} \\ i < j}} (w_i - w_j)^{2k_i k_j / t} \\ &\cdot \prod_{i \in \mathcal{A}} \left(-\frac{u}{w_i - 1} + \frac{x}{w_i - z} \right)^{-k_i} (1 - u)^{2j_2 + 2j_3 - R + St} u^{-2j_3 - 1} \end{aligned} \quad (5.33)$$

where

$$\begin{aligned} \mathcal{A} &= \mathcal{I} \cup \mathcal{O} \cup \{R + S\} \\ w &= w_{R+S} \\ k_{R+S} &= t \end{aligned} \quad (5.34)$$




Figure 5.6: Integration contours for an s-channel block corresponding to fusion rule II. The corresponding u integration is simply along the unit circle.

To see that the above formula produces the right singular behaviour in the limit $z \rightarrow 0$ followed by $x \rightarrow 0$, we may scale all the w_j 's with $j \in \mathcal{O}$ as

$$w_j \rightarrow zw_j, \quad \text{for all } j \in \mathcal{O} \quad (5.35)$$

and scale all the w_i 's with $i \in \mathcal{I}$ as

$$w_i \rightarrow 1/w_i, \quad \text{for all } i \in \mathcal{I} \quad (5.36)$$

and also

$$w \rightarrow \frac{uz-x}{u-x}w \quad (5.37)$$

One can then show that

$$W_{(r,s)}^{(R,S)}(z, x) \sim z^{-\Delta(j_1)-\Delta(j_2)+\Delta(j_{II})}(-x)^{j_1+j_2-j_{II}} \quad (5.38)$$

where

$$j_{II} = -j_1 - j_2 + r - st - 1 \quad (5.39)$$

This is precisely the expected singular behaviour.

Finally we consider the conformal blocks in the t-channel corresponding to fusion rule I, see Fig. 5.7

$$\begin{aligned} & W_{(r,s)}^{(R,S)}(j_1, j_2, j_3, j_4; z, x) \\ &= z^{2j_1j_2/t}(1-z)^{2j_2j_3/t} \oint_{\mathcal{C}_u} \frac{du}{2\pi i} \oint_{\mathcal{C}_w} \frac{dw}{2\pi i} \int_0^{-\infty} \prod_{i \in \mathcal{I}} \frac{dw_i}{2\pi i} \int_z^1 \prod_{i \in \mathcal{O}} \frac{dw_i}{2\pi i} \\ & \cdot w_i^{2k_i j_1/t} (w_i - z)^{2k_i j_2/t} (1 - w_i)^{2k_i j_3/t} \prod_{\substack{i,j \in \mathcal{A} \\ i < j}} (w_i - w_j)^{2k_i k_j/t} \\ & \cdot \prod_{i \in \mathcal{A}} \left(-\frac{u}{w_i - 1} + \frac{x}{w_i - z} \right)^{-k_i} (1-u)^{2j_2+2j_3-R+St} u^{-2j_3-1} \end{aligned} \quad (5.40)$$




Figure 5.7: Integration contours for the screening charges in the case of a t-channel block corresponding to fusion rule I. The u integration is along a closed contour starting in 1 and surrounding x .

where

$$\begin{aligned}\mathcal{A} &= \mathcal{I} \cup \mathcal{O} \cup \{r+s\} \\ w &= w_{r+s} \\ k_{r+s} &= t\end{aligned}\tag{5.41}$$

To see that the above formula produces the right singular behaviour in the limit $z \rightarrow 1$ followed by $x \rightarrow 1$, we may scale all the w_j 's with $j \in \mathcal{O}$ as

$$w_j \rightarrow 1 - (1-z)w_j, \quad \text{for all } j \in \mathcal{O}\tag{5.42}$$

and scale all w_i 's with $i \in \mathcal{I}$ as

$$w_i \rightarrow w_i/(w_i - 1), \quad \text{for all } i \in \mathcal{I}\tag{5.43}$$

and also

$$w \rightarrow 1 - \frac{(1-z)}{u-x}uw\tag{5.44}$$

We also scale u as

$$u \rightarrow x + (1-x)u\tag{5.45}$$

It should be noticed that the final u contour starts at 1 and goes along the unit circle such that it surrounds 0 and the other points which are away from 0 by a distance of order $(1-z)/(1-x)$. This means that we can not deform the u contour to the form $\int_0^1 du$, or in terms of the original u variable, that cannot be deformed into $\int_x^1 du$. Using these scalings, we show that in the presence of at least one screening charge of the second kind in the scaling region (the region close to 1 and z) the singular behaviour is

$$W_{(r,s)}^{(R,S)}(z, x) \sim (1-z)^{-\Delta(j_2)-\Delta(j_3)+\Delta(j_I)}(x-1)^{j_2+j_3-j_I}\tag{5.46}$$

where

$$j_I = j_2 + j_3 - r + st\tag{5.47}$$

What happens if there is no screening charge of the second kind in the scaling region? Then the above method does not apply, but in that case $j_2 + j_3 - j_I$ is an integer, and

$$W_{(r,0)}^{(R,S)}(z,x) \sim (1-z)^{-\Delta(j_2)-\Delta(j_3)+\Delta(j_I)}(x-1)^{j_2+j_3-j_I} \quad (5.48)$$

is a polynomial in x . There will be no extra singularities present in w 's, such as at $\delta = \frac{x-z}{x-1}$, if we integrate over u first. Thus we could choose the following contours ($w = w_{r+s} = w_r \in \mathcal{O}$)

$$\begin{aligned} \int_z^1 dw_j & \quad , \quad j \in \mathcal{O} \\ \int_{-\infty}^0 dw_i & \quad , \quad i \in \mathcal{I} \\ \int_0^1 du & \end{aligned} \quad (5.49)$$

These contours are effectively closed in the sense that a total derivative integrated along them vanishes, such as is required for the KZ equations to be satisfied. Notice, however, that these contours are not closed (in the same sense) when there is a screening charge of the second kind in the scaling region. However, it is difficult to determine explicitly the $(1-x)$ behaviour for these contours, but since we know that our formula is both projective and $SL(2)$ invariant, we could express the above formula in terms of $x_3 = 0$ and $x_1 = 1$, where the $(1-x)$ behaviour is manifest.

It may seem surprising that one could not make use of the $j_1 \leftrightarrow j_3$ symmetry to obtain t-channel contours from s-channel ones and vice versa. The reason is that the simple form of the 4-point function we have given with only one auxiliary u integration, breaks this symmetry, since not all 4 primary fields are treated on the same footing. For a more symmetric treatment, more u integrations have to be introduced, which is also inconvenient, however.

5.2 Andreev's Representation

In this section we base our discussion on the integral realization of Andreev [8]. In a subsequent section we discuss the equivalence between that realization and the one described in the preceding section [110, 112] and in the sequel denoted PRY. Here we show how to choose simple integration contours [112] so that we produce both s- and t-channel blocks corresponding to both fusion rules I and II. It will turn out that the t-channel blocks are obtained in a very simple way from the s-channel blocks so we mostly concentrate on the latter. It is the specification of the integration contours which is the contribution here and in [113] over [8]. The advantage of the realization of [8] is that contrary to the case discussed so far, there is no auxiliary integration in addition to the integration over positions of screening charges. The disadvantage is that the Andreev representation (so far) has no underlying free field realization and is therefore only known for 4-point blocks. For later purposes it is convenient to have different names for s- and t-channel blocks. We denote them by letters $\mathcal{S}$ or S and $\mathcal{T}$ or T . The differences will be explained.

5.2.1 Case of Fusion Rule I

We define the complex block in the s-channel for fusion rule I with r screenings of the first kind and s screenings of the second kind at the right vertex as follows

$$\begin{aligned}
\mathcal{S}_{(r,s,0)}^{(R,S)}(z, x) = & z^{2j_1j_2/t}(1-z)^{2j_2j_3/t} \int_0^z \prod_{i \in I_1, k \in J_1} du_i dv_k \int_1^\infty \prod_{j \in I_2, l \in J_2} du_j dv_l \\
& \cdot u_i^{a'}(1-u_i)^{b'}(z-u_i)^{c'} \prod_{i < i', \in I_1} (u_i - u_{i'})^{2\rho'} u_j^{a'}(u_j-1)^{b'}(u_j-z)^{c'} \\
& \cdot \prod_{j < j', \in I_2} (u_j - u_{j'})^{2\rho'} \prod_{i \in I_1, j \in I_2} (u_j - u_i)^{2\rho'} \\
& \cdot v_k^a(1-v_k)^b(z-v_k)^c \prod_{k < k', \in J_1} (v_k - v_{k'})^{2\rho} v_l^a(v_l-1)^b(v_l-z)^c \\
& \cdot \prod_{l < l', \in J_2} (v_l - v_{l'})^{2\rho} \prod_{k \in J_1, l \in J_2} (v_l - v_k)^{2\rho} \\
& \cdot \prod_{i,k} (u_i - v_k)^{-2} \prod_{i,l} (u_i - v_l)^{-2} \prod_{j,k} (u_j - v_k)^{-2} \prod_{j,l} (u_j - v_l)^{-2} \\
& \cdot \prod_{i,j,k,l} (u_i - x)(u_j - x)(v_k - x)^{-\rho}(v_l - x)^{-\rho}
\end{aligned} \tag{5.50}$$

Here we have introduced the following index sets

$$\begin{aligned}
I_1 &= \{1, \dots, r\} \\
I_2 &= \{r+1, \dots, R\} \\
J_1 &= \{1, \dots, s\} \\
J_2 &= \{s+1, \dots, S\}
\end{aligned} \tag{5.51}$$

where R and S are the total numbers of screenings of the first and second kinds respectively. Variables u and v belong to screenings of the first and second kind respectively, although this language is rather symbolic, since as yet there exists no known free field realization which directly gives this form. Also the integrals are taken along complex Dotsenko-Fateev contours shown in Fig. 5.8. Notice that expressions of the form $(u_i - u_{i'})^{2\rho'}$ have a phase defined by the fact that the first of the two integration variables have a lower imaginary part than the last variable. Finally

$$\begin{aligned}
a &= -2j_3 + t + R - St - 1 \\
b &= -2j_1 + t + R - St - 1 \\
c &= 2j_1 + 2j_2 + 2j_3 - R + St + 1 \\
\rho &= t, \quad \rho' = 1/t \\
a' &= -a/t, \quad b' = -b/t, \quad c' = -c/t
\end{aligned} \tag{5.52}$$

The integrand of this expression is provided in a slightly different form in [8]. In fact, there the j 's are replaced by their parametrisations (4.18) giving rise to 4 independent forms [8] for the integrand depending on whether the j_i^+ or the j_i^- form is used. The above form holds in general. By analysing the small z and small x behaviour of this form it is easy to establish




Figure 5.8: Integration contours for u and v variables for an s-channel block for fusion rule I.




Figure 5.9: Integration contours for u and v variables for an s-channel block for fusion rule II.

that this conformal block corresponds to the s-channel diagram Fig. 5.1 with the intermediate j given by fusion rule I. Indeed by scaling $u_i \rightarrow zu_i$, $v_k \rightarrow zv_k$, $i \in I_1$, $k \in J_1$ we find

$$\mathcal{S}_{(r,s,0)}^{(R,S)}(z, x) \sim z^{-\Delta(j_1)-\Delta(j_2)+\Delta(j_I)} (-x)^{j_1+j_2-j_I} \quad (5.53)$$

with

$$j_I = j_1 + j_2 - r + st \quad (5.54)$$

The contours in Fig. 5.8 are essentially equal to the contours in [42] for minimal models.

5.2.2 Case of Fusion Rule II

Fig. 5.9 shows the integration contours. The s-channel block for fusion rule II is given by

$$\begin{aligned} \mathcal{S}_{(r,s,1)}^{(R,S)}(z, x) &= z^{2j_1j_2/t} (1-z)^{2j_2j_3/t} \int_0^z du_i dv_k \int_1^\infty du_j dv_l \oint_{C_v} \frac{dv}{2\pi i} \\ &\cdot u_i^{a'} (1-u_i)^{b'} (z-u_i)^{c'} \prod_{i < i'} (u_i - u_{i'})^{2\rho'} u_j^{a'} (u_j - 1)^{b'} (u_j - z)^{c'} \\ &\cdot \prod_{j < j'} (u_j - u_{j'})^{2\rho'} \prod_{i,j} (u_j - u_i)^{2\rho'} \\ &\cdot v_k^a (1-v_k)^b (z-v_k)^c \prod_{k < k'} (v_k - v_{k'})^{2\rho} v_l^a (v_l - 1)^b (v_l - z)^c \prod_{l < l'} (v_l - v_{l'})^{2\rho} \end{aligned}$$

$$\begin{aligned}
& \cdot \prod_{k,l} (v_l - v_k)^{2\rho} \prod_{i,k} (u_i - v_k)^{-2} \prod_{i,l} (u_i - v_l)^{-2} \prod_{j,k} (u_j - v_k)^{-2} \prod_{j,l} (u_j - v_l)^{-2} \\
& \cdot (u_i - x)(u_j - x)(v_k - x)^{-\rho}(v_l - x)^{-\rho} \\
& \cdot v^a(1-v)^b(v-z)^c(v-v_k)^{2\rho}(v_l - v)^{2\rho}(v-u_i)^{-2}(v-u_j)^{-2}(v-x)^{-\rho}
\end{aligned} \tag{5.55}$$

Here the variables, u_i, u_j, v_k, v_l are taken along approximately real (for z real) contours as for fusion rule I. The indices indicate: $i = 1, \dots, r; j = r+1, \dots, R; k = 1, \dots, s; l = s+1, \dots, S-1$, whereas v runs along the contour $\mathcal{C}_v$ which starts at x and surrounds both 0 and z , cf. Fig. 5.9. In addition to the s-channel blocks we have defined above, we define additional ones in analogy to the case for minimal models [42]. Namely, instead of using complex contours close to the real axis (for real z), we may use real time ordered integrations, with an ordering so that all terms in the integral expression for $\mathcal{S}_{(r,s,0)}^{(R,S)}$ become real. This block is denoted $S_{(r,s,0)}^{(R,S)}$. Similar to the case of 3-point functions one finds

$$\begin{aligned}
\mathcal{S}_{(r,s,0)}^{(R,S)}(z, x) &= \lambda_r(\rho') \lambda_{R-r}(\rho') \lambda_s(\rho) \lambda_{S-s}(\rho) S_{(r,s,0)}^{(R,S)}(z, x) \\
S_{(r,s,0)}^{(R,S)}(z, x) &= s_{(r,s,0)}^{(R,S)}(z, x) N_{(r,s,0)}^{(R,S)}
\end{aligned} \tag{5.56}$$

where $s_{(r,s,0)}^{(R,S)}(z, x)$ is normalised in such a way that the behaviour as $z \rightarrow 0, x \rightarrow 0$ is

$$s_{(r,s,0)}^{(R,S)}(z, x) = z^{-\Delta(j_1) - \Delta(j_2) + \Delta(j_I)} (-x)^{j_1 + j_2 - j_I} (1 + \mathcal{O}(z, x)) \tag{5.57}$$

The λ -functions were defined in (4.65). Similarly for fusion rule II we write

$$\begin{aligned}
\mathcal{S}_{(r,s,1)}^{(R,S)}(z, x) &= \lambda_r(\rho') \lambda_s(\rho) \lambda_{R-r}(\rho') \lambda_{S-s-1}(\rho) S_{(r,s,1)}^{(R,S)}(z, x) \\
S_{(r,s,1)}^{(R,S)}(z, x) &= s_{(r,s,1)}^{(R,S)}(z, x) N_{(r,s,1)}^{(R,S)} \\
s_{(r,s,1)}^{(R,S)}(z, x) &= z^{-\Delta(j_1) - \Delta(j_2) + \Delta(j_{II})} (-x)^{j_1 + j_2 - j_{II}} (1 + \mathcal{O}(z, x))
\end{aligned} \tag{5.58}$$

where

$$j_{II} = -j_I - 1 \tag{5.59}$$

The leading behaviour (for $z \rightarrow 0$ followed by $x \rightarrow 0$) in this case of fusion rule II is determined by the scalings

$$\begin{aligned}
u_i &\rightarrow zu_i \\
v_k &\rightarrow zv_k \\
v &\rightarrow (-x)v
\end{aligned} \tag{5.60}$$

In the limit $z, x \rightarrow 0$ the v integral becomes trivial and is carried using (2.131). The normalisation constants $N_{(r,s,0)}^{(R,S)}$ and $N_{(r,s,1)}^{(R,S)}$ are found in terms of the Dotsenko-Fateev integral (4.58) We shall need these normalisations in the calculation of crossing matrices. After some calculations we obtain

$$\begin{aligned}
N_{(r,s,0)}^{(R,S)} &= (-)^{R-r+S-s} \mathcal{J}_{r,s}(a, c; \rho) \mathcal{J}_{R-r, S-s}(a + c - 2(r - \rho s) - \rho, b; \rho) \\
&\cdot \prod_{i=0}^{R-r-1} \frac{s(a' + c' - 2(s - \rho' r) + 1 + i\rho')}{s(a' + b' + c' - 2(s - \rho' r) + 1 + \rho'(R - r - 1 + i))} \\
&\cdot \prod_{i=0}^{S-s-1} \frac{s(a + c - 2(r - \rho s) - \rho + i\rho)}{s(a + b + c - 2(r - \rho s) - \rho + \rho(S - s - 1 + i))}
\end{aligned} \tag{5.61}$$

and

$$N_{(r,s,1)}^{(R,S)} = N_{(r,s+1,0)}^{(R,S)} \frac{\Gamma(\rho)\Gamma(1-\rho)\Gamma(2-2r+a+c+2s\rho)}{\Gamma(-a-c-2\rho s+2r)\Gamma(1-r+a+s\rho)\Gamma(1-r+c+s\rho)} \cdot \frac{1}{\Gamma((s+1)\rho-r)\Gamma(2-r+a+c+(s-1)\rho)} \quad (5.62)$$

For the integral realization [8] considered here it is trivial to obtain the t-channel forms once the s-channel forms above are given. In fact we have in an obvious notation ($\epsilon = 0, 1$ for fusion rules I and II)

$$\mathcal{T}_{(r,s,\epsilon)}^{(R,S)}(z, x; j_1, j_2, j_3, j_4) = \mathcal{S}_{(r,s,\epsilon)}^{(R,S)}(1-z, 1-x; j_3, j_2, j_1, j_4) \quad (5.63)$$

We notice the following. When in the integral realization, we also transform all integration variables as $u \rightarrow 1-u$, $v \rightarrow 1-v$, the integrand for the t-channel block is identical to the one for the s-channel block, up to phases. In particular, whenever we have $(u-x)$ or $(v-x)^{-\rho}$ in the s-channel, we would have $(x-u)$ and $(x-v)^{-\rho}$ in the t-channel. Also, after transformation of the variables, the integration contours in the t-channel are between z and 1 and between 0 and $-\infty$, and the complex contour for v in the case of fusion rule II surrounds z and 1. The above factors, $(u-x)$ etc. are real provided $x < 0$ in the s-channel, or $x > 1$ in the t-channel. These two possibilities map to each other under $x \rightarrow 1-x$.

5.3 Proof of Equivalence

As previously indicated there is no absolute need for proving the equivalence between the 4-point function by PRY and the one by Andreev since both satisfy the Knizhnik-Zamolodchikov equations. Nevertheless, it is of some interest to understand better how two such seemingly very different expressions can agree, and it is rather nice to be aware of the clarification provided by the relation to the minimal model case treated in Chapter 2. Here [112] we go over several of the steps needed for a direct analytic proof. In fact, we investigate the singularity structure of the two expressions in the double limits, $z, x \rightarrow 0$, $z, x \rightarrow 1$, $z, x \rightarrow \infty$ and in the single limit, $z \rightarrow x$. We restrict ourselves to just one of the s-channel conformal blocks.

Proposition

$$\begin{aligned} & \int_0^1 du \prod_{k=1}^N dv_k v_k^{a'} (1-v_k)^{b'} (1-zv_k)^{c'} \left(1 - \frac{1-v_k}{1-zv_k} \frac{z}{x} u\right) \prod_{k < k'} (v_k - v_{k'})^{2\rho'} \\ & \cdot \prod_{i=1}^M dw_i w_i^a (1-w_i)^b (1-zw_i)^c \left(1 - \frac{1-w_i}{1-zw_i} \frac{z}{x} u\right)^{-\rho} \prod_{i < i'} (w_i - w_{i'})^{2\rho} \\ & \cdot \prod_{k,i}^{N,M} (v_k - w_i)^{-2} u^{-c-1} (1-u)^{b+c-N+(M-1)\rho} \\ & = K_{NM}^x \int_0^1 \prod_{k=1}^N dv_k v_k^{a'-\delta'} (1-v_k)^{b'+\delta'} (1-zv_k)^{c'-\delta'} \left(1 - \frac{z}{x} v_k\right) \prod_{k < k'} (v_k - v_{k'})^{2\rho'} \\ & \cdot \prod_{i=1}^M dw_i w_i^{a-\delta} (1-w_i)^{b+\delta} (1-zw_i)^{c-\delta} \left(1 - \frac{z}{x} w_i\right)^{-\rho} \prod_{i < i'} (w_i - w_{i'})^{2\rho} \end{aligned}$$

$$\cdot \prod_{k,i}^{N,M} (v_k - w_i)^{-2} \quad (5.64)$$

where

$$K_{NM}^x = \frac{\Gamma(-c)\Gamma(b+c+1-N+(M-1)\rho)}{\Gamma(b+1-N+(M-1)\rho)} K_{NM} \quad (5.65)$$

and K_{NM} is given by (2.165). Here, up to irrelevant common pre-factors, the left hand side is the PRY form of the conformal block (5.21) for $r = R = N$ and $s = S = M$ in the s-channel. We denote this by S^{PRY} . Similarly up to the same pre-factors and the new normalisation constant, K_{NM}^x , the right hand side is essentially (5.50). We denote it by S^A . Notice in particular that now we put

$$a = 2j_1, \quad b = 2j_2 + \rho, \quad c = 2j_3, \quad \rho = t \quad (5.66)$$

δ, δ' are given by the same expressions as for the minimal models

$$\begin{aligned} \delta &= a + c + 1 - N + (M - 1)\rho \\ \delta' &= a' + c' + 1 - M + (N - 1)\rho' \end{aligned} \quad (5.67)$$

Then $a - \delta$ is what was called a in previous sections, $b + \delta$ was previously called c and $c - \delta$ was previously called b . In subsequent sections we shall revert to this notation, but in this section we stick to the present notation in order to emphasise the similarity with minimal models. Also notice, that because all integrations are between 0 and 1 (after scaling by z), it is possible to deform the u integration in Fig. 5.3 to being along the real axis from 0 to 1.

Proof

We demonstrate that both the left hand side and the right hand side of the claimed identity have the same singularities in the limits $z, x \rightarrow 0$, $z, x \rightarrow 1$, $z, x \rightarrow \infty$ and $z \rightarrow x$. The z limits are the *dominant* ones, the ones taken first. The proof turns out to be rather more laborious than for the minimal models. This is due to the x dependence and the u integration in the case of S^{PRY} . However, the general strategy is entirely analogous, so we will not go over all the steps on the way.

$z, x \rightarrow 0$

The limit $z, x \rightarrow 0$ is simple to deal with and it gives rise to the normalisation constant K_{NM}^x differing from the one in the minimal models K_{NM} because of the u integration.

$z, x \rightarrow 1$

We first deal with S^A . Exactly as in the case of minimal models, we split the ordered integration ranges for the v 's in n v_k 's and m w_i 's in $(0, 1 - \epsilon)$ and $N - n$ v_l 's and $M - m$ w_j 's in $(1 - \epsilon, 1)$. Omitting integration signs and products for brevity we find

$$\begin{aligned} S_{nm}^A &\sim (1 - z)^{(N-n)(b'+c'+1)+(N-n)(N-n-1)\rho'+(M-m)(b+c+1)+(M-m)(M-m-1)\rho-2(N-n)(M-m)} \\ &\cdot (x - 1)^{N-n-(M-m)\rho} K_{NM}^x \\ &\cdot v_k^{a'-\delta'} (1 - v_k)^{b'+c'+(N-n)2\rho'-2(M-m)} (x - v_k) (v_k - v_{k'})^{2\rho'} \\ &\cdot w_i^{a-\delta} (1 - w_i)^{b+c+(M-m)2\rho-2(N-n)} (x - w_i)^{-\rho} (w_i - w_{i'})^{2\rho} (v_k - w_i)^{-2} \\ &\cdot v_l^{-b'-c'-2-(N-n-1)2\rho'+2(M-m)} (1 - v_l)^{b'+\delta'} (v_l - v_{l'})^{2\rho'} \\ &\cdot w_j^{-b-c-2-(M-m-1)2\rho+2(N-n)} (1 - w_j)^{b+\delta} (w_j - w_{j'})^{2\rho} (v_l - w_j)^{-2} \end{aligned} \quad (5.68)$$

where we have performed the same scalings as for minimal models. The above has to be summed over n and m , but for a fixed value we pick up the pure $(1-z)$ and $(x-1)$ singularity indicated. The l, j part of the integration gives immediately rise to a standard Dotsenko-Fateev integral. For the k, i part we perform the further split and scalings

$$\begin{aligned}
\int_0^1 &= \int_0^{1-\epsilon} + \int_{1-\epsilon}^1 \\
v &\rightarrow 1 - (1 - 1/x) \frac{1-v}{v} \\
\int_{1-\epsilon}^1 dv &\rightarrow \int_0^1 (1 - 1/x) \frac{dv}{v^2} \\
\int_0^1 \prod_{k=1}^n dv_k \prod_{i=1}^m dw_i &= \sum_{n_0, m_0} \int_0^{1-\epsilon} \prod_{k_0=n-n_0+1}^n dv_{k_0} \prod_{i_0=m-m_0+1}^m dw_{i_0} \\
&\cdot \int_{1-\epsilon}^1 \prod_{k=1}^{n-n_0} dv_k \prod_{i=1}^{m-m_0} dw_i
\end{aligned} \tag{5.69}$$

$$\tag{5.70}$$

In the limit $x \rightarrow 1$ we extract the $(1-x)$ power and find the coefficient again to be given by the product of two DF integrals. Analysing the gamma-functions of these we see that we can only get a non-vanishing result if $(n_0, m_0) = (n, m)$ or if $(n_0, m_0) = (n, m-1)$. We denote these cases by S_{nm}^{AI} and S_{nm}^{AII} . They will turn out to be related to fusion rules I and II. Combining everything we find the following singularities in the limit $z, x \rightarrow 1$

$$\begin{aligned}
S_{nm}^{AI} &= (1-z)^{(N-n)(b'+c'+1)+(N-n)(N-n-1)\rho'+(M-m)(b+c+1)+(M-m)(M-m-1)\rho-2(N-n)(M-m)} \\
&\cdot (x-1)^{N-n-(M-m)\rho} N(S_{nm}^{AI}) \\
S_{nm}^{AII} &= (1-z)^{(N-n)(b'+c'+1)+(N-n)(N-n-1)\rho'+(M-m)(b+c+1)+(M-m)(M-m-1)\rho-2(N-n)(M-m)} \\
&\cdot (x-1)^{b+c+1-N+n+(M-m-1)\rho} N(S_{nm}^{AII})
\end{aligned} \tag{5.71}$$

One checks that the singularities exactly correspond to fusion rules I and II. The normalisations $N(S_{nm}^{AI})$ and $N(S_{nm}^{AII})$ are found explicitly in terms of products of DF integrals to be lengthy expressions involving many products of ratios of gamma-functions.

We now turn to a similar analysis of S^{PRY} in the same limit $z \rightarrow 1$ followed by $x \rightarrow 1$. We replace $u \rightarrow 1-u$ and perform the same split and the same scalings of the v and w variables as in the case of minimal models. Omitting again integration signs and products we find

$$\begin{aligned}
S_{nm}^{PRY} &\sim (1-z)^{(N-n)(b'+c'+1)+(N-n)(N-n-1)\rho'+(M-m)(b+c+1)+(M-m)(M-m-1)\rho-2(N-n)(M-m)} \\
&\cdot v_k^{a'} (1-v_k)^{b'+c'+(N-n)2\rho'-2(M-m)} (v_k - v_{k'})^{2\rho'} \\
&\cdot w_i^a (1-w_i)^{b+c+(M-m)2\rho-2(N-n)} (w_i - w_{i'})^{2\rho} (v_k - w_i)^{-2} \\
&\cdot v_l^{-b'-c'-2-(N-n-1)2\rho'+2(M-m)} (1-v_l)^{b'} (x - (1-v_l)(1-u)) (v_l - v_{l'})^{2\rho'} \\
&\cdot w_j^{-b-c-2-(M-m-1)2\rho+2(N-n)} (1-w_j)^b (x - (1-w_j)(1-u))^{-\rho} (w_j - w_{j'})^{2\rho} \\
&\cdot (v_l - w_j)^{-2} u^{b+c-N+(M-1)\rho} (1-u)^{-c-1} (x-1+u)^{n-m\rho}
\end{aligned} \tag{5.72}$$

Here the k, i integrations are independent of x and u and are readily evaluated in terms of DF integrals, so we concentrate on the l, j part. We perform the split and the scalings

$$\int_0^1 \prod_{l=1}^{N-n} dv_l \prod_{j=1}^{M-m} dw_j = \sum_{n_0, m_0} \int_0^\epsilon \prod_{l_0=N-n-n_0+1}^{N-n} dv_{l_0} \prod_{j_0=M-m-m_0+1}^{M-m} dw_{j_0}$$

$$\begin{aligned}
& \cdot \int_{\epsilon}^1 \prod_{l=1}^{N-n-n_0} dv_l \prod_{j=1}^{M-m-m_0} dw_j \\
w & \rightarrow (x-1) \frac{1-w}{w} \\
\int_0^{\epsilon} dw & \rightarrow \int_1^{\frac{x-1}{\epsilon+x-1}} (1-x) \frac{dw}{w^2} \sim (x-1) \int_0^1 \frac{dw}{w^2}
\end{aligned} \tag{5.73}$$

and similarly for $\int_0^1 du$. To be able to distinguish we write

$$\begin{aligned}
\int_{\epsilon}^1 du & \rightarrow \int_0^1 du \\
\int_0^{\epsilon} du & \rightarrow (x-1) \int_0^1 \frac{dy}{y^2} \\
u & \rightarrow (x-1) \frac{1-y}{y}
\end{aligned} \tag{5.74}$$

and denote them the u and y cases respectively. In the u case the arising DF integrals turn out to vanish unless $n_0 = m_0 = 0$, and we find in the u case

$$\begin{aligned}
S_{nm}^{PRY,u} & \sim (1-z)^{(N-n)(b'+c'+1)+(N-n)(N-n-1)\rho'+(M-m)(b+c+1)+(M-m)(M-m-1)\rho} \\
& \cdot (1-z)^{-2(N-n)(M-m)} \\
& \cdot \prod_{k=1}^n \prod_{i=1}^m v_k^{a'} (1-v_k)^{b'+c'+(N-n)2\rho'-2(M-m)} (v_k - v_{k'})^{2\rho'} \\
& \cdot w_i^a (1-w_i)^{b+c+(M-m)2\rho-2(N-n)} (w_i - w_{i'})^{2\rho} (v_k - w_i)^{-2} \\
& \cdot \prod_{l=1}^{N-n} \prod_{j=1}^{M-m} v_l^{-b'-c'-2-(N-n-1)2\rho'+2(M-m)} (1-v_l)^{b'} (v_l - v_{l'})^{2\rho'} \\
& \cdot w_j^{-b-c-2-(M-m-1)2\rho+2(N-n)} (1-w_j)^b (w_j - w_{j'})^{2\rho} (v_l - w_j)^{-2} \\
& \cdot (1 - (1-v_l)(1-u))(1 - (1-w_j)(1-u))^{-\rho} \\
& \cdot u^{b+c-N+n+(M-m-1)\rho} (1-u)^{-c-1}
\end{aligned} \tag{5.75}$$

Now we want to establish

$$S_{NM}^{PRY,u} = S_{NM}^{AI} \tag{5.76}$$

and

$$S_{nm}^{PRY,u} = 0 \quad , \quad (n, m) \neq (N, M) \tag{5.77}$$

A straightforward analysis shows that the first one is satisfied. The second identity follows from the first generalised DF integral in (5.95).

In the y case we introduce a similar further splitting resulting in objects $S_{nm;n_0m_0}^{PRY}$. It turns out to be possible to demonstrate that

$$\begin{aligned}
S_{nm;00}^{PRY} & = S_{nm}^{AII} \\
S_{nm;01}^{PRY} & = S_{nm}^{AI} \quad , \quad m < M
\end{aligned} \tag{5.78}$$

The analysis contains no new ideas over the situation encountered for minimal models, but again the calculations involved are quite lengthy. In principle we should check that higher

values of n_0, m_0 give zero. We anticipate no interesting problems here [112]. The seemingly missing PRY counterpart of S_{nM}^{AI} for $n < N$ might be 'recovered' in the framework of the comments following (5.48).

$z, z/x, x \rightarrow \infty$

Again we first analyse the S^A case. We introduce the same splitting of integrations and the same variable transformations as for minimal models, and find

$$\begin{aligned}
S_{nm}^A &\sim K_{NM}^x (-z)^{n(-a'+\delta'-1)+(N-n)(c'-\delta'+1)-n(n-1)\rho'} \\
&\cdot (-z)^{m(-a+\delta-1)+(M-m)(c-\delta-\rho)-m(m-1)\rho+2nm} \\
&\cdot x^{-(N-n)+(M-m)\rho} \\
&\cdot v_k^{-a'-c'+2\delta'-2+2m-(n-1)2\rho'} (1-v_k)^{a'-\delta'} (v_k-v_{k'})^{2\rho'} \\
&\cdot w_i^{-a-c+2\delta-2+2n-(m-1)2\rho} (1-w_i)^{a-\delta} (w_i-w_{i'})^{2\rho} (v_k-w_i)^{-2} \\
&\cdot \left(1 + \frac{1-v_k}{xv_k}\right) \left(1 + \frac{1-w_i}{xw_i}\right)^{-\rho} \\
&\cdot v_l^{a'+c'-2\delta'+1-2m+n2\rho'} (1-v_l)^{b'+\delta'} (v_l-v_{l'})^{2\rho'} \\
&\cdot w_j^{a+c-2\delta-2n+(2m-1)\rho} (1-w_j)^{b+\delta} (w_j-w_{j'})^{2\rho} (v_l-w_j)^{-2}
\end{aligned} \tag{5.79}$$

again omitting integration signs and products, which are just as for the case of minimal models. The l, j integration is seen to result in DF integrals. In the k, i integrals we perform a split of integrals from 0 to ϵ and from ϵ to 1. In the $\int_0^\epsilon$ we transform variables like

$$\begin{aligned}
v &\rightarrow \frac{1-v}{xv} \\
1 + \frac{1-v}{xv} &\rightarrow \frac{1}{1-v} - \frac{1}{x} \sim \frac{1}{1-v} \\
\int_0^\epsilon dv &\rightarrow x^{-1} \int_0^1 \frac{dv}{v^2} \\
\int_0^1 \prod_{k=1}^n dv_k \prod_{i=1}^m dw_i &= \sum_{n_0, m_0} \int_0^\epsilon \prod_{k_0=n-n_0+1}^n dv_{k_0} \prod_{i_0=m-m_0+1}^m dw_{i_0} \\
&\cdot \int_\epsilon^1 \prod_{k=1}^{n-n_0} dv_k \prod_{i=1}^{m-m_0} dw_i
\end{aligned} \tag{5.80}$$

An analysis of the coefficients of the singularities reveals that this is non-vanishing only if $(n_0, m_0) = (0, 0)$ or if $(n_0, m_0) = (0, 1)$. These two cases we term again (we use the same notation as before, even though now we consider a different limit), S_{nm}^{AI} and S_{nm}^{AII} for what turns out to be fusion rules I and II. We find

$$\begin{aligned}
S_{nm}^{AI} &= (-z)^{-(N-n)a'+nc'-(N-n)(N-n-1)\rho'-(M-m)a+mc-(M-m)(M-m)\rho+(M-m)(2N-2n-1)} \\
&\cdot x^{-(N-n)+(M-m)\rho} N(S_{nm}^{AI}) \\
S_{nm}^{AII} &= (-z)^{-(N-n)a'+nc'-(N-n)(N-n-1)\rho'-(M-m)a+mc-(M-m)(M-m)\rho} \\
&\cdot (-z)^{(M-m)(2N-2n-1)} \\
&\cdot x^{-a-c-1+N-n+(-M+m)\rho} N(S_{nm}^{AII})
\end{aligned} \tag{5.81}$$

where the normalisations (different of course to the ones in the previous limit $z, x \rightarrow 1$), $N(S_{nm}^{AI})$ and $N(S_{nm}^{AII})$ are given (in terms of DF integrals) by lengthy products of ratios of gamma-functions. The singularities shown indicate that indeed we are dealing with fusion rules I and II.

We then treat the S^{PRY} case. Again we first perform the same splittings and variable transformations as for S^A with the same meaning of v_k, v_l, w_i, w_j . The i, k part is again simple, whereas the l, j part is treated with a split of the ordered integrations as

$$\begin{aligned} \int_0^1 \prod_{l=1}^{N-n} dv_l \prod_{j=1}^{M-m} dw_j &= \sum_{n_0, m_0} \int_0^\epsilon \prod_{l_0=N-n-n_0+1}^{N-n} dv_{l_0} \prod_{j_0=M-m-m_0+1}^{M-m} dw_{j_0} \\ &\cdot \int_\epsilon^1 \prod_{l=1}^{N-n-n_0} dv_l \prod_{j=1}^{M-m-m_0} dw_j \end{aligned} \quad (5.82)$$

followed by the scalings

$$\begin{aligned} v_{l_0} &\rightarrow \frac{1 - v_{l_0}}{x v_{l_0}} \\ w_{j_0} &\rightarrow \frac{1 - w_{j_0}}{x w_{j_0}} \end{aligned} \quad (5.83)$$

We then seek to demonstrate that

$$\begin{aligned} S_{nm;00}^{PRY} &= S_{N-n, M-m}^{AI} \\ S_{nm;01}^{PRY} &= S_{N-n, M-m}^{AII} \\ S_{nm; n_0 m_0}^{PRY} &= 0, \quad (n_0, m_0) \neq (0, 0), (0, 1) \end{aligned} \quad (5.84)$$

The proofs of the first two identities are lengthy, but with no new ideas introduced. An explicit proof of the last identity is lacking, but again we anticipate no problems.

$z \rightarrow x$

This case is the most complicated one. We omit nearly all the details, most of which are similar to what have been described above. Let us outline the strategy. First one may check that the nature of the singularity is a linear combination of just two different powers of $(z - x)$ namely either $(z - x)^0$ or $(z - x)^{(c+1-\rho)}$. Second one must investigate whether the coefficients of these two powers are the same for the two sides of (5.64). That coefficient is a function of x in the limit $z \rightarrow x$, so we must investigate whether the coefficient functions defined by the two sides of (5.64) are equal. As above the technique is to investigate the singularity structure in the singular limits $x = 0, 1, \infty$.

Let us return to the first issue. Here it is suggested to denote the channel or basis corresponding to the limit $z \rightarrow x$ by v , inspired by the alphabet s, t, u, v . Consider for fusion rule I the general real s -channel block $S_{(r,s,0)}^{(R,S)}(z, x)$ and perform the transformations $u_i, v_k \rightarrow z u_i, z v_k$ and $u_j, v_l \rightarrow 1/u_j, 1/v_l$ whereby all integrations become $\int_0^1$

$$S_{(r,s,0)}^{(R,S)}(z, x) \sim u_i^{a'} (1 - z u_i)^{b'} (1 - u_i)^{c'} (u_i - u_{i'})^{2\rho'}$$

$$\begin{aligned}
& \cdot u_j^{-a'-b'-c'}(1-u_j)^{b'}(1-zu_j)^{c'}(u_j-u_{j'})^{2\rho'} \\
& \cdot u_j^{-2\rho'(R-r-1)}u_j^{-2\rho'}(1-zu_iu_j)^{2\rho'}u_j^{-2} \\
& \cdot v_k^a(1-zv_k)^b(1-v_k)^c(v_k-v_{k'})^{2\rho} \\
& \cdot v_l^{-a-b-c}(1-v_l)^b(1-zv_l)^c(v_l-v_{l'})^{2\rho}v_l^{-2\rho(S-s-1)} \\
& \cdot v_l^{-2\rho}(1-zv_kv_l)^{2\rho}v_l^{-2} \\
& \cdot (u_i-v_k)^{-2}(1-zu_iv_l)^{-2}v_l^2(1-zv_ku_j)^{-2}u_j^2(u_j-v_l)^{-2}u_j^{2(S-s-1)}v_l^{2(R-r-1)} \\
& \cdot \left(1-\frac{z}{x}u_i\right)\left(1-\frac{z}{x}v_k\right)^{-\rho}u_j^{-1}(1-xu_j)v_l^\rho(1-xv_l)^{-\rho}
\end{aligned} \tag{5.85}$$

In these considerations we leave out integration signs and products. We notice that only u_i and v_k may scale wrt z/x and let us denote the r_0, s_0 ones that do by w_i and w_k

$$\begin{aligned}
u_i & \rightarrow 1 - \left(1 - \frac{z}{x}\right) \frac{1-w_i}{w_i} \\
v_k & \rightarrow 1 - \left(1 - \frac{z}{x}\right) \frac{1-w_k}{w_k}
\end{aligned} \tag{5.86}$$

In the limit $z \rightarrow x$ the w integrations decouple

$$\begin{aligned}
& \prod_{i=1}^{r_0} dw_i \prod_{k=1}^{s_0} dw_k w_i^{-c'-2\rho'(r_0-1)-2+2s_0-1} (1-w_i)^{c'} (w_i-w_{i'})^{2\rho'} \\
& \cdot w_k^{-c-2\rho(s_0-1)-2+2r_0+\rho} (1-w_k)^c (w_k-w_{k'})^{2\rho} (w_i-w_k)^{-2}
\end{aligned} \tag{5.87}$$

from which one deduces that $r_0 = 0$ and $s_0 = 0, 1$. This gives that the only possible powers of $z-x$ are

$$(z-x)^{s_0(c+1-\rho)}, \quad s_0 = 0, 1 \tag{5.88}$$

A special feature arises for $s = 0$ since then $s_0 = 0$ and in the corresponding rows there is only one matrix element different from zero. The two possible v-channel blocks are easily expressed in terms of simple contours and will be denoted $V_1^{(R,S)}$ and $V_2^{(R,S)}$

$$\begin{aligned}
V_1^{(R,S)} &= S_{(0,0,0)}^{(R,S)} \\
V_2^{(R,S)} &= \int_1^\infty \prod_{j=1, l=1}^{R, S-1} du_j dv_l \int_z^x dw \\
& \cdot u_j^{a'}(u_j-1)^{b'}(u_j-z)^{c'}(u_j-x)(u_j-u_{j'})^{2\rho'} \\
& \cdot v_l^a(v_l-1)^b(v_l-z)^c(v_l-x)^{-\rho}(v_l-v_{l'})^{2\rho}(u_j-v_l)^{-2} \\
& \cdot w^a(1-w)^b(w-z)^c(x-w)^{-\rho}(v_l-w)^{2\rho}(u_j-w)^{-2}
\end{aligned} \tag{5.89}$$

We have already seen that V_1 has the correct $(z-x)^0$ behaviour. To see that V_2 has the correct $z-x$ singularity we make the transformations $u_j, v_l \rightarrow 1/u_j, 1/v_l$ and $v \rightarrow (x-z)v + z$ whereby all integrations become $\int_0^1$ and one easily finds the singular behaviour

$$(x-z)^{c-\rho+1} \tag{5.90}$$

The situation for fusion rule II is very analogous. We make the transformations $u_i, v_k, v \rightarrow zu_i, zv_k, xv$ and $u_j, v_l \rightarrow 1/u_j, 1/v_l$ whereby we obtain (up to irrelevant phases)

$$\begin{aligned}
S_{(r,s,1)}^{(R,S)} \sim & \oint_{S^1} \prod \int_0^1 u_i^{a'} (1 - zu_i)^{b'} (1 - u_i)^{c'} (u_i - u_{i'})^{2\rho'} \\
& \cdot u_j^{-a'-b'-c'} (1 - u_j)^{b'} (1 - zu_j)^{c'} (u_j - u_{j'})^{2\rho'} u_j^{-2\rho'(R-r-1)} \\
& \cdot u_j^{-2\rho'} (1 - zu_i u_j)^{2\rho'} u_j^{-2} \\
& \cdot v_k^a (1 - zv_k)^b (1 - v_k)^c (v_k - v_{k'})^{2\rho} \\
& \cdot v_l^{-a-b-c} (1 - v_l)^b (1 - zv_l)^c (v_l - v_{l'})^{2\rho} v_l^{-2\rho(S-s-1)} \\
& \cdot v_l^{-2\rho} (1 - zv_k v_l)^{2\rho} v_l^{-2} \\
& \cdot (u_i - v_k)^{-2} (1 - zu_i v_l)^{-2} v_l^2 (1 - zv_k u_j)^{-2} u_j^2 (u_j - v_l)^{-2} \\
& \cdot u_j^{2(S-s-1)} v_l^{2(R-r-1)} \left(1 - \frac{z}{x} u_i\right) \left(1 - \frac{z}{x} v_k\right)^{-\rho} \\
& \cdot u_j^{-1} (1 - xu_j) v_l^\rho (1 - xv_l)^{-\rho} \\
& \cdot v^a (1 - xv)^b \left(1 - \frac{x}{z} v\right)^c \left(v - \frac{z}{x} v_k\right)^{2\rho} (1 - xvv_l)^{2\rho} v_l^{-2\rho} \\
& \cdot \left(v - \frac{z}{x} u_i\right)^{-2} (1 - xvu_i)^{-2} u_j^2 (1 - v)^{-\rho}
\end{aligned} \tag{5.91}$$

As for fusion rule I it is relevant to scale u_i, v_k but this time also v . If v is not scaled we find exactly the same decoupling of the transformed variables (denoted w_i, w_k). Again this indicates that we only need $V_1^{(R,S)}$ and $V_2^{(R,S)}$. However, if v does scale the situation is much more complicated to handle. Though, we anticipate that it does not lead to any new v-blocks.

Let us turn to the second issue. It turns out that the sought equality (5.64) depends on the following identities

$$\begin{aligned}
& \int_0^1 dw dy w^{-a-2+\rho} (1-w)^a (1-(1-w)(1-(1-x)y))^{-\rho} \\
& \cdot y^{-b-c+N-2+(-M+2)\rho} (1-y)^{b+c-N+(M-1)\rho} \\
= & (1-x)^{-a-1} \frac{\Gamma(a+1)\Gamma(b+c+1-N+(M-1)\rho)\Gamma(-a-b-c-2+N+(-M+2)\rho)}{\Gamma(\rho)}
\end{aligned} \tag{5.92}$$

and

$$\int_0^1 dw \int_0^w dy w^{-a} (1-w)^{a+c-2} (w-y)^{-c} y^{b+c-2} (1-y)^{-b} = 0 \tag{5.93}$$

which are not too difficult to prove. Next define the Dotsenko-Fateev integrand

$$\begin{aligned}
& DF(N, M; a, b, \rho; \{v_k\}, \{w_i\}) \\
= & \prod_{k=1}^N v_k^{a'} (1 - v_k)^{b'} \prod_{k < k'} (v_k - v_{k'})^{2\rho'} \prod_{i=1}^M w_i^a (1 - w_i)^b \prod_{i < i'} (w_i - w_{i'})^{2\rho} \prod_{k,i}^{N,M} (v_k - w_i)^{-2}
\end{aligned} \tag{5.94}$$

Then we find that the equality of the two sides of (5.64) depends on the following three identities (generalised DF integrals)

$$\begin{aligned}
(I) \quad & \int_0^1 du \prod_{k=1}^N dv_k \prod_{i=1}^M dw_i DF(N, M; a, b, \rho; \{v_k\}, \{w_i\}) \\
& \cdot (1 - (1 - v_k)u)(1 - (1 - w_i)u)^{-\rho} u^{-c-1} (1 - u)^{b+c-N+(M-1)\rho} \\
& = \frac{\Gamma(-c)\Gamma(b+c+1-N+(M-1)\rho)\Gamma(a+b+c+2-N+(M-2)\rho)}{\Gamma(b+1-\rho)\Gamma(a+b+c+2-2N+(2M-2)\rho)} \\
& \cdot \int_0^1 \prod_{k=1}^N dv_k \prod_{i=1}^M dw_i DF(N, M; a, b-\rho, \rho; \{v_k\}, \{w_i\}) \\
(II) \quad & \int_0^1 du \prod_{k=1}^n dv_k \prod_{i=1}^{m+1} dw_i \\
& \cdot DF(n, m+1; a, b+c-2N+2n+(2M-2m-1)\rho, \rho; \{v_k\}, \{w_i\}) \\
& \cdot (1 - v_k - u)(1 - w_i - u)^{-\rho} u^{-b-c+N-n-2+(-M+m+2)\rho} (1 - u)^{b+c-N+(M-1)\rho} \\
& \sim \frac{\Gamma(b+c+1-N+(M-1)\rho)\Gamma(a+b+c+2-N+(M-2)\rho)}{\Gamma(a+b+c+2-2N+(2M-2)\rho)} \\
& \cdot \frac{\Gamma(-b-c+N-n-1+(-M+m+2)\rho)\Gamma(-N+n+(M-m)\rho)}{\Gamma(\rho)} \\
& \cdot \int_0^1 \prod_{k=1}^n dv_k \prod_{i=1}^m dw_i \\
& \cdot DF(n, m; a, b+c-2N+2n+(2M-2m-1)\rho, \rho; \{v_k\}, \{w_i\}) \\
(III) \quad & \int_0^1 du \prod_{k=1}^n dv_k \prod_{i=1}^m dw_i \\
& \cdot DF(n, m; -a-c+2n-2+(-2m+2)\rho, a, \rho; \{v_k\}, \{w_i\}) \\
& \cdot (1 - v_k u)(1 - w_i u)^{-\rho} u^{-c-1} (1 - u)^{b+c-N+(M-1)\rho} \\
& = \frac{\Gamma(-c)\Gamma(b+c+1-N+(M-1)\rho)\Gamma(a+b+c+2-N+(M-2)\rho)}{\Gamma(b+1-N+n+(M-m-1)\rho)\Gamma(a+b+c+2-N-n+(M+m-2)\rho)} \\
& \cdot \int_0^1 \prod_{k=1}^n dv_k \prod_{i=1}^m dw_i DF(n, m; -a-c+2n-2+(-2m+2)\rho, a, \rho; \{v_k\}, \{w_i\})
\end{aligned} \tag{5.95}$$

All the final integrals are of course DF integrals. In the second identity there is a phase depending on the precise choice of the integration contour for u . These last three identities have not been proven directly, but are easily checked for low values of N, M . One may take the attitude that the undoubted identity of the PRY realization and that of Andreev, i.e. the unquestionable correctness of (5.64), implies these somewhat remarkable integral identities in addition to the ones underlying the cases discussed above.

Let us supply with some details on the limit $z \rightarrow x$. We already know that the rhs of (5.64) vanishes except in the two cases $(r_0, s_0) = (0, 0), (0, 1)$. The notation used there is different from the one used otherwise and to which we will stick here. The translation of $(r_0, s_0) = (0, 0), (0, 1)$

gives $(n, m) = (N, M), (N, M - 1)$. In conclusion we find for the Andreev representation

$$\begin{aligned}
S_{NM}^A &\sim K_{NM}^x \prod_{k=1}^N v_k^{a'-\delta'} (1-v_k)^{b'+\delta'+1} (1-xv_k)^{c'-\delta'} (v_k-v_{k'})^{2\rho'} \\
&\cdot \prod_{i=1}^M w_i^{a-\delta} (1-w_i)^{b+\delta-\rho} (1-xw_i)^{c-\delta} (w_i-w_{i'})^{2\rho} (v_k-w_i)^{-2} \\
&= \frac{\Gamma(-c)\Gamma(b+c+1-N+(M-1)\rho)\Gamma(a+b+c+2-N+(M-2)\rho)}{\Gamma(b+1-\rho)\Gamma(a+b+c+2-2N+(2M-2)\rho)} \\
&\cdot v_k^{a'} (1-v_k)^{b'+1} (1-xv_k)^{c'} (v_k-v_{k'})^{2\rho'} \\
&\cdot w_i^a (1-w_i)^{b-\rho} (1-xw_i)^c (w_i-w_{i'})^{2\rho} (v_k-w_i)^{-2}
\end{aligned} \tag{5.96}$$

and

$$\begin{aligned}
S_{N,M-1}^A &\sim (1-z/x)^{b+\delta-\rho+1} (1-x)^{c-\delta} \\
&\cdot K_{NM}^x \prod_{k=1}^N v_k^{a'-\delta'} (1-v_k)^{b'+\delta'-1} (1-xv_k)^{c'-\delta'} (v_k-v_{k'})^{2\rho'} \\
&\cdot \prod_{i=1}^{M-1} w_i^{a-\delta} (1-w_i)^{b+\delta+\rho} (1-xw_i)^{c-\delta} (w_i-w_{i'})^{2\rho} (v_k-w_i)^{-2} \\
&\cdot w^{-b-\delta+\rho-2} (1-w)^{b+\delta} \\
&= (1-z/x)^{a+b+c+2-N+(M-2)\rho} (1-x)^{-a-1+N+(-M+1)\rho} \\
&\cdot \frac{\Gamma(a+1)\Gamma(b+c+1-N+(M-1)\rho)\Gamma(-a-b-c-2+N+(-M+2)\rho)}{\Gamma(\rho)} \\
&\cdot \prod_{k=1}^N v_k^{a'-1} (1-v_k)^{b'} (1-xv_k)^{c'-1} (v_k-v_{k'})^{2\rho'} \\
&\cdot \prod_{i=1}^{M-1} w_i^{a+\rho} (1-w_i)^b (1-xw_i)^{c+\rho} (w_i-w_{i'})^{2\rho} (v_k-w_i)^{-2}
\end{aligned} \tag{5.97}$$

The rewritings are due to the integral identity for minimal models (2.163). Again we have used the same notation as in the other limits, but as long we do not make any direct comparisons between results obtained in the different limits, it should not lead to any confusion.

Now we turn to PRY. We will have to divide into two cases; $V^{PRY,u}$ where u is not scaled and $V^{PRY,y}$ where u is scaled

$$u \rightarrow (1-z/x) \frac{1-y}{y} \tag{5.98}$$

In both cases we let n v 's and m w 's transform and denote the cases $S_{nm}^{PRY,u}$ and $S_{nm}^{PRY,y}$. Several lengthy and cumbersome manipulations using the various integral identities, lead to the following identities

$$S_{NM}^A \sim S_{00}^{PRY,u}, \quad S_{N,M-1}^A \sim S_{01}^{PRY,y} \tag{5.99}$$

Without going into these final and comprehensive computations we conclude that the above proof presents substantial evidence for the claimed integral identity (5.64) at the level of direct

verification. Taking the aforementioned attitude of accepting the equivalence of the PRY and Andreev representations from the point of view of both representations producing solutions to the KZ equations, the above assumptions are all justified.

□

5.4 Over-screening

The 4-point blocks considered so far are ones where we have either fusion rule I operating at both vertices, or fusion rule II operating at both vertices. We now describe how to obtain 4-point blocks for the case where we have either fusion rule I for $j_1 j_2 j$ and fusion rule II for $j j_3 j_4$, we denote that case by (II,I), or fusion rule II for $j_1 j_2 j$ and fusion rule I for $j j_3 j_4$, we denote that case by (I,II). We emphasise that for a collection of spins considered so far, so that fusion rule I (or fusion rule II) is possible at both vertices, neither (I,II) nor (II,I) will be possible. Hence there will be no mixing in the crossing matrix calculations.

Our technique is based on the discussion of fusion rules I and II for the 3-point function in Chapter 4. Now the charge conservation reads

$$j_1 + j_2 + j_3 - j_4 = R - St \quad (5.100)$$

and as for the charge conservation of the 3-point function there is a considerable freedom in choosing the numbers of screenings. Let us introduce the following three choices

Standard Screening

$$\begin{aligned} 2R &= r_1 + r_2 + r_3 - r_4 - 2 \\ 2S &= s_1 + s_2 + s_3 - s_4 \end{aligned} \quad (5.101)$$

Over-screening

$$\begin{aligned} 2R &= r_1 + r_2 + r_3 - r_4 - 2 + p \\ 2S &= s_1 + s_2 + s_3 - s_4 + q \end{aligned} \quad (5.102)$$

Reducible Over-screening

$$\begin{aligned} 2R &= r_1 + r_2 + r_3 - r_4 - 2 + 2p \\ 2S &= s_1 + s_2 + s_3 - s_4 + 2q \end{aligned} \quad (5.103)$$

Any 4-point function may be factorised into a product of two 3-point functions, but in the previous work based on (5.101) this fact has been somewhat unclear in the case of fusion rule II. There is apparently a very simple explanation for that. According to the discussion of 3-point functions, the 'natural' way to obtain a fusion rule II 4-point function is to use (5.103), an over-screening by p, q at both vertices. However, the total addition of $2p, 2q$ might be reducible since it involves only even numbers. Indeed this is what we find since we have been able to define 4-point functions for fusion rule II using the standard screening (5.101). The reducibility obscures the factorisation.

Let us first notice that (5.102) indeed provides us with a new family of 4-point functions (if we can find contours such that the blocks are well-defined and non-vanishing). This follows

from the fact that there is no overlap between fusion rule I and II, such that a block based on (II,I) will be different from blocks based on other combinations. Here the notation (II,I) refers to the fusion rules involved in the (left,right) vertices. There are two cases of interest

(I,II)

$$\begin{aligned}(\rho_1, \sigma_1) &= \left(\frac{r_1 + r_2 - r - 1 + p}{2}, \frac{s_1 + s_2 - s + q}{2} \right) \\(\rho_2, \sigma_2) &= \left(\frac{r + r_3 - r_4 - 1}{2}, \frac{s + s_3 - s_4}{2} \right) \\(R, S) &= (\rho_1 + \rho_2, \sigma_1 + \sigma_2)\end{aligned}\tag{5.104}$$

(II,I)

$$\begin{aligned}(\rho_1, \sigma_1) &= \left(\frac{r_1 + r_2 - r - 1}{2}, \frac{s_1 + s_2 - s}{2} \right) \\(\rho_2, \sigma_2) &= \left(\frac{r + r_3 - r_4 - 1 + p}{2}, \frac{s + s_3 - s_4 + q}{2} \right) \\(R, S) &= (\rho_1 + \rho_2, \sigma_1 + \sigma_2)\end{aligned}\tag{5.105}$$

(ρ_1, σ_1) are the numbers of screenings around the right vertex $j_1 j_2 j$ while (ρ_2, σ_2) are the numbers of screenings around the left vertex $j j_3 j_4$. The internal state is parameterised as

$$2j + 1 = r - st\tag{5.106}$$

R and S are the total numbers of screenings.

At both vertices we choose contours similar to the 3-point functions. Compared to the contours in Fig. 5.8 this amounts to substitute (in the case (I,II)) the σ_1 DF contours for the screenings of the second kind around the right vertex by Felder contours. More precisely, we may take $v_1, \dots, v_s$ to run along circle like contours passing through the point z , and surrounding the contours for $u_1, \dots, u_r$, i.e. surrounding the point 0, in such a way that the v_1 -contour lies inside the v_2 -contour, etc. Similarly for the case of (II,I), we modify the contours for $v_{s+1}, \dots, v_S$ into circle like contours surrounding 0 (actually surrounding ∞) and passing through the point 1, in such a way that the contour for v_{s+1} lies inside the contour for v_{s+2} etc.

Thus, in the case (I,II) we want to establish the following

(I,II)

$$\lambda_{\rho_1}(1/t) \chi_{\sigma_1}^{(2)}(S - s_3 - 1; t) \lambda_{\rho_2}(1/t) \lambda_{\sigma_2}(t) N_{(\rho_1, \sigma_1, 0)}^{(R, S)} \neq 0, \infty\tag{5.107}$$

The argument of the χ function follows from the fact that the deformation of Felder contours only depends on the number of integration variables and (minus) the fractional part of the common power of the w_i 's (not on the power of $(1 - w_i)$). In this case the power is $a = -2j_3 + t + R - St - 1$ so the fractional part is $(s_3 - S + 1)t$. Using (4.60) one obtains the normalisation

$$\begin{aligned}N_{(\rho_1, \sigma_1, 0)}^{(R, S)} &= \mathcal{J}_{\rho_1, \sigma_1}(a, c; t) \tilde{\mathcal{J}}_{R - \rho_1, S - \sigma_1}(a + c - 2(\rho_1 - \sigma_1)t - t, b; t) \\&= \mathcal{J}_{\rho_1, \sigma_1}(a, c; t) \mathcal{J}_{\rho_2, \sigma_2}(-a - b - c + 2(R - 1) - 2(S - 1)t + t, b; t) \\&= t^{2\rho_1\sigma_1} \prod_{j=1}^{\rho_1} \frac{\Gamma(j/t)}{\Gamma(1/t)} \prod_{j=1}^{\sigma_1} \frac{\Gamma(jt - \rho_1)}{\Gamma(t)}\end{aligned}$$

$$\begin{aligned}
& \cdot \prod_{j=0}^{\rho_1-1} \frac{\Gamma(S - s_3 + (r_3 - R + j)/t)}{\Gamma(1 - 2\sigma_1 + s_1 + s_2 + (1 + \rho_1 - r_1 - r_2 + j)/t)} \\
& \cdot \prod_{j=0}^{\rho_1-1} \Gamma(1 + s_1 + s_2 + s_3 - S + (2 + R - r_1 - r_2 - r_3 + j)/t) \\
& \cdot \prod_{j=0}^{\sigma_1-1} \frac{\Gamma(1 - \rho_1 + R - r_3 + (s_3 - S + 1 + j)t)}{\Gamma(r_1 + r_2 - \rho_1 + (\sigma_1 - s_1 - s_2 + j)t)} \\
& \cdot \prod_{j=0}^{\sigma_1-1} \Gamma(r_1 + r_2 + r_3 - 1 - \rho_1 - R + (S - s_1 - s_2 - s_3 + j)t) \\
& \cdot t^{2\rho_2\sigma_2} \prod_{j=1}^{\rho_2} \frac{\Gamma(j/t)}{\Gamma(1/t)} \prod_{j=1}^{\sigma_2} \frac{\Gamma(jt - \rho_2)}{\Gamma(t)} \\
& \cdot \prod_{j=0}^{\rho_2-1} \frac{\Gamma(S - s_2 + (r_2 - R + j)/t) \Gamma(S - s_1 + (r_1 - R + j)/t)}{\Gamma(2\sigma_1 - s_1 - s_2 + (r_1 + r_2 - 1 - R - \rho_1 + j)/t)} \\
& \cdot \prod_{j=0}^{\sigma_2-1} \frac{\Gamma(1 + \rho_1 - r_2 + (s_2 + 1 - S + j)t) \Gamma(1 + \rho_1 - r_1 + (s_1 + 1 - S + j)t)}{\Gamma(2 + R + \rho_1 - r_1 - r_2 + (1 - S - \sigma_1 + s_1 + s_2 + j)t)}
\end{aligned} \tag{5.108}$$

According to both fusion rule I and II

$$\rho_i < p \quad \sigma_i < q \tag{5.109}$$

so

$$\lambda_{\rho_i} \neq 0 \neq \lambda_{\sigma_i} \tag{5.110}$$

and we may concentrate on χ and N . The remaining analysis is straightforward (but cumbersome) and equivalent to the one employed in the discussion of 3-point functions: using the relevant fusion rules, we consider the gamma- and sine-functions one by one in order to determine the netto number of $\Gamma(0)$'s ($\sim 1/s(0)$'s). As expected, one finds the cancellation $(\Gamma(0)/\Gamma(0))^2$ from the part concerning the right vertex (fusion rule II). The expectation is due to the factorisation into 3-point functions and the similar experience from considering 3-point functions.

The analysis of the alternative possibility (5.105) is analogous. Now we want to show

(II,I)

$$(-1)^{\sigma_2} \lambda_{\rho_1}(1/t) \lambda_{\sigma_1}(t) \lambda_{\rho_2}(1/t) \chi_{\sigma_2}^{(2)}(S - s_2 - 1; t) N_{(\rho_1, \sigma_1, 0)}^{(R, S)} \neq 0, \infty \tag{5.111}$$

This time we encounter the cancellation $(\Gamma(0)/\Gamma(0))^2$ at the left vertex, again as expected.

Having performed the analysis of the 4-point function based on over-screening (5.102) it is a simple matter to verify that reducible over-screening (5.103) indeed leads to well-defined and non-vanishing 4-point functions. We choose ρ_1 DF contours and σ_1 Felder contours around the right vertex along with ρ_2 DF contours and σ_2 Felder contours around the left vertex. The numbers of screenings are

(II,II)

$$(\rho_1, \sigma_1) = \left(\frac{r_1 + r_2 - r - 1 + p}{2}, \frac{s_1 + s_2 - s + q}{2} \right)$$

$$\begin{aligned}
(\rho_2, \sigma_2) &= \left(\frac{r + r_3 - r_4 - 1 + p}{2}, \frac{s + s_3 - s_4 + q}{2} \right) \\
(R, S) &= (\rho_1 + \rho_2, \sigma_1 + \sigma_2)
\end{aligned} \tag{5.112}$$

The analysis of the corresponding 4-point function is most easily performed using the techniques above

$$\mathcal{S}_{(\rho_1, \sigma_1, (II, II))}^{(R, S)} = (-1)^{\sigma_2} \lambda_{\rho_1}(1/t) \chi_{\sigma_1}^{(2)}(S - s_3 - 1; t) \lambda_{\rho_2}(1/t) \chi_{\sigma_2}^{(2)}(S - s_2 - 1; t) N_{(\rho_1, \sigma_1, 0)}^{(R, S)} \tag{5.113}$$

Remembering that for the over-screened 4-point function based on (II,I) and (I,II) all $\Gamma(0)$'s appeared at the over-screened vertices, we may conclude that in the present case there will be exactly 4 cancellations of the form $\Gamma(0)/\Gamma(0)$ leaving $\mathcal{S}_{(\rho_1, \sigma_1, (II, II))}^{(R, S)}$ well-defined and non-vanishing.

The representation (5.113) gives us an alternative form of that block from the one considered so far, but one with more screenings (more integrations in the case of Andreev's representation). Our previous treatment is the most economic as far as the numbers of screenings are concerned, and indeed it is more convenient when considering the crossing matrix (next section) since our technique there is based on contour deformations and becomes 'singular' when different numbers of contours are in play. We will have more to say about 4-point functions based on over-screening when discussing operator algebra coefficients in a subsequent section.

5.5 Crossing Matrix

The crossing matrix $\alpha_{(r, s, \epsilon), (r', s', \epsilon')}^{(R, S)}$ is defined by the equation

$$\begin{aligned}
S_{(r, s, \epsilon)}^{(R, S)}(z, x) &= \sum_{r'=0}^R \sum_{s'=0}^S \alpha_{(r, s, \epsilon), (r', s', 0)}^{(R, S)} T_{(r', s', 0)}^{(R, S)}(z, x) \\
&+ \sum_{r'=0}^R \sum_{s'=0}^{S-1} \alpha_{(r, s, \epsilon), (r', s', 1)}^{(R, S)} T_{(r', s', 1)}^{(R, S)}(z, x)
\end{aligned} \tag{5.114}$$

As explained in [42], it is enough to calculate one column and one row of this matrix in order to determine monodromy invariant Greens functions. It turns out that a moderate modification of the techniques described there suffices for completing the corresponding calculations here following [112]. The main new feature is the fact that we have to observe also the x dependence, and the presence of the complex contour in the case of conformal blocks for fusion rule II, cf. Fig. 5.9.

We will not discuss the case of over-screening in this context but refer to the section below on Greens functions.

5.5.1 The Column of the Transformation Matrix

Following the idea of [42] we define the following object [112] (suppressing several variables)

$$J(r_1, s_1, r_2, s_2, r_3, s_3) = z^{2j_1 j_2 / t} (1 - z)^{2j_2 j_3 / t}$$

$$\begin{aligned}
& \cdot \int_0^z \prod_{i=1}^{r_1} du_i \prod_{k=1}^{s_1} dv_k \int_z^1 \prod_{m=r_1+1}^{r_1+r_2} du_m \prod_{n=s_1+1}^{s_1+s_2} dv_n \int_1^\infty \prod_{j=r_1+r_2+1}^{r_1+r_2+r_3} du_j \prod_{l=s_1+s_2+1}^{s_1+s_2+s_3} dv_l \\
& \cdot \prod_i u_i^{a'} (1-u_i)^{b'} (z-u_i)^{c'} \prod_{i<i'} (u_i-u_{i'})^{2\rho'} (u_i-x) \\
& \cdot \prod_m u_m^{a'} (1-u_m)^{b'} (u_m-z)^{c'} \prod_{m<m'} (u_m-u_{m'})^{2\rho'} (x-u_m) \\
& \cdot \prod_j u_j^{a'} (u_j-1)^{b'} (u_j-z)^{c'} \prod_{j<j'} (u_j-u_{j'})^{2\rho'} (u_j-x) \\
& \cdot \prod_{m,i} (u_m-u_i)^{2\rho'} \prod_{j,i} (u_j-u_i)^{2\rho'} \prod_{j,m} (u_j-u_m)^{2\rho'} \\
& \cdot \prod_k v_k^a (1-v_k)^b (z-v_k)^c \prod_{k<k'} (v_k-v_{k'})^{2\rho} (v_k-x)^{-\rho} \\
& \cdot \prod_n v_n^a (1-v_n)^b (v_n-z)^c \prod_{n<n'} (v_n-v_{n'})^{2\rho} (x-v_n)^{-\rho} \\
& \cdot \prod_l v_l^a (v_l-1)^b (v_l-z)^c \prod_{l<l'} (v_l-v_{l'})^{2\rho} (v_l-x)^{-\rho} \\
& \cdot \prod_{n,k} (v_n-v_k)^{2\rho} \prod_{l,k} (v_l-v_k)^{2\rho} \prod_{l,n} (v_l-v_n)^{2\rho} \\
& \cdot \prod_{\alpha,\beta} (v_\alpha-u_\beta)^{-2}
\end{aligned} \tag{5.115}$$

In other words, there are r_1 and s_1 u and v integrations between 0 and z , r_2 and s_2 u and v integrations between z and 1 and r_3 and s_3 u and v integrations between 1 and ∞ . Also the variables, u_i, v_k, u_j, v_l are taken along contours similar to the ones in Fig. 5.8, whereas the variables, u_m, v_n are taken along similar ones lying between z and 1. We notice that

$$\begin{aligned}
J(r, s, 0, 0, R-r, S-s) &= \mathcal{S}_{(r,s,0)}^{(R,S)} \\
J(0, 0, R, S, 0, 0) &= \mathcal{T}_{(R,S,0)}^{(R,S)}
\end{aligned} \tag{5.116}$$

Therefore we may start from $J(r, s, 0, 0, R-r, S-s)$ and gradually move integration contours by contour deformations on to the interval $(z, 1)$. In the process we pick up contributions from integrals between $-\infty$ and 0, but these may be neglected in the calculation of the column, $\alpha_{(r,s,\epsilon),(R,S,0)}^{(R,S)}$. The calculational procedure consists in deforming upper and lower u and v contours in appropriate ways, and forming suitable linear combinations of the result. As explained in [42], one may then derive identities for the functions, $J(r_1, s_1, r_2, s_2, r_3, s_3)$, by carefully keeping track of the phases arising between the result of the deformations, and the definitions of the J 's. The useful identities turn out to be after some calculations

$$\begin{aligned}
& J(r_1, s_1, r_2, s_2, r_3, s_3) \\
& = e^{i\pi\rho'(r_2-r_1+1)} \frac{s(b'+\rho'(r_2+r_3))}{s(b'+c'+\rho'(r_1-1+2r_2+r_3))} J(r_1-1, s_1, r_2+1, s_2, r_3, s_3) + \dots \\
& = -e^{i\pi\rho(s_2-s_1+2)} \frac{s(b+\rho(s_2+s_3))}{s(b+c+\rho(s_1-1+2s_2+s_3))} J(r_1, s_1-1, r_2, s_2+1, r_3, s_3) + \dots \\
& \cdot J(0, 0, r_2, s_2, r_3, s_3) \\
& = e^{i\pi\rho'(r_2-r_3+1)} \frac{s(c'+\rho'r_2)}{s(b'+c'+\rho'(2r_2+r_3-1))} J(0, 0, r_2+1, s_2, r_3-1, s_3) + \dots
\end{aligned}$$

$$= -e^{i\pi\rho(s_2-s_3+2)} \frac{s(c+\rho s_2)}{s(b+c+\rho(2s_2+s_3-1))} J(0,0,r_2,s_2+1,r_3,s_3-1) + \dots \quad (5.117)$$

Here the dots stand for terms that cannot contribute to the crossing matrix element. Let us illustrate the techniques involved by deriving the very first identity in (5.117). We deform the upper u variable in the interval $(0, z)$ on to the intervals $(-\infty, 0)$, $(1, z)$ and $(+\infty, 1)$. This will result in a relation of the form

$$J(r_1, s_1, r_2, s_2, r_3, s_3) = \phi_1 J(r_1 - 1, s_1, r_2 + 1, s_2, r_3, s_3) + \phi_2 J(r_1 - 1, s_1, r_2, s_2, r_3 + 1, s_3) + \dots \quad (5.118)$$

where the dots stand for contributions from integration over the interval $(-\infty, 0)$ which cannot have a projection onto the t-channel block we are interested in, which is $J(0, 0, R, S, 0, 0)$. We call the upper u variable simply u and only consider the u parts of the various integrands. Let us denote the sets of indices for the u variables as I_1, I_2, I_3 when the variables are in the intervals $(0, z)$, $(z, 1)$, $(1, \infty)$ respectively. Likewise we denote the index sets for the v variables as J_1, J_2, J_3 when the v variables are in the same intervals. Then we have before we transform $J(r_1, s_1, r_2, s_2, r_3, s_3)$

$$\int_0^z du u^{a'} (1-u)^{b'} (z-u)^{c'} \prod_{I_1} (u_i - u)^{2\rho'} \prod_{I_2} (u_m - u)^{2\rho'} \prod_{I_3} (u_j - u)^{2\rho'} (u-x) \prod_{J_1, J_2, J_3} (v_\alpha - u)^{-2} \quad (5.119)$$

The u part in $J(r_1 - 1, s_1, r_2 + 1, s_2, r_3, s_3)$ would be

$$\int_z^1 du u^{a'} (1-u)^{b'} (u-z)^{c'} \prod_{I_1} (u - u_i)^{2\rho'} \prod_{I_2} (u - u_m)^{2\rho'} \prod_{I_3} (u_j - u)^{2\rho'} (x-u) \prod_{J_1, J_2, J_3} (v_\alpha - u)^{-2} \quad (5.120)$$

Thus we find

$$\phi_1 = e^{-i\pi c'} e^{-i\pi 2\rho'(r_1-1)} (-1)^2 \quad (5.121)$$

The u part in $J(r_1 - 1, s_1, r_2, s_2, r_3 + 1, s_3)$ would be

$$\int_1^\infty du u^{a'} (u-1)^{b'} (u-z)^{c'} \prod_{I_1} (u - u_i)^{2\rho'} \prod_{I_2} (u - u_m)^{2\rho'} \prod_{I_3} (u_j - u)^{2\rho'} (u-x) \prod_{J_1, J_2, J_3} (v_\alpha - u)^{-2} \quad (5.122)$$

so we find

$$\phi_2 = -e^{-i\pi(b'+c'+2\rho'(r_1-1)+2\rho'r_2)} \quad (5.123)$$

We then transform the lowest u contour similarly. Before transforming we have for the u part of $J(r_1, s_1, r_2, s_2, r_3, s_3)$

$$\int_0^z du u^{a'} (1-u)^{b'} (z-u)^{c'} \prod_{I_1} (u - u_i)^{2\rho'} \prod_{I_2} (u_m - u)^{2\rho'} \prod_{I_3} (u_j - u)^{2\rho'} \prod_{J_1, J_2, J_3} (v_\alpha - u)^{-2} (u-x) \quad (5.124)$$

We want to find ψ_1, ψ_2 such that

$$J(r_1, s_1, r_2, s_2, r_3, s_3) = \psi_1 J(r_1 - 1, s_1, r_2 + 1, s_2, r_3, s_3) + \psi_2 J(r_1 - 1, s_1, r_2, s_2, r_3 + 1, s_3) + \dots \quad (5.125)$$

The u part of $J(r_1 - 1, s_1, r_2 + 1, s_2, r_3, s_3)$ would be

$$\int_z^1 du u^{a'} (1-u)^{b'} (u-z)^{c'} \prod_{I_1} (u - u_i)^{2\rho'} \prod_{I_2} (u - u_m)^{2\rho'} \prod_{I_3} (u_j - u)^{2\rho'} \prod_{J_1, J_2, J_3} (v_\alpha - u)^{-2} (x-u) \quad (5.126)$$

hence

$$\psi_1 = (-1)^2 e^{i\pi(c' + 2\rho' r_2)} \quad (5.127)$$

The u part of $J(r_1 - 1, s_1, r_2, s_2, r_3 + 1, s_3)$ would be

$$\int_1^\infty du u^{a'} (u-1)^{b'} (u-z)^{c'} \prod_{I_1} (u-u_i)^{2\rho'} \prod_{I_2} (u-u_m)^{2\rho'} \prod_{I_3} (u-u_j)^{2\rho'} \prod_{J_1, J_2, J_3} (v_\alpha - u)^{-2} (u-x) \quad (5.128)$$

so we find

$$\psi_2 = -e^{i\pi(b' + c' + 2\rho' r_2 + 2\rho' r_3)} \quad (5.129)$$

Finally, we have

$$(\phi_2^* - \psi_2^*) J(r_1, s_1, r_2, s_2, r_3, s_3) = (\phi_1 \phi_2^* - \psi_1 \psi_2^*) J(r_1 - 1, s_1, r_2 + 1, s_2, r_3, s_3) + \dots \quad (5.130)$$

which gives the promised identity.

After several further but in principle straightforward calculations, we obtain

$$\begin{aligned} \alpha_{(r,s,0),(R,S,0)}^{(R,S)} &= (-1)^S e^{i\pi S \rho} \alpha_{r,R}^{(R)}(a', b', c'; \rho') \alpha_{s,S}^{(S)}(a, b, c; \rho) \\ \alpha_{s,S}^{(S)}(a, b, c; \rho) &= \frac{\prod_{j=1}^S s(j\rho)}{\prod_{k=1}^{S-s} s(k\rho) \prod_{m=1}^s s(m\rho)} \\ &\quad \cdot \prod_{j=0}^{s-1} \frac{s(b + \rho(S - s + j))}{s(b + c + \rho(S + j - 1))} \prod_{l=0}^{S-s-1} \frac{s(c + \rho(s + l))}{s(b + c + \rho(s + S + l - 1))} \end{aligned} \quad (5.131)$$

and where $\alpha_{r,R}^{(R)}(a', b', c'; \rho')$ is given by a completely similar expression. The phase is the result of multiplying many phases together. This completes the calculation of the matrix elements of the relevant column as far as fusion rule I is concerned. The result has a form identical to what is found for minimal models [42].

Concerning fusion rule II it turns out that a simple trick allows to obtain the result rather easily. In fact, a suitable contour deformation of the complex contour $\mathcal{C}_v$ allows one to obtain an equation of the form

$$\mathcal{S}_{(r,s,1)}^{(R,S)} = -\frac{1}{\pi} \left(e^{i\pi \rho s} s(c + \rho s) \mathcal{S}_{(r,s+1,0)}^{(R,S)} + s(a + c + 2\rho s) \int_x^0 \dots \right) \quad (5.132)$$

The integral from x to 0 (we imagine $x < 0$ in the s -channel) cannot have a contribution with a $(1-z)$ singularity appropriate for $\mathcal{T}_{(R,S,0)}^{(R,S)}$, and so we do not specify the integrand, and we drop the integral in the calculation. Now it is an easy matter to obtain the missing matrix elements from the ones we have already given. One finds

$$\alpha_{(r,s,1),(R,S,0)}^{(R,S)} = -\frac{1}{\pi} \frac{s(c + \rho s) s((s+1)\rho)}{s(\rho)} \alpha_{(r,s+1,0),(R,S,0)}^{(R,S)} \quad (5.133)$$

where

$$s = 0, \dots, S-1 \quad (5.134)$$

5.5.2 The Row of the Transformation Matrix

The procedure is to consider the s-channel block $S_{(R,S,0)}^{(R,S)}$ and then isolate the t-channel singularities in $(1-z)$ and $(x-1)$. The strengths of these singularities will tell us which t-channel block is obtained. In this way we determine modified crossing matrix elements

$$\begin{aligned} S_{(r,s,\epsilon)}^{(R,S)}(z, x) &= \sum_{r'=0}^R \sum_{s'=0}^S \alpha_{(r,s,\epsilon),(r',s',0)}^{(R,S)'} t_{(r',s',0)}^{(R,S)}(z, x) \\ &+ \sum_{r'=0}^R \sum_{s'=0}^{S-1} \alpha_{(r,s,\epsilon),(r',s',1)}^{(R,S)'} t_{(r',s',1)}^{(R,S)}(z, x) \end{aligned} \quad (5.135)$$

These matrix elements are related to the ones we have previously considered by the normalisation constants of the last section. Denoting the corresponding normalisations in the t-channel by $\tilde{N}_{(r,s,0)}^{(R,S)}(a, b, c; \rho)$, we have

$$\begin{aligned} \alpha_{(R,S,0),(r,s,0)}^{(R,S)} &= \alpha_{(R,S,0),(r,s,0)}^{(R,S)'} / \tilde{N}_{(r,s,0)}^{(R,S)}(a, b, c; \rho) \\ &= \alpha_{(R,S,0),(r,s,0)}^{(R,S)'} / N_{(r,s,0)}^{(R,S)}(b, a, c; \rho) \end{aligned} \quad (5.136)$$

We consider the real time ordered form of the integral representation

$$\begin{aligned} &e^{i\pi(R-S\rho)} S_{(R,S,0)}^{(R,S)} \\ &= z^{2j_1 j_2 / t} (1-z)^{2j_2 j_3 / t} \\ &\cdot z^P T \int_0^1 \prod_{I=1}^R du_I \prod_{K=1}^S dv_K u_I^{a'} (1-zu_I)^{b'} (1-u_I)^{c'} (x-zu_I) \prod_{I < I'} (u_I - u_{I'})^{2\rho'} \\ &\cdot v_K^a (1-zv_K)^b (1-v_K)^c (x-zv_K)^{-\rho} \prod_{K < K'} (v_K - v_{K'})^{2\rho} \prod_{I,K} (u_I - v_K)^{-2} \end{aligned} \quad (5.137)$$

The phase on the left hand side takes into account that s- and t-channel blocks are defined with different phase conventions as far as the factors $(x-u)$ and $(x-v)^{-\rho}$ are concerned. The pre-factor z^P is obtained from scaling the integration variables with z . When $z \rightarrow 1$ it is regular and we shall ignore it in the following.

Next, we use analytic tricks similar to those used when discussing the integral identity, in order to isolate the singularities in $(1-z)$ and $(x-1)$. We consider the integration region, where the first r u_i 's are integrated (ordered) from $1-\epsilon$ to 1, (ϵ small > 0) the first s v_k 's similarly from $(1-\epsilon)$ to 1, and the remaining variables from 0 to $1-\epsilon$. The first u_i 's and v_k 's are indexed by i and k and the remaining ones by j, l . The first u_i 's are transformed as $u_i \rightarrow 1-u_i$, followed by $u_i \rightarrow (1-z)u_i$, and likewise $v_k \rightarrow 1-v_k$ followed by $v_k \rightarrow (1-z)v_k$. Inserting that in (5.137) we find the singular behaviour as $z \rightarrow 1$

$$(1-z)^{-h(j_2)-h(j_3)+h(j_I)} \quad (5.138)$$

where

$$j_I = j_2 + j_3 - r + st \quad (5.139)$$

Furthermore one isolates the $x \rightarrow 1$ behaviour

$$(x-1)^{r-st} = (x-1)^{j_2+j_3-j_I} \quad (5.140)$$

Therefore we may calculate the coefficient of $t_{(r,s,0)}^{(R,S)}$ in the expansion of $S_{(R,S,0)}^{(R,S)}$. After some work one finds the result

$$\begin{aligned} \alpha_{(R,S,0),(r,s,0)}^{(R,S)'} &= e^{i\pi(-R+S\rho)} \mathcal{J}_{r,s}(-b-c+2(r-1-(s-1)\rho), c; \rho) \\ &\cdot \mathcal{J}_{R-r, S-s}(b+c-\rho-2(r-\rho s), a; \rho) \end{aligned} \quad (5.141)$$

Some further calculations give

$$\begin{aligned} \alpha_{(R,S,0),(r,s,0)}^{(R,S)} &= (-1)^S e^{i\pi S\rho} \alpha_{R,r}^{(R)}(a', b', c', d'; \rho') \alpha_{S,s}^{(S)}(a, b, c, d; \rho) \\ \alpha_{S,s}^{(S)}(a, b, c, d; \rho) &= \prod_{i=0}^{s-1} \frac{s(b+i\rho)}{s(b+c+(s-1+i)\rho)} \\ &\cdot \prod_{i=0}^{S-s-1} \frac{s(a+b+c+d+2(S-1)\rho-i\rho)}{s(b+c+d+2(S-1)\rho-(S-s-1+i)\rho)} \end{aligned} \quad (5.142)$$

For convenience of writing we have defined

$$\begin{aligned} d &= -\rho \\ d' &= -d/\rho = 1 \end{aligned} \quad (5.143)$$

The presence of the d dependence is the only difference from the corresponding expression in minimal models [42]. It originates directly from the factors $(u-x)$ and $(v-x)^{-\rho}$ in the integral realization. Such factors are not present in the case of minimal models.

In order to isolate the singularity which corresponds to the t-channel blocks for fusion rule II, we supplement the above specification of the integration region by the requirement that the variable v_{s+1} should be integrated between $1-\epsilon$ and 1, and then transformed as $v_{s+1} \rightarrow 1-v_{s+1}$ followed by $v_{s+1} \rightarrow (x-1)v_{s+1}$. After some calculations we find

$$\begin{aligned} \alpha_{(R,S,0),(r,s,1)}^{(R,S)'} &= e^{i\pi(-R+S\rho)} \mathcal{J}_{r,s}(-b-c+2(r-s\rho)+2(\rho-1), c; \rho) \\ &\cdot \mathcal{J}_{R-r, S-s-1}(a, b+c+\rho-2(r-\rho s); \rho) \\ &\cdot \frac{\Gamma(\rho-1-b-c-2\rho s+2r)\Gamma(b+c+2\rho s-2r+1)}{\Gamma(\rho)} \end{aligned} \quad (5.144)$$

and using the normalisations and various gamma-function identities

$$\alpha_{(R,S,0),(r,s,1)}^{(R,S)} = \pi \alpha_{(R,S,0),(r,s+1,0)}^{(R,S)} \frac{s(\rho)}{s(b+\rho s)s(b+c+\rho(s-1))} \quad (5.145)$$

5.6 Monodromy Invariant Greens Functions

Following the discussion in [42], monodromy invariant 4-point Greens functions

$$G_{j_1, j_2, j_3, j_4}(z, \bar{z}, x, \bar{x}) \quad (5.146)$$

can be obtained by writing

$$G_{j_1, j_2, j_3, j_4}(z, \bar{z}, x, \bar{x}) = \sum_{r,s,\epsilon} |S_{(r,s,\epsilon)}^{(R,S)}(j_1, j_2, j_3, j_4; z, x)|^2 X_{(r,s,\epsilon)}^{(R,S)} \quad (5.147)$$

This form ensures single valuedness in the limits $z \rightarrow 0$ and $x \rightarrow 0$. Single valuedness in the limits $z \rightarrow 1$ and $x \rightarrow 1$ is ensured provided the X 's are chosen to satisfy [42]

$$X_{(r,s,\epsilon)}^{(R,S)} \propto \frac{\alpha_{(R,S,0),(r,s,\epsilon)}^{(R,S)}(b, a, c, d; \rho)}{\alpha_{(r,s,\epsilon),(R,S,0)}^{(R,S)}(a, b, c; \rho)} \quad (5.148)$$

Using re-scaling tricks similar to [42] we may throw away terms (products of sine-functions) independent of r and s . In this case we obtain [112]

$$\begin{aligned} X_{(r,s,0)}^{(R,S)} &= X_r^{(R)}(a', b', c', d'; \rho') X_s^{(S)}(a, b, c, d; \rho) \\ X_s^{(S)}(a, b, c, d; \rho) &= \prod_{i=1}^s s(i\rho) \prod_{i=1}^{S-s} s(i\rho) \prod_{i=0}^{s-1} \frac{s(a+i\rho)s(c+i\rho)}{s(a+c+(s-1+i)\rho)} \\ &\quad \cdot \prod_{i=0}^{S-s-1} \frac{s(b+i\rho)s(1-a-b-c-d-2(S-1)\rho+i\rho)}{s(1-a-c-d-2(S-1)\rho+(S-s-1+i)\rho)} \end{aligned} \quad (5.149)$$

where $X_r^{(R)}(a', b', c', d'; \rho')$ is given by a quite similar expression. Let us illustrate the re-scaling technique by demonstrating the appearance of $\prod_{i=0}^{S-s-1} s(b+i\rho)$. From the ratio between the α 's in (5.148) we find that the only term involving only b is

$$\begin{aligned} \prod_0^{s-1} \frac{1}{s(b+\rho(S-s+j))} &= \prod_{S-s}^{S-1} \frac{1}{s(b+\rho j)} \\ &= \prod_0^{S-1} \frac{1}{s(b+\rho j)} \prod_0^{S-s-1} s(b+\rho j) \end{aligned} \quad (5.150)$$

The factor with the product from 0 to $S-1$ may be scaled away and we are left with the aforementioned term. Generally, it is allowed to renormalise the coefficients by arbitrary functions of R, S and a, b, c, d .

The expression (5.149) is very similar to the result for minimal models except for the presence of the $d = -\rho$ and $d' = 1$. Finally

$$X_{(r,s,1)}^{(R,S)} = -\pi^2 \frac{s^2(\rho)}{s(c+\rho s)s((s+1)\rho)s(a+\rho s)s(a+c+\rho(s-1))} X_{(r,s+1,0)}^{(R,S)} \quad (5.151)$$

Consistency requires that the Greens functions thus defined automatically are single valued also around $z \rightarrow x$. It has been checked that indeed for one screening charge of the second kind this is the case [112]. It is expected to be true generally.

It is convenient for studies of the operator algebra coefficients to also introduce the following expansion

$$G_{j_1, j_2, j_3, j_4}(z, \bar{z}, x, \bar{x}) = \sum_{r,s,\epsilon} |s_{(r,s,\epsilon)}^{(R,S)}(j_1, j_2, j_3, j_4; z, x)|^2 f_{(r,s,\epsilon)}^{(R,S)} \quad (5.152)$$

where $s_{(r,s,\epsilon)}^{(R,S)}$ are defined in (5.56) and (5.58). The coefficients $f_{(r,s,\epsilon)}^{(R,S)}$ differ from the coefficients $X_{(r,s,\epsilon)}^{(R,S)}$ by a factor $(N_{(r,s,\epsilon)}^{(R,S)})^2$. It is fairly straightforward to collect all the results and obtain the

expressions for $f_{(r,s,\epsilon)}^{(R,S)}$. On the way we use the following symmetrised version of (4.58), which is proven using the identity (2.170),

$$\begin{aligned}
\mathcal{J}_{nm}(a, b; \rho) &= \rho^{2nm} \prod_{i,j=1}^{n,m} \frac{1}{(-i+j\rho)} \prod_{i=1}^n \frac{\Gamma(i\rho')}{\Gamma(\rho')} \prod_{i=1}^m \frac{\Gamma(i\rho)}{\Gamma(\rho)} \\
&\cdot \prod_{i,j=0}^{n-1,m-1} \frac{1}{(a+j\rho-i)(b+j\rho-i)(a+b+(m-1+j)\rho-(n-1+i))} \\
&\cdot \prod_{i=0}^{n-1} \frac{\Gamma(1+a'+i\rho')\Gamma(1+b'+i\rho')}{\Gamma(2-2m+a'+b'+(n-1+i)\rho')} \\
&\cdot \prod_{i=0}^{m-1} \frac{\Gamma(1+a+i\rho)\Gamma(1+b+i\rho)}{\Gamma(2-2n+a+b+(m-1+i)\rho)} \tag{5.153}
\end{aligned}$$

We then obtain

$$\begin{aligned}
f_{(r,s,0)}^{(R,S)} &= \Lambda_{r,s}^{(R,S)}(\rho) \\
&\cdot \prod_{i,j=0}^{r-1,s-1} \frac{1}{(a+j\rho-i)^2(c+j\rho-i)^2(a+c+\rho(s-1+j)-(r-1+i))^2} \\
&\cdot \prod_{i,j=0}^{R-r-1,S-s-1} \frac{1}{(b-i+j\rho)^2(e-i+j\rho)^2(e+b-(R-r-1+i)+(S-s-1+j)\rho)^2} \\
&\cdot \prod_{i=0}^{r-1} \frac{G(1+a'+i\rho')G(1+c'+i\rho')}{G(2-2s+a'+c'+(r-1+i)\rho')} \prod_{i=0}^{s-1} \frac{G(1+a+i\rho)G(1+c+i\rho)}{G(2-2r+a+c+(s-1+i)\rho)} \\
&\cdot \prod_{i=0}^{R-r-1} \frac{G(1+b'+i\rho)G(1+e'+i\rho)}{G(2+e'+b'-2(S-s)+(R-r-1+i)\rho')} \\
&\cdot \prod_{i=0}^{S-s-1} \frac{G(1+b+i\rho)G(1+e+i\rho)}{G(2+e+b-2(R-r)+(S-s-1+i)\rho)} \tag{5.154}
\end{aligned}$$

where we have defined

$$G(x) = \frac{\Gamma(x)}{\Gamma(1-x)} = \frac{1}{G(1-x)} \tag{5.155}$$

and where

$$\begin{aligned}
\Lambda_{r,s}^{(R,S)}(\rho) &= \rho^{4rs+4(R-r)(S-s)} \prod_{i=1}^s G(i\rho) \prod_{i=1}^{S-s} G(i\rho) \prod_{i=1}^r G(i\rho') \prod_{i=1}^{R-r} G(i\rho') \\
&\cdot \prod_{i,j=1}^{r,s} \frac{1}{(i-j\rho)^2} \prod_{i,j=1}^{R-r,S-s} \frac{1}{(i-j\rho)^2} \tag{5.156}
\end{aligned}$$

and where we have defined

$$\begin{aligned}
e &= -a - b - c - d - 2\rho(S-1) + 2(R-1) \\
e' &= -e/\rho \tag{5.157}
\end{aligned}$$

These expressions are like the ones for minimal models except for the appearance of the terms d, d' in the definition of e, e' . Finally we have

$$f_{(r,s,1)}^{(R,S)} = f_{(r,s+1,0)}^{(R,S)} \cdot \frac{G(2+a+c+2s\rho-2r)G(1+a+c+2s\rho-2r)G(1-(s+1)\rho+r)}{G(1-r+a+s\rho)G(1-r+c+s\rho)G(2-r+a+c+(s-1)\rho)} \quad (5.158)$$

The following useful identity for the G -functions defined above will be used repeatedly in the sequel

$$G(x) = (-1)^n G(x-n) \prod_{i=1}^n (x-i)^2 \quad (5.159)$$

5.6.1 Case of Mixed Fusion Rules

The construction of crossing matrices and monodromy coefficients in the case of mixed fusion rules (or over-screening) is made essentially trivial by the following observations. We have previously seen that the integral for the DF contours Fig. 5.8 is related to a corresponding integral for time ordered integrations by the factor

$$\lambda_r(1/t)\lambda_s(t)\lambda_{R-r}(1/t)\lambda_{S-s}(t) \quad (5.160)$$

Similarly for the present case of some contours being of Felder type, we get instead a factor

$$\lambda_r(1/t)\chi_s^{(2)}(S-s_3-1;t)\lambda_{R-r}(1/t)\lambda_{S-s}(t) \quad (5.161)$$

for (I,II), and a factor

$$\lambda_r(1/t)\lambda_s(t)\lambda_{R-r}(1/t)\chi_{S-s}^{(2)}(S-s_2-1;t) \quad (5.162)$$

for (II,I). These rather trivial new normalisations allow us to follow completely the treatment for fusion rule I (i.e. the case (I,I)) described above and insert appropriate $\frac{\chi}{\lambda}$ factors as normalisations. It is fairly easy to see, that for the new monodromy coefficients $f_{(r,s,(I,II))}^{(R,S)}$ and $f_{(r,s,(II,I))}^{(R,S)}$ (in a self explanatory notation), the only $\frac{\chi}{\lambda}$ factors multiplying $f_{(r,s,(I,I))}^{(R,S)}$ which survive, are ones which do not depend on (r, s) or $(R-r, S-s)$, but only on (R, S) and hence may be absorbed into renormalisations.

5.7 Operator Algebra Coefficients

All properties of a CFT are encoded in the knowledge of the symmetry algebra $W \oplus \bar{W}$ (W is an extension of the Virasoro algebra Vir , here a semi-direct sum of $sl(2)$ and Vir), the spectrum of primary fields and the operator algebra (OA) coefficients or structure constants. From the analysis in the preceding sections we are now in a position to determine the OA coefficients and thereby 'solve' the theory.

In the $SL(2)$ theory the operator algebra coefficients $C_{\lambda_1 \lambda_2}^{\lambda_3}$ are defined by

$$\phi_{j_2}(z, \bar{z}, x, \bar{x})\phi_{j_1}(0, 0, 0, 0) = \sum_j \frac{|x|^{2(j_1+j_2-j)}}{|z|^{2(\Delta(j_1)+\Delta(j_2)-\Delta(j))}} C_{\lambda_1 \lambda_2}^{\lambda} \phi_j(0, 0, 0, 0) + \dots \quad (5.163)$$

where the dots represent contributions from descendants and where $\lambda = 2j + 1$. The OA coefficients are only defined up to normalisation of the fields

$$\begin{aligned}\phi_{j_i}(z, \bar{z}, x, \bar{x}) &\rightarrow \rho_{\lambda_i} \phi_{j_i}(z, \bar{z}, x, \bar{x}), & \rho_1 = 1 \\ C_{\lambda_1 \lambda_2}^{\lambda_3} &\rightarrow \rho_{\lambda_1} \rho_{\lambda_2} \rho_{\lambda_3}^{-1} C_{\lambda_1 \lambda_2}^{\lambda_3}\end{aligned}\quad (5.164)$$

Define the conjugate field $\phi^j(z, \bar{z}, x, \bar{x})$ by

$$\langle 0 | \phi^{j_i}(z, \bar{z}, x, \bar{x}) \phi_{j_j}(0, 0, 0, 0) | 0 \rangle = \delta_{\lambda_j}^{\lambda_i} |z|^{-4\Delta(j_i)} |x|^{4j_i} \quad (5.165)$$

Then the OPE coefficients can be calculated by considering the 3-point function

$$\begin{aligned}&\langle 0 | \phi^{j_3}(z_3, \bar{z}_3, x_3, \bar{x}_3) \phi_{j_2}(z_2, \bar{z}_2, x_2, \bar{x}_2) \phi_{j_1}(z_1, \bar{z}_1, x_1, \bar{x}_1) | 0 \rangle \\ &= C_{\lambda_1 \lambda_2}^{\lambda_3} |z_1 - z_2|^{2(\Delta_3 - \Delta_1 - \Delta_2)} |z_2 - z_3|^{2(\Delta_1 - \Delta_2 - \Delta_3)} |z_1 - z_3|^{2(\Delta_2 - \Delta_1 - \Delta_3)} \\ &\cdot |x_1 - x_2|^{2(j_1 + j_2 - j_3)} |x_2 - z_3|^{2(j_2 + j_3 - j_1)} |x_1 - x_3|^{2(j_1 + j_3 - j_2)}\end{aligned}\quad (5.166)$$

Defining

$$\overline{\phi^j}(z, \bar{z}, x, \bar{x}) = \phi^j(-1/z, -1/\bar{z}, -1/x, -1/\bar{x}) |z|^{4\Delta_j} |x|^{-4j} \quad (5.167)$$

we see that

$$C_{\lambda_1 \lambda_2}^{\lambda_3} = \langle 0 | \overline{\phi^{j_3}}(0, 0, 0, 0) \phi_{j_2}(1, 1, 1, 1) \phi_{j_1}(0, 0, 0, 0) | 0 \rangle \quad (5.168)$$

Consider now

$$\langle 0 | \phi_{j_i}(z, \bar{z}, x, \bar{x}) \phi_{j_j}(0, 0, 0, 0) | 0 \rangle = C_{\lambda_i \lambda_j}^1 |z|^{-4\Delta(j_i)} |x|^{4j_i} \quad (5.169)$$

In the $SL(2)$ theory the fields are self-conjugate so $C_{\lambda_i \lambda_j}^1$ is not only symmetric in λ_i, λ_j but also diagonal, and it may be used as a metric, raising and lowering indices

$$\begin{aligned}C_{\lambda_i \lambda_j}^1 &= c_{\lambda_i} \delta_{\lambda_i \lambda_j} = g_{\lambda_i \lambda_j} \\ C_{\lambda_i \lambda_j}^{\lambda_l} &= g^{\lambda_l \lambda_k} C_{\lambda_i \lambda_j \lambda_k} = c_{\lambda_l}^{-1} C_{\lambda_i \lambda_j \lambda_l}\end{aligned}\quad (5.170)$$

Here $C_{\lambda_i \lambda_j \lambda_k}$ is totally symmetric in the λ_i 's, and c_λ is a normalisation constant. We also have

$$\begin{aligned}\phi^{j_i}(z, \bar{z}, x, \bar{x}) &= g^{\lambda_i \lambda_j} \phi_{j_j}(z, \bar{z}, x, \bar{x}) \\ &= c_{\lambda_i}^{-1} \phi_{j_i}(z, \bar{z}, x, \bar{x})\end{aligned}\quad (5.171)$$

One can fix the normalisation constant $\hat{c}_\lambda = 1$ in a basis where $\hat{\phi}_j = \rho_\lambda^{-1} \phi_j$, such that the fields are orthonormal

$$\hat{C}_{\lambda_1 \lambda_2}^1 = \delta_{\lambda_1 \lambda_2} \quad (5.172)$$

Let the totally symmetric OA coefficients in the orthonormal basis be denoted as $\hat{C}_{\lambda_i \lambda_j \lambda_l}$, then in the other basis the OA coefficients $C_{\lambda_i \lambda_j}^{\lambda_k}$ can be written as

$$\begin{aligned}C_{\lambda_i \lambda_j}^{\lambda_k} &= \rho_{\lambda_i} \rho_{\lambda_j} \rho_{\lambda_k}^{-1} \hat{C}_{\lambda_i \lambda_j \lambda_k} \\ \rho_\lambda^2 &= c_\lambda\end{aligned}\quad (5.173)$$

Here the normalisation constants correspond to the original basis without 'hats'.

5.7.1 Case of Fusion Rule I

Since there is no overlap between the two sets of fusion rules, a certain 4-point Greens function

$$G_{j_1 j_2 j_3 j_4}(z, \bar{z}, x, \bar{x}) = \langle j_4 | \phi_{j_3}(1, 1, 1, 1) \phi_{j_2}(z, \bar{z}, x, \bar{x}) | j_1 \rangle \quad (5.174)$$

behaves, in the limit $z, x \rightarrow 0$, either like

$$\begin{aligned} G_{j_1 j_2 j_3 j_4}(z, \bar{z}, x, \bar{x}) &= \sum_j C_{\lambda_3 \lambda}^{\lambda_4} C_{\lambda_1 \lambda_2}^{\lambda} |z|^{2(-\Delta_{j_1} - \Delta_{j_2} + \Delta_j)} |x|^{2(j_1 + j_2 - j)} (1 + \mathcal{O}(z, \bar{z}, x, \bar{x})) \\ &= \sum_j C_{\lambda_3 \lambda}^{\lambda_4}(I) C_{\lambda_1 \lambda_2}^{\lambda}(I) |z|^{2(-\Delta_{j_1} - \Delta_{j_2} + \Delta_j)} |x|^{2(j_1 + j_2 - j)} (1 + \mathcal{O}(z, \bar{z}, x, \bar{x})) \\ &+ \sum_j C_{\lambda_3 \lambda}^{\lambda_4}(II) C_{\lambda_1 \lambda_2}^{\lambda}(II) |z|^{2(-\Delta_{j_1} - \Delta_{j_2} + \Delta_j)} |x|^{2(j_1 + j_2 - j)} (1 + \mathcal{O}(z, \bar{z}, x, \bar{x})) \end{aligned} \quad (5.175)$$

or like

$$\begin{aligned} G_{j_1 j_2 j_3 j_4}(z, \bar{z}, x, \bar{x}) &= \sum_j C_{\lambda_3 \lambda}^{\lambda_4}(I) C_{\lambda_1 \lambda_2}^{\lambda}(II) |z|^{2(-\Delta_{j_1} - \Delta_{j_2} + \Delta_j)} |x|^{2(j_1 + j_2 - j)} (1 + \mathcal{O}(z, \bar{z}, x, \bar{x})) \\ &+ \sum_j C_{\lambda_3 \lambda}^{\lambda_4}(II) C_{\lambda_1 \lambda_2}^{\lambda}(I) |z|^{2(-\Delta_{j_1} - \Delta_{j_2} + \Delta_j)} |x|^{2(j_1 + j_2 - j)} (1 + \mathcal{O}(z, \bar{z}, x, \bar{x})) \end{aligned} \quad (5.176)$$

depending on the 4-tuple (j_1, j_2, j_3, j_4) . First we concentrate on the former case where

$$f_{(r,s,\epsilon)}^{(R,S)}(j_1, j_2, j_3, j_4) = C_{\lambda_3 \lambda_\epsilon}^{\lambda_4} C_{\lambda_1 \lambda_2}^{\lambda_\epsilon} \quad (5.177)$$

with $\lambda_\epsilon = \lambda_I, \lambda_{II}$ for $\epsilon = 0, 1$. Such a split requires that the monodromy coefficients are properly normalised. The normalisation adopted so far (5.154) (and thereby (5.158)) follows the prescription of Dotsenko and Fateev in the case of minimal models, but turns out to be inadequate here [112]. Indeed it is completely essential that the above factorisation takes place in such a way that the operator algebra coefficients only depend on the variables indicated and not on anything else. In particular, $C_{\lambda_1 \lambda_2}^{\lambda_\epsilon}$ is allowed to depend on r, s which are given in terms of the spins (the λ 's) indicated, but it is not allowed to depend on R, S for example. Likewise $C_{\lambda_3 \lambda_\epsilon}^{\lambda_4}$ is allowed to depend on $R - r, S - s$ but again, not on R, S . However, it is allowed (as utilised above) to renormalise the coefficients by arbitrary functions of $R, S, \lambda_1, \lambda_2, \lambda_3$, ($j_4 = j_1 + j_2 + j_3 - R + St$). It turns out [112] to be possible to devise such a normalisation with the above criterion satisfied. This we first do for fusion rule I. We have to use (cf. (5.52))

$$\begin{aligned} a &= -\lambda_3 + R - St + t \\ b &= -\lambda_1 + R - St + t \\ e &= -\lambda_2 + R - St + t \\ c &= \lambda_4 + R - St \\ e + b &= -\lambda_I + 2(R - r) - 1 - 2t(S - s - 1) = +\lambda_{II} + 2(R - r) - 1 - 2t(S - s - 1) \\ a + c &= \lambda_I + 2r - 2st - 1 + t = -\lambda_{II} + 2r - 2st - 1 + t \end{aligned} \quad (5.178)$$

Then we may express the monodromy coefficient as

$$\begin{aligned}
f_{(r,s,0)}^{(R,S)} &= C_{\lambda_1 \lambda_2}^{\lambda_I}(I) C_{\lambda_3 \lambda_I}^{\lambda_4}(I) \\
&= t^{4rs} \prod_1^s G(it - r) \prod_1^r G(i/t) \\
&\cdot \prod_0^{r-1} \frac{G(S + (\lambda_3 - R + j)/t) G(S + 1 + (-\lambda_4 - R + j)/t)}{G(1 - (\lambda_I + 1 + j)/t)} \\
&\cdot \prod_0^{s-1} \frac{G(1 - \lambda_3 + R - r - (S - s + j)t) G(1 + \lambda_4 + R - r - (S - s + 1 - j)t)}{G(1 + \lambda_I + r - (s - j)t)} \\
&\cdot t^{4(R-r)(S-s)} \prod_1^{S-s} G(it - R + r) \prod_1^{R-r} G(i/t) \\
&\cdot \prod_0^{R-r-1} \frac{G(S + (\lambda_1 - R + i)/t) G(S + (\lambda_2 - R + i)/t)}{G((\lambda_I - 1 - i)/t)} \\
&\cdot \prod_0^{S-s-1} \frac{G(1 - \lambda_1 + r - (s + i)t) G(1 - \lambda_2 + r - (s + i)t)}{G(1 - \lambda_I + R - r - (S - s - 1 - i)t)} \tag{5.179}
\end{aligned}$$

This is certainly not in the form required, but we may renormalise it. Thus we consider terms involving only λ_1

$$\begin{aligned}
&\prod_{i=0}^{R-r-1} G(S + (\lambda_1 - R + i)/t) \prod_{j=0}^{S-s-1} G(1 - \lambda_1 + r - (s + j)t) \\
&= t^{2r(S-s)} \prod_{i=0}^{r-1} G(1 - s + (1 - \lambda_1 + i)/t) \prod_{j=0}^{s-1} G(\lambda_1 + jt) \\
&\cdot \prod_{i=0}^{R-1} G(S + (\lambda_1 - R + i)/t) \prod_{j=0}^{S-1} G(1 - \lambda_1 + (1 - S + j)t) \tag{5.180}
\end{aligned}$$

The last line may now be scaled away. Symmetry arguments in the λ_i 's render considerations of terms including $\lambda_2, \lambda_3, \lambda_4$ trivial, while the λ_I part contributes as

$$\begin{aligned}
&\prod_{i=0}^{R-r-1} G^{-1}((\lambda_I - 1 - i)/t) \prod_{j=0}^{S-s-1} G^{-1}(1 - \lambda_I + R - r - (S - s - 1 - j)t) \\
&\cdot \prod_{i=0}^{r-1} G^{-1}(1 - (\lambda_I + 1 + i)/t) \prod_{j=0}^{s-1} G^{-1}(1 + \lambda_I + r - (s - j)t) \\
&= t^{-2(R-r)(S-s)-2rs} \prod_{i=0}^{R-r-1} G^{-1}(S - s - (1 - \lambda_I + i)/t) \prod_{j=0}^{S-s-1} G^{-1}(1 - \lambda_I - jt) \\
&\cdot \prod_{i=0}^{r-1} G^{-1}(1 + s - (1 + \lambda_I + i)/t) \prod_{j=0}^{s-1} G^{-1}(1 + \lambda_I - (j + 1)t) \tag{5.181}
\end{aligned}$$

The total power of t becomes

$$t^{4RS-2(R-r)(S-s)-2rs} \tag{5.182}$$

where we may absorb the t^{4RS} in the normalisation. Finally we find

$$\begin{aligned}
C_{\lambda_1 \lambda_2}^\lambda(r, s; I) &= t^{-2rs} \prod_{i=1}^r G(i/t) \prod_{i=1}^s G(it - r) \\
&\cdot \prod_{i=0}^{r-1} \frac{G(1-s+(1-\lambda_1+i)/t)G(1-s+(1-\lambda_2+i)/t)}{G(1+s-(1+\lambda+i)/t)} \\
&\cdot \prod_{i=0}^{s-1} \frac{G(\lambda_1+it)G(\lambda_2+it)}{G(1+\lambda-(1+i)t)} \\
C_{\lambda_3 \lambda}^{\lambda_4}(R-r, S-s; I) &= t^{-2(R-r)(S-s)} \prod_{i=1}^{R-r} G(i/t) \prod_{i=1}^{S-s} G(it - (R-r)) \\
&\cdot \prod_{i=0}^{R-r-1} \frac{G(1-(S-s)+(1-\lambda_3+i)/t)}{G(S-s-(1-\lambda+i)/t)} \\
&\cdot \prod_{i=0}^{R-r-1} G(-(S-s)+(1+\lambda_4+i)/t) \\
&\cdot \prod_{i=0}^{S-s-1} \frac{G(\lambda_3+it)G(-\lambda_4+(1+i)t)}{G(1-\lambda-it)} \tag{5.183}
\end{aligned}$$

and notice the symmetry

$$C_{\lambda_1 \lambda_2}^{\lambda_3}(r, s; I) = C_{\lambda_1, -\lambda_3+t}^{-\lambda_2+t}(r, s; I) \tag{5.184}$$

which reflects the involutory invariance (4.43). Here, for clarity we have indicated the dependencies on r, s or $R-r, S-s$ and on the fusion rule (here I). In fact this is somewhat superfluous. The point is, that from three spins, it is always clear by which of the two fusion rules they couple, and for each case there is a unique possible value of r, s (or $R-r, S-s$). Furthermore, a trivial consistency check reveals that indeed the two expressions in (5.183) are equivalent.

We may now compute the normalisation constants c_λ and ρ_λ for $\lambda = r - st$

$$c_\lambda = t^{-2s} \frac{G(\lambda/t)}{G(1/t)} \quad , \quad \rho_\lambda = t^{-s} \frac{G^{1/2}(\lambda/t)}{G^{1/2}(1/t)} \tag{5.185}$$

Using this we want to find the normalised and symmetric OA coefficient $\hat{C}_{\lambda_1 \lambda_2 \lambda_3}(I)$. Let us introduce the notation

$$\begin{aligned}
\hat{R} &= (r_1 + r_2 + r_3 - 1)/2 \\
\hat{S} &= (s_1 + s_2 + s_3)/2 \\
R_i &= \hat{R} - r_i \\
S_i &= \hat{S} - s_i
\end{aligned} \tag{5.186}$$

where $\lambda_i = 2j_i + 1 = r_i - s_i t$. Employing (5.159) one obtains

$$\hat{C}_{\lambda_1 \lambda_2 \lambda_3}(I) = G^{1/2}(1/t) \prod_{i=1}^{\hat{R}} G(i/t) \prod_{j=1}^{\hat{S}} G(jt - \hat{R})$$

$$\cdot \prod_{k=1}^3 \frac{t^{s_k}}{G^{1/2}(\lambda_k/t)} \frac{\prod_{i=1}^{R_k} G(i/t) \prod_{j=1}^{S_k} G(jt - R_k)}{\prod_{i=1}^{r_k-1} G(i/t) \prod_{j=1}^{s_k} G(jt - r_k + 1)} \quad (5.187)$$

In order to compare with Andreev's result we introduce the functions $P(n, m)$ [8], here expressed in terms of the G -functions (5.155)

$$\begin{aligned} P(n, m) &= \prod_{j=1}^{n-1} \prod_{i=1}^{m-1} (jt - i)^{-2} \prod_{j=1}^{n-1} G(jt) \prod_{i=1}^{m-1} G(i/t) \\ &= (-1)^{(n-1)(m-1)} \prod_{i=1}^{m-1} G(i/t) \prod_{j=1}^{n-1} G(jt - m + 1) \end{aligned} \quad (5.188)$$

whereby we may rewrite (5.187) as

$$\hat{C}_{\lambda_1 \lambda_2 \lambda_3}(I) = t^{2\hat{S}} (-1)^{\hat{S}} G^{1/2}(1/t) P(\hat{S} + 1, \hat{R} + 1) \prod_{k=1}^3 G^{1/2}(1 - \lambda_k/t) \frac{P(S_k + 1, R_k + 1)}{P(s_k + 1, r_k)} \quad (5.189)$$

This is in accordance with Andreev's result. However, so far we have only considered the case of fusion rule I while the case of fusion rule II is somewhat more subtle. Indeed, in [8] Andreev is able to determine the OA coefficients in that case only up to normalisation. Nevertheless, an inspection of 4-point functions with mixed fusion rules determines the normalisation of the fields, after which (5.158) fixes unambiguously the structure coefficients, as will become clear below. It should be stressed that the fixing of the OA coefficients for fusion rule II is only *relative to the choice* of normalisation of the OA coefficients for fusion rule I (5.183).

5.7.2 Case of Fusion Rule II

A priori it is not clear how to normalise the fields in a fusion rule II coupling. This is due to the fact that the identity operator ($j = 0$; $r = 1, s = 0$) is not present according to fusion rule II, hence $C_{\lambda_i \lambda_j}^1(II)$ cannot be used to determine the normalisations. However, when extracting the OA coefficients from a 4-point Greens function with mixed fusion rules (5.176), the fields should have the same normalisations, whether they appear in the fusion rule I coefficient or in the fusion rule II coefficient. Therefore, the normalisation constants in the case of fusion rule II are inherited from the case of fusion rule I and are thus given by the same expressions (5.185).

In the notation (5.186) the numbers of screenings are $R_3 + p/2, S_3 + q/2$, and we have, up to an overall renormalisation constant,

$$\begin{aligned} C_{\lambda_1 \lambda_2}^{\lambda_3}(II) &\propto t^{-2(R_3+p/2)(S_3+q/2)} \prod_{i=1}^{R_3+p/2} G(i/t) \prod_{j=1}^{S_3+q/2} G(jt - (R_3 + p/2)) \\ &\cdot \prod_{i=0}^{R_3+p/2-1} \frac{G(1 - (S_3 + q/2) + (1 - \lambda_1 + i)/t) G(1 - (S_3 + q/2) + (1 - \lambda_2 + i)/t)}{G(1 + S_3 + q/2 - (1 + \lambda_3 + i)/t)} \\ &\cdot \prod_{j=0}^{S_3+q/2-1} \frac{G(\lambda_1 + jt) G(\lambda_2 + jt)}{G(1 + \lambda_3 - (1 + j)t)} \\ &= t^{-4} \frac{G(1 + \lambda_3)}{G(1 + \hat{R} - \hat{S}t)} \prod_{i=1}^{R_3+p/2} G(i/t) \prod_{j=1}^{S_3+q/2} G(jt - R_3 - p/2) \end{aligned}$$

$$\begin{aligned}
& \cdot \frac{\prod_{i=1}^{p/2-1-R_2} G(i/t) \prod_{j=1}^{q/2-1-S_2} G(jt - p/2 + R_2 + 1)}{\prod_{i=1}^{r_1-1} G(i/t) \prod_{j=1}^{s_1} G(jt - r_1 + 1)} \\
& \cdot \frac{\prod_{i=1}^{p/2-1-R_1} G(i/t) \prod_{j=1}^{q/2-1-S_1} G(jt - p/2 + R_1 + 1)}{\prod_{i=1}^{r_2-1} G(i/t) \prod_{j=1}^{s_2} G(jt - r_2 + 1)} \\
& \cdot \frac{\prod_{i=1}^{\hat{R}-p/2} G(i/t) \prod_{j=1}^{\hat{S}-q/2-1} G(jt - \hat{R} + p/2)}{\prod_{i=1}^{p-r_3-1} G(i/t) \prod_{j=1}^{q-s_3} G(jt - p + r_3 + 1)} \quad (5.190)
\end{aligned}$$

where we have used (5.159). In order to determine the proportionality constant, let us further rewrite (5.190). Thus, with the same proportionality constant one finds

$$\begin{aligned}
C_{\lambda_1 \lambda_2}^{\lambda_3}(II) & \propto t^{-4-2(q-s_3)} \frac{G(1+\lambda_3)G(-\lambda_3/t)}{G(1+R_1-S_1t)G(1+R_2-S_2t)} C_{\lambda_1 \lambda_2}^{-\lambda_3}(I) \\
& = t^{-2(q-s_3)} \frac{G(2-\lambda_3/t)G(-R_3+S_3t)}{G(-\hat{R}+(\hat{S}+1)t)G(1+r_3-(1+s_3)t)} C_{\lambda_1 \lambda_2}^{-\lambda_3+2t}(I) \quad (5.191)
\end{aligned}$$

Let us now make a comparison with (5.158)

$$\begin{aligned}
f_{(r,s,1)}^{(R,S)} & = C_{\lambda_3 \lambda_{II}}^{\lambda_4}(II) C_{\lambda_1 \lambda_2}^{\lambda_{II}}(II) \\
& = C_{\lambda_3, -\lambda_4+t}^{-\lambda_{II}+t}(II) C_{\lambda_1 \lambda_2}^{\lambda_{II}}(II) \\
& \propto t^{-2q-6} C_{\lambda_3, -\lambda_4+t}^{\lambda_{II}-t}(I) C_{\lambda_1 \lambda_2}^{-\lambda_{II}+2t}(I) \\
& \cdot \frac{G(t-\lambda_{II})G(1-\lambda_{II}+t)G(-j_1-j_2+j_{II})}{G(-1+t-j_1-j_2-j_{II})G(1+j_3+j_4-j_{II})G(t-j_3-j_4-j_{II})} \\
& = t^{-2q-6} f_{(r,s+1,0)}^{(R,S)} \\
& \cdot \frac{G(1-\lambda_{II}+t)G(-\lambda_{II}+t)G(1-(s+1)t+r)}{G(1-\lambda_3+R-r-(S-s-1)t)G(1+\lambda_4+R-r-(S-s)t)G(1-\lambda_{II}+r-st)} \quad (5.192)
\end{aligned}$$

From this we may read off the proportionality constant in (5.190) to be [112]

$$t^{q+3} \quad (5.193)$$

In conclusion, with the relevant expressions for the numbers of screenings, now denoted r, s

$$r = (r_1 + r_2 - r_3 - 1 + p)/2, \quad s = (s_1 + s_2 - s_3 + q)/2 \quad (5.194)$$

the result (5.183) for $C_{\lambda_1 \lambda_2}^{\lambda_3}$ is valid also for fusion rule II provided one multiplies the rhs by t^{q+3} .

As a consistency check of the result $C_{\lambda_1 \lambda_2}^{\lambda_3}$ in (5.183) one may analyse the products of G -functions in order to verify that up to cancellations of the form $\Gamma(0)/\Gamma(0)$ raised to some power, (5.183) is non-vanishing and well-defined for both sets of fusion rules. It turns out that for fusion rule I no cancellations are needed. For fusion rule II where the extra factor of t^{q+3} is of no importance in this analysis, one finds

$$\Gamma(0) \sim \prod_{i=0}^{r-1} G(1-s+(1-\lambda_1+i)/t) \sim \prod_{i=0}^{r-1} G(1-s+(1-\lambda_2+i)/t)$$

$$\begin{aligned}
\Gamma(0) &\sim \prod_{i=0}^{r-1} \frac{1}{G(1+s-(1+\lambda_3+i)/t)} \\
1/\Gamma(0) &\sim \prod_{i=0}^{s-1} G(\lambda_1+it) \sim \prod_{i=0}^{s-1} G(\lambda_2+it) \sim \prod_{i=0}^{s-1} \frac{1}{G(1+\lambda_3-(1+i)t)}
\end{aligned} \tag{5.195}$$

leaving (5.183) non-vanishing and well-defined upon the cancellation $(\Gamma(0)/\Gamma(0))^3$.

In order to make a direct comparison with Andreev's result we use (5.159) to first obtain the symmetric OA coefficient

$$\begin{aligned}
\hat{C}_{\lambda_1 \lambda_2 \lambda_3}(II) &= t^{(q-1)(4\hat{R}-2p-1)} G^{\frac{1}{2}}(1/t) \prod_{i=1}^{\hat{R}-p/2} G(i/t) \prod_{j=1}^{\hat{S}-q/2} G(jt - \hat{R} + p/2) \\
&\cdot \prod_{k=1}^3 \frac{t^{s_k}}{G^{\frac{1}{2}}(\lambda_k/t)} \frac{\prod_{i=1}^{R_k+p/2} G(i/t) \prod_{j=1}^{S_k+q/2} G(jt - R_k - p/2)}{\prod_{i=1}^{p-r_k} G(i/t) \prod_{j=1}^{q-s_k-1} G(jt - p + r_k)}
\end{aligned} \tag{5.196}$$

where we have included the renormalisation constant t^{q+3} . Furthermore, we should change our notation in (5.186) for λ_3 (the definitions for $\hat{R}$, R_i , $\hat{S}$, S_i and $\lambda_{\{1,2\}}$ remain unchanged)

$$\lambda_3 = p - r_3 - (q - s_3)t = -r_3 + s_3t \tag{5.197}$$

We see that in terms of the present notation, the original one has been transformed according to

$$\begin{aligned}
r_1 &\rightarrow r_1 & , & & s_1 &\rightarrow s_1 \\
r_2 &\rightarrow r_2 & , & & s_2 &\rightarrow s_2 \\
r_3 &\rightarrow p - r_3 & , & & s_3 &\rightarrow q - s_3 \\
R_1 &\rightarrow p/2 - 1 - R_2 & , & & S_1 &\rightarrow q/2 - S_2 \\
R_2 &\rightarrow p/2 - 1 - R_1 & , & & S_2 &\rightarrow q/2 - S_1 \\
R_3 &\rightarrow \hat{R} - p/2 & , & & S_3 &\rightarrow \hat{S} - q/2 \\
\hat{R} &\rightarrow R_3 + p/2 & , & & \hat{S} &\rightarrow S_3 + q/2
\end{aligned} \tag{5.198}$$

In this new notation one can establish the identity

$$\begin{aligned}
\frac{\prod_{i=1}^{R_k} G(i/t) \prod_{j=1}^{S_k-1} G(jt - R_k)}{\prod_{i=1}^{r_k-1} G(i/t) \prod_{j=1}^{s_k} G(jt - r_k + 1)} &= t^{2(q-1)(r_k-1-R_k)} \\
&\cdot \frac{\prod_{i=1}^{p-1-R_k} G(i/t) \prod_{j=1}^{q-S_k} G(jt - p + 1 + R_k)}{\prod_{i=1}^{p-r_k} G(i/t) \prod_{j=1}^{q-s_k-1} G(jt - p + r_k)}
\end{aligned} \tag{5.199}$$

valid only for $k = 1, 2$. Finally one finds

$$\begin{aligned}
\hat{C}_{\lambda_1 \lambda_2 \lambda_3}(II) &= t^{2S-1} \frac{G^{\frac{1}{2}}(\lambda_3/t) G^{\frac{1}{2}}(1/t)}{G^{\frac{1}{2}}(\lambda_1/t) G^{\frac{1}{2}}(\lambda_2/t)} \frac{G(1+\lambda_3)}{G(1+R_3-S_3t)} \\
&\cdot \prod_{i=1}^{\hat{R}} G(i/t) \prod_{j=1}^{\hat{S}} G(jt - \hat{R}) \prod_{k=1}^3 \frac{\prod_{i=1}^{R_k} G(i/t) \prod_{j=1}^{S_k-1} G(jt - R_k)}{\prod_{i=1}^{r_k-1} G(i/t) \prod_{j=1}^{s_k} G(jt - r_k + 1)}
\end{aligned} \tag{5.200}$$

while Andreev's formula may be written (in the notation employed here, and up to a power of t) as

$$C_{\lambda_1 \lambda_2 \lambda_3}(II) \propto G(S_3 t - R_3) \prod_{i=1}^{\hat{R}} G(i/t) \prod_{j=1}^{\hat{S}} G(jt - \hat{R}) \prod_{k=1}^3 \prod_{i=1}^{R_k} G(i/t) \prod_{j=1}^{S_k-1} G(jt - R_k) \quad (5.201)$$

Chapter 6

Conclusions and Outlook

In this thesis we have shown how to deal with fractional powers of free fields by virtue of fractional calculus, and thereby we have elucidated the meaning of the second screening charge proposed by Bershadsky and Ooguri [24] for affine $SL(2)$ WZNW models. The real merit of this is that it has enabled us to use free field realizations to build the most general chiral blocks for such theories on the sphere, even in the case of admissible representations with fractional levels. The ensuing integral formulas are tractable and have allowed us to verify many formal properties, such as projective invariances and the fact that the chiral blocks satisfy the Knizhnik-Zamolodchikov equations. We have presented an explicit and simple proof of the proposal in [64] for how Hamiltonian reduction works at the level of correlators, reducing affine $SL(2)$ current algebra for admissible representations to conformal minimal models. We have further verified that the blocks found indeed reduce in that way. Also the relation to more standard formulations of Hamiltonian reduction has been discussed. Employing the notion of over-screening has allowed us to re-derive the fusion rules on the basis of an analysis of 3-point functions in terms of our free field realization.

The proposal in this thesis for how to treat free ghost fields raised to fractional powers involves a variety of expansions of certain functions, with different monodromies for the individual expansion terms. The physical condition imposed is then that the final expression must respect the known monodromy for the item in consideration. This idea has now found applications in other respects, notably in the operator approach to Liouville theory [130].

In order to generalise a work [42] by Dotsenko and Fateev on minimal models, we have investigated in great detail the 4-point blocks. In particular we have devised integration contours appropriate for suitable bases of blocks, both using our own representation based on free fields and one by Andreev [8] applicable only to 4-point functions. The equivalence of these two representations have been analysed thoroughly at the level of highly non-trivial integral manipulations. As a by-product, we have presented a proof of a remarkable duality in minimal models. In order to compute the monodromy invariant Greens functions we have calculated the crossing matrix. Based on the monodromy coefficients we have isolated the operator algebra coefficients of the theory for both fusion rules. Again the notion of over-screening has been utilised, and in such a way that we have been able to normalise the operator algebra coefficients unambiguously, contrary to the results in [8].

We have presented an explicit (generalised) Wakimoto free field realization of affine Lie algebras by a Gauss decomposition and by determining anomalous (quantum) terms. We

have discussed the screening currents of both kinds generalising the one of the second kind by Bershadsky and Ooguri, and found partial proofs of their existence. Complete proofs were presented in some cases. We have undertaken the study of primary fields in the framework of the isotopic x variable representation and found solutions in special cases of $SL(n)$. In the case of $SL(3)$ we have found the general solution for the primary fields and we have been able to prove the consistency conditions for the existence of the screening currents. This means that we are now in a position to generalise our program in chapters 4 and 5 for $SL(2)$ to $SL(3)$. In principle we should then be able to determine the fusion rules based on 3-point functions, find the 4-point Greens functions and eventually the operator algebra coefficients. We hope to come back to this elsewhere.

Very recently de Boer and Fehér [37] have also obtained an explicit construction of Wakimoto realizations of current algebras, based on quantising a Poisson bracket realization obtained using Hamiltonian reduction. However, a direct comparison is not obvious and their proof is announced to be presented elsewhere. Furthermore, they write down the general result for screening currents of the *first* kind only and claim to have and intend to present a proof of that elsewhere. If this is the case we have then shown in this thesis that screening currents of the *second* kind do exist for $SL(n)$. We believe it should be possible to generalise our existence proofs to the other simple groups. This belief is supported by our very latest developments. Utilising a new and more compact notation to be presented in [113], we hope to be able to carry out such proofs and discuss further general results on the issue of primary fields. Generalisations to supergroups are also currently under investigation.

Some steps have already been taken in the direction of generalising to supergroups. In [29] the representation theory of the affine Lie superalgebra $\widehat{sl}(2|1)$ at fractional level and the free field realization of the corresponding affine current algebra have been discussed.

A future work lies in the computation of correlation functions in the $SL(2)/SL(2)$ approach to non-critical strings where conformal minimal matter is coupled to gravity. Attempts in this direction are found in [9]. Using the results established in this thesis, a possible outline is the following. In the $SL(2)/SL(2)$ approach one deals with a direct sum of 3 copies of affine $SL(2)$ current algebra $\widehat{sl}(2)$ with levels k , $-k-4$ and 4 respectively, $k = p/q - 2$ is fractional and the co-prime integer pair (p, q) parametrises the conformal minimal model. The former two copies correspond to the original $SL(2)$ group and to the gauge group which is also an $SL(2)$ but with shifted (dressed) level. The last copy with level 4 is expressed in terms of a fermionic ghost pair arising in a gauge fixing procedure. Utilising this splitting it should be possible to calculate the correlation functions. However, subtleties from the BRST conditions make the computation non-trivial. When the necessary techniques eventually are developed, a generalisation to higher groups and supergroups seems feasible.

Though not final, yet indeed crucial steps in the direction of formulating non-critical strings in the G/G framework, the work in this thesis presents some new and definite results, notably the completion of the solution of CFT based on affine $SL(2)$ current algebra for admissible representations in terms of free field realizations, the completion of the Wakimoto construction for simple affine current algebras, and discussions on free field realizations of screening currents and primary fields for simple groups.

Bibliography

- [1] E. Abdalla, M.C.B. Abdalla, D. Dalmazi and A. Zadra, *2D-Gravity in Non-Critical Strings*, Lecture Notes in Physics **m20** (Springer, 1994)
- [2] O. Aharony, O. Ganor, J. Sonnenschein, S. Yankielowicz and N. Sochen, Nucl. Phys. **B 399** (1993) 527
- [3] O. Aharony, J. Sonnenschein and S. Yankielowicz, Phys. Lett. **B 289** (1992) 309
- [4] L. Alvarez-Gaumé, C. Gomez and G. Sierra, Phys. Lett. **B 220** (1989) 142;
L. Alvarez-Gaumé, C. Gomez and G. Sierra, Nucl. Phys. **B 319** (1989) 155;
L. Alvarez-Gaumé, C. Gomez and G. Sierra, Nucl. Phys. **B 330** (1990) 347
- [5] L. Alvarez-Gaumé, C. Gomez and G. Sierra, in *Physics and Mathematics of Strings*, Eds. L. Brink, D. Friedan and A.M. Polyakov (World Scientific, 1990)
- [6] K. Amoto, J. Math. Soc. Japan **39** (1987) 191
- [7] G. Anderson and G. Moore, Commun. Math. Phys. **117** (1988) 441
- [8] O. Andreev, Int. J. Mod. Phys. **A 10** (1995) 3221;
O. Andreev, Phys. Lett. **B 363** (1995) 166
- [9] O. Andreev, Phys. Lett. **B 375** (1996) 60
- [10] O. Andreev and B. Feigin, Nucl. Phys. **B 433** (1995) 685
- [11] H. Awata, A. Tsuchiya and Y. Yamada, Nucl. Phys. **B 365** (1991) 680;
H. Awata, Prog. Theor. Phys. Suppl. **110** (1992) 303
- [12] H. Awata and Y. Yamada, Mod. Phys. Lett. **A7** (1992) 1185
- [13] F.A. Bais and P.G. Bouwknegt, Nucl. Phys. **B 279** (1987) 561
- [14] F.A. Bais, P. Bouwknegt, K. Schoutens and M. Surridge, Nucl. Phys. **B 304** (1988) 348;
F.A. Bais, P. Bouwknegt, K. Schoutens and M. Surridge, Nucl. Phys. **B 304** (1988) 371
- [15] J. Balog, L. Fehér, L. O’Raifeartaigh, P. Forgács and A. Wipf, Ann. of Phys. **203** (1990) 76;
L. Fehér, L. O’Raifeartaigh, P. Ruelle, I Tsutsui and A. Wipf, Phys. Rep. **222** (1992) 1;
P. Forgács, A. Wipf, J. Balog, L. Fehér and L. O’Raifeartaigh, Phys. Lett. **B 227** (1989) 214;

- [16] M. Bauer, Ph. Di Francesco, C. Itzykson and J.-B. Zuber, Nucl. Phys. **B 362** (1991) 515
- [17] M. Bauer and N. Sochen, Phys. Lett. **B 275** (1992) 82
- [18] G.G.A. Bäuerle and E.A. de Kerf, *Lie Algebra, Part 1* (Elsevier Science Publishers B.V., North Holland, 1990)
- [19] C. Becchi, A. Rouet and R. Stora, Phys. Lett. **B 52** (1975) 344;
C. Becchi, A. Rouet and R. Stora, Ann. Phys. **98** (1976) 287;
I.V. Tyutin, Lebedev Preprint FIAN 39 (1975)
- [20] A.A. Belavin, in *Proc. of the second Yukawa Symposium, Nishinomiya, Japan*, Springer Proceedings in Physics, Vol. 31 (1988) 132
- [21] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, Nucl. Phys. **B 241** (1984) 333;
A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, J. Stat. Phys. **34** (1984) 763
- [22] D. Bernard and G. Felder, Commun. Math. Phys. **127** (1990) 145
- [23] M.A. Bershadsky, V.G. Knizhnik and M.G. Teitelman, Phys. Lett. **B 151** (1985) 31
- [24] M. Bershadsky and H. Ooguri, Commun. Math. Phys. **126** (1989) 49
- [25] W. Boucher, D. Friedan and A. Kent, Phys. Lett. **B 172** (1986) 316
- [26] P. Bouwknegt, J. McCarthy and K. Pilch, Phys. Lett. **B 234** (1990) 297;
P. Bouwknegt, J. McCarthy and K. Pilch, Commun. Math. Phys. **131** (1990) 125;
P. Bouwknegt, J. McCarthy and K. Pilch, Prog. Theor. Phys. Suppl. **102** (1990) 67
- [27] P. Bouwknegt and K. Schoutens, Phys. Rep. **223** (1993) 183
- [28] P. Bouwknegt and K. Schoutens (Eds.), *W-symmetry*, Adv. Series in Math. Phys. **22** (World Scientific, 1995)
- [29] P. Bowcock and A. Taormina, DTP/96/21, hep-th/9605220, preprint;
P. Bowcock, R.-L.K. Koktava and A. Taormina, DTP/96/11, hep-th/9606015, preprint
- [30] E. Braaten, T.L. Curtright and C.K. Zachos, Nucl. Phys. **B 260** (1985) 630
- [31] J.L. Cardy, in *Les Houches, Session XLIX 1988, Fields, Strings and Critical Phenomena*, E. Brezin and J.Zinn-Justin (Eds.) (Elsevier, 1989)
- [32] V. Chari and A. Pressley, *A Guide to Quantum Groups* (Cambridge Univ. Press, 1995)
- [33] M. Caselle and G. Ponzano, Phys. Lett. **B 224** (1989) 303;
M. Caselle, G. Ponzano and F. Ravanini, Phys. Lett. **251** (1990) 260
- [34] P. Christe and R. Flume, Nucl. Phys. **B 282** (1987) 466
- [35] S. Coleman, Commun. Math. Phys. **31** (1973) 259

- [36] J. de Boer, *Extended Conformal Symmetry in Non-Critical String Theory* (Univ. of Utrecht, 1993) Ph.D. thesis
- [37] J. de Boer and L. Fehér, ITP-SB-96-22, INS-Rep.-1140, hep-th/9605102, preprint
- [38] R. Dijkgraaf and E. Verlinde, Nucl. Phys. **B** (Proc. Suppl.) **5B** (1988) 87
- [39] P. Di Vecchia, B. Durhuus and J.L. Petersen, Phys. Lett. **B** **144** (1984) 245
- [40] P. Di Vecchia, J.L. Petersen and H.B. Zheng, Phys. Lett. **B** **162** (1985) 327;
P. Di Vecchia, J.L. Petersen and M. Yu, Phys. Lett. **B** **172** (1986) 211
- [41] V.I.S. Dotsenko, Nucl. Phys. **B** **338** (1990) 747;
V.I.S. Dotsenko, Nucl. Phys. **B** **358** (1991) 547
- [42] V.I.S. Dotsenko and V.A. Fateev, Nucl. Phys. **B** **240**[FS12] (1984) 312;
V.I.S. Dotsenko and V.A. Fateev, Nucl. Phys. **B** **251**[FS13] (1985) 691
- [43] V.I.S. Dotsenko and V.A. Fateev, Phys. Lett. **B** **154** (1985) 291
- [44] V.G. Drinfeld and V.V. Sokolov, J. Sov. Math. **30** (1984) 1975
- [45] T. Eguchi and A. Taormina, Phys. Lett. **B** **210** (1988) 125
- [46] H. Eichenherr, Phys. Lett. **B** **151** (1985) 26
- [47] V.A. Fateev and S.L. Lykyanov, Int. J. Mod. Phys. **A** **3** (1988) 507;
V.A. Fateev and S.L. Lykyanov, Sov. Phys. JETP **67** (1988) 447
- [48] V.A. Fateev and A.B. Zamolodchikov, Nucl. Phys. **B** **280**[FS18] (1987) 644
- [49] B.L. Feigin and E. Frenkel, Phys. Lett. **B** **246** (1990) 75
- [50] B.L. Feigin and E.V. Frenkel, Commun. Math. Phys. **128** (1990) 161;
B.L. Feigin and E.V. Frenkel, Lett. Math. Phys. **19** (1990) 307;
B.L. Feigin and E.V. Frenkel, in *Physics and Mathematics of Strings*, L. Brink *et al* (Eds.) (World Scientific 1990)
- [51] B.L. Feigin and D.B. Fuks, Funct. Anal. Appl. **16** (1982) 114;
B.L. Feigin and D.B. Fuks, Funct. Anal. Appl. **17** (1983) 241
- [52] B.L. Feigin and D.B. Fuks, Springer Lecture Notes in Mathematics 1060 (1984) 230
- [53] B.L. Feigin and D.B. Fuks, in *Representations of Lie Groups and Related Topics*, A. Vershik and D. Zhelobenko (Eds.) (Gordon and Breach, 1990)
- [54] B. Feigin and F. Malikov, Adv. Sov. Math. **17** (1993) 15, hep-th/9306137;
B. Feigin and F. Malikov, q-alg/9511011, preprint;
K. Iohara and F. Malikov, Mod. Phys. Lett. **A** **8** (1993) 3613
- [55] B. Feigin and F. Malikov, Lett. Math. Phys **31** (1994) 315

- [56] G. Felder, Nucl. Phys. **B 317** (1989) 215 [Erratum: **B 324** (1989) 548]
- [57] G. Felder and A. Varchenko, hep-th/9502165, preprint;
G. Felder and C. Wierczkowski, hep-th/9411004, preprint
- [58] P. Fre' and P. Soriani, *The N=2 Wonderland: From Calabi-Yau Manifolds to Topological Field Theories* (World Scientific, 1995)
- [59] D. Friedan, in *Les Houches, Session XXXIX 1982, Recent Advances in Field Theory and Statistical Mechanics*, J.B. Zuber and R. Stora (Eds.) (Elsevier, 1984)
- [60] D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. **B 271** (1986) 93
- [61] D. Friedan, Z. Qui and S. Shenker, in *Vertex Operators in Mathematics and Physics*, J. Lepowski, S. Mandelstam and I.M. Singer (Eds.) (Berkeley, 1983);
D. Friedan, Z. Qui and S. Shenker, Phys. Rev. Lett. **52** (1984) 1575;
D. Friedan, Z. Qui and S. Shenker, Commun. Math. Phys. **107** (1986) 535
- [62] D. Friedan, Z. Qui and S. Shenker, Phys. Lett. **B 151** (1985) 37
- [63] J. Fuchs, *Affine Lie Algebras and Quantum Groups* (Cambridge Univ. Press, 1992)
- [64] P. Furlan, A.Ch. Ganchev, R. Paunov and V.B. Petkova, Phys. Lett. **B 267** (1991) 63;
P. Furlan, A.Ch. Ganchev, R. Paunov and V.B. Petkova, Nucl. Phys. **B 394** (1993) 665;
A.Ch. Ganchev and V.B. Petkova, Phys. Lett. **B 293** (1992) 56
- [65] P. Furlan, A.Ch. Ganchev and V.B. Petkova, ESI-362-96, hep-th/9608018, preprint
- [66] A.Ch. Ganchev and V.B. Petkova, Phys. Lett. **B318** (1993) 74
- [67] K. Gawedzki, in *New Symmetry Principles in Quantum Field Theory*, J. Fröhlich *et al* (Eds.), proceedings of NATO Advanced Study Institute, Cargese 1991 (Plenum Press, 1992)
- [68] K. Gawedzki and A. Kupiainen, Nucl. Phys. **B 320** (1989) 625
- [69] D. Gepner and E. Witten, Nucl. Phys. **B 278** (1986) 493
- [70] A. Gerasimov, A. Marshakov, A. Morozov, M. Olshanetskii, and S. Shatashvili, Int. J. Mod. Phys. **A 5** (1990) 2495
- [71] P. Ginsparg, in *Les Houches, Session XLIX 1988, Fields, Strings and Critical Phenomena*, E. Brezin and J.Zinn-Justin (Eds.) (Elsevier, 1989)
- [72] P. Ginsparg and G. Moore, YCTP-P23-92, LA-UR-92-3479, hep-th/9304011, lecture notes from the 1992 TASI Summer School
- [73] P. Goddard and D. Olive, Int. J. Mod. Phys. **A 1** (1986) 303
- [74] P. Goddard, A. Kent and D. Olive, Phys. Lett. **B 152** (1985) 88;
P. Goddard, A. Kent and D. Olive, Commun. Math. Phys. **103** (1986) 105

- [75] C. Gomez and G. Sierra, Phys. Lett. **B 240** (1990) 149;
C. Gomez and G. Sierra, Nucl. Phys. **B 352** (1991) 791
- [76] C. Gomez and G. Sierra, in *Proc. of the Trieste 1990 Summer School* (World Scientific, 1991);
C. Gomez, M. Ruiz-Altaba and G. Sierra, *Quantum Groups in Two-dimensional Physics* (Cambridge Univ. Press, 1996)
- [77] M.B. Green, J.H. Schwarz and E. Witten, *Superstring Theory* Vols. 1 and 2 (Cambridge Univ. Press, 1987)
- [78] M.B. Halpern, E. Kiritsis, N. Obers and K. Clubok, Phys. Rep. **265** (1996) 1
- [79] S. Helgason, in *Battelle Rencontres, 1967 Lectures in Mathematics and Physics*, Eds. C.M. DeWitt and J.A. Wheeler (W.A. Benjamin Inc., 1968)
- [80] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton Univ. Press, 1992)
- [81] H.-L. Hu and M. Yu, Phys. Lett. **B 289** (1992) 302;
H.-L. Hu and M. Yu, Nucl. Phys. **B 391** (1993) 389
- [82] K. Ito, Phys. Lett. **B 252** (1990) 69
- [83] C. Itzykson and J.-M. Drouffe, *Statistical Field Theory, Vol. 2* (Cambridge Univ. Press, 1989)
- [84] C. Itzykson, H. Saleur and J.-B. Zuber, Eds., *Conformal Invariance and Applications to Statistical Mechanics* (World Scientific, 1988)
- [85] V.G. Kac, Lecture Notes in Phys. **94** (1979)
- [86] V.G. Kac, *Infinite Dimensional Lie Algebras*, (3rd edition, Cambridge Univ. Press, 1990)
- [87] V.G. Kac and D.A. Kazhdan, Adv. Math. **34** (1979) 79
- [88] V.G. Kac and A.K. Raina, *Bombay Lectures on Highest Weight Representations of Infinite Dimensional Lie Algebras*, Advanced Series in Mathematical Physics Vol. 2 (World Scientific, 1987)
- [89] V.G. Kac and M. Wakimoto, Proc. Natl. Acad. Sci. USA **85** (1988) 4956;
V.G. Kac and D.A. Kazhdan, Adv. Ser. Math. Phys., Vol. 7 (World Scientific, 1989), p. 138
- [90] D. Karabali, Q-H. Park, H.J. Schnitzer and Z. Yang, Phys. Lett. **B 216** (1989) 307
- [91] D. Karabali and H.J. Schnitzer, Nucl. Phys. **B 329** (1990) 649
- [92] C. Kassel, *Quantum Groups*, GTM 155 (Springer, 1995)
- [93] M. Kato and S. Matsuda, Phys. Lett. **B 172** (1986) 216

- [94] M. Kato and Y. Yamada, Prog. Theor. Phys. Suppl. **110** (1992) 291
- [95] S.V. Ketov, *Conformal Field Theory* (World Scientific, 1995)
- [96] V. Kiryakova, *Generalized Fractional Calculus and Applications*, Pitman Research Notes in Mathematics Series 301 (Longman Group, UK, 1994)
- [97] V.G. Knizhnik and A.B. Zamolodchikov, Nucl. Phys. **B 247** (1984) 83
- [98] T. Kugo and I Ojima, Prog. Theor. Phys. Suppl. **66** (1979) 1
- [99] G. Kuroki, Commun. Math. Phys. **142** (1991) 511
- [100] S. Majid, *Foundations of Quantum Group Theory* (Cambridge Univ. Press, 1995)
- [101] F.G. Malikov, B.L. Feigin and D.B. Fuks, Funkt. Anal. Prilozhen **20** (1986) 25
- [102] A.C. McBride and G.F. Roach (Eds.), *Fractional Calculus* (Pitman Advanced Publishing Program, Boston, 1985)
- [103] R.V. Moody and A. Pianzola, *Lie Algebras with Triangular Decompositions* (John Wiley and Sons, 1995)
- [104] G. Moore and N. Reshetikhin, Nucl. Phys. **B 328** (1989) 557
- [105] G. Moore and N. Seiberg, Phys. Lett. **B 212** (1988) 451;
G. Moore and N. Seiberg, Nucl. Phys. **B 313** (1989) 16;
G. Moore and N. Seiberg, Commun. Math. Phys. **123** (1989) 177
- [106] A. Neveu and J. Schwarz, Nucl. Phys. **B 31** (1971) 86
- [107] S.P. Novikov, Sov. Math. Dokl **24** (1981) 222;
S.P. Novikov, Usp. Math. Nauk. **37** (1982) 3
- [108] J.L. Petersen, *Lecture Notes on BRST Quantisation* from Nordita Summer School in High Energy Physics, Gräftvallen, August 1986, unpublished
- [109] J.L. Petersen, *Notes on Conformal Field Theory*, lecture notes prepared with the help of J. Rasmussen (NBI, University of Copenhagen, 1994)
- [110] J.L. Petersen, J. Rasmussen and M. Yu, Nucl. Phys. **B 457** (1995) 309;
J.L. Petersen, J. Rasmussen and M. Yu, in *Proceedings of the workshop Gauge Theories, Applied Supersymmetry and Quantum Gravity, Leuven July 1995, Leuven Notes in Mathematical and Theoretical Physics, Vol. 6*, B. de Wit *et al* (Eds.) (Leuven 1996), hep-th/9510059
- [111] J.L. Petersen, J. Rasmussen and M. Yu, Nucl. Phys. **B 457** (1995) 343;
J.L. Petersen, J. Rasmussen and M. Yu, Nucl. Phys. **B** (Proc. Suppl.) **49** (1996) 27

- [112] J.L. Petersen, J. Rasmussen and M. Yu, NBI-HE-96-18, AS-ITP-96-13, hep-th/9607129, accepted for publication in Nucl. Phys. **B**;
J.L. Petersen, J. Rasmussen and M. Yu, to appear in *Proceedings on Inauguration Conference of Asia Pacific Center for Theoretical Physics, Seoul June 1996*
- [113] J.L. Petersen, J. Rasmussen and M. Yu, preprint in preparation
- [114] J. Polchinski, hep-th/9511157, talk given at YKIS '95;
J. Polchinski, NSF-ITP-96-60, hep-th/9607050
- [115] A.M. Polyakov, Sov. Phys. JETP Lett. **12** (1970) 538
- [116] A.M. Polyakov, Phys. Lett. **B 103** (1981) 207;
A.M. Polyakov, Phys. Lett. **B 103** (1981) 211
- [117] A.M. Polyakov, in *Physics and Mathematics of Strings*, Eds. L. Brink, D. Friedan and A.M. Polyakov (World Scientific, 1990)
- [118] A.M. Polyakov and P.B. Wiegman, Phys. Lett. **B 131** (1983) 121;
A.M. Polyakov and P.B. Wiegman, Phys. Lett. **B 141** (1984) 223
- [119] A. Pressley and G. Segal, *Loop Groups*, (Oxford Univ. Press, 1986)
- [120] C. Ramirez, H. Ruegg and M. Ruiz-Altaba, Phys. Lett. **B 247** (1990) 499;
C. Ramirez, H. Ruegg and M. Ruiz-Altaba, Nucl. Phys. **B 364** (1991) 195
- [121] P. Ramond, Phys. Rev. **D 3** (1971) 2415
- [122] J. Rasmussen, *W Algebra*, Master Thesis (in danish), NBI at the Univ. of Copenhagen (1993)
- [123] A. Rocha-Charidi, Proc. Symp. Pure. Math. **49** (1989) 259
- [124] B. Ross (Ed.), *Fractional Calculus and Its Applications*, Lecture Notes in Mathematics 457 (Springer 1975)
- [125] V. Sador, Int. Jour. Mod. Phys. **A 8** (1993) 5115
- [126] S.G. Samko, A.A. Kilbas and O.L. Marichev, *Fractional Integrals and Derivatives* (Gordon and Breach, 1993)
- [127] V.V. Schechtman and A.N. Varchenko, Invent. Math. **106** (1991) 139
- [128] A.N. Schellekens, *Conformal Field Theory*, lecture notes for *Grundlagen und neue Methoden der Theoretische Physik*, Saalburg, Germany, September 1995
- [129] A.N. Schellekens and N.P. Warner, Phys. Rev. **D 34** (1986) 3092
- [130] J. Schnittger, Nucl. Phys. **B 471** (1996) 521
- [131] J.H. Schwarz, CALT-68-2065, hep-th/9607201, lectures presented at the ICTP Spring School 1996 and at the TASI Summer School 1996

- [132] C. Sommerfeld, Phys. Rev. **176** (1968) 2019
- [133] H. Sugawara, Phys. Rev. **170** (1968) 1659
- [134] W. Taylor IV, LBL-34507, hep-th/9310040, Ph.D. thesis
- [135] C. Thorn, Nucl. Phys. **B 248** (1984) 551
- [136] C. Vafa, Phys. Lett. **B 206** (1988) 421
- [137] C. Vafa, Nucl. Phys. **B 469** (1996) 403
- [138] E. Verlinde, Nucl. Phys. **B 300** [FS22] (1988) 360
- [139] M. Wakimoto, Commun. Math. Phys. **104** (1986) 605
- [140] J. Wess and B. Zumino, Phys. Lett. **B 37** (1971) 95
- [141] E. Witten, Commun. Math. Phys. **92** (1984) 455
- [142] A.B. Zamolodchikov, Theor. Math. Phys. **63** (1985) 1205
- [143] A.B. Zamolodchikov and V.A. Fateev, Sov. J. Nucl. Phys. **43** (1986) 657
- [144] A.B. Zamolodchikov and V.A. Fateev, JETP **63** (1986) 913